

## A system of probabilistic inequalities

ten Raa, Thijs

*Published in:*  
SIAM Review

*Publication date:*  
1983

[Link to publication](#)

*Citation for published version (APA):*  
Ten Raa, M. H. (1983). A system of probabilistic inequalities. *SIAM Review*, 25(4), 569-570.

### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

### Take down policy

If you believe that this document breaches copyright, please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Problem 83-17, A System of Probabilistic Inequalities

Author(s): T. Ten Raa

Source: *SIAM Review*, Vol. 25, No. 4, (Oct., 1983), pp. 569-570

Published by: Society for Industrial and Applied Mathematics

Stable URL: <http://www.jstor.org/stable/2029472>

Accessed: 15/04/2008 08:41

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=siam>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

---

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We enable the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## PROBLEMS AND SOLUTIONS

EDITED BY MURRAY S. KLAMKIN

COLLABORATING EDITORS: HENRY E. FETTIS      CECIL C. ROUSSEAU  
 YUDELL L. LUKE                                      OTTO G. RUEHR

*All problems and solutions should be sent, typewritten in duplicate, to Murray S. Klamkin, Department of Mathematics, University of Alberta, Edmonton, Alberta, Canada T6G 2G1. An asterisk placed beside a problem number indicates that the problem was submitted without solution. Proposers and solvers whose solutions are published will receive 10 reprints of the corresponding problem section. Other solvers will receive just one reprint provided a self-addressed stamped (U.S.A. or Canada) envelope is enclosed. Proposers and solvers desiring acknowledgment of their contributions should include a self-addressed stamped postcard (no stamp necessary outside the U.S.A. and Canada). Solutions should be received by February 15, 1984.*

### PROBLEMS

#### Two Legendre Polynomial Identities

*Problem 83-16, by A. D. RAWLINS (Brunel University, Middlesex, UK).*

Show that


$$(1) \quad P_n \{x(x^2 - \alpha^2)^{-1/2}\} = \frac{(-1)^n}{n!} (x^2 - \alpha^2)^{(n+1)/2} \frac{d^n}{dx^n} (x^2 - \alpha^2)^{-1/2}, \quad n \geq 0,$$

$$(2) \quad P_n \left\{ \frac{1 + Bx}{B + x} \right\} = B^{-1} (n!)^{-2} \left\{ \frac{-(B + x)^{n+1}}{2} \right. \\ \left. \cdot \prod_{p=1}^n \left\{ -\frac{d}{dx} \left[ (1 - x^2) \frac{d}{dx} \right] - p(p - 1) \right\} \right\} \left\{ \frac{x - B}{x + B} \right\}, \quad n \geq 1.$$

#### A System of Probabilistic Inequalities

*Problem 83-17\*, by T. TEN RAA (Erasmus University, Rotterdam, The Netherlands).*

Consider  $n$  locations,  $n \geq 2$ :



Initially, there is *one* particle at each location. Then there are two consecutive transitions, governed by real nonnegative matrices  $(p_{ij})$  and  $(x_{ij})$ , respectively. Thus  $p_{ij}$  is the probability that the particle which is originally at location  $i$  will move to  $j$  in the first round. Similarly,  $x_{ij}$  is the probability that a particle which after the first round is at location  $i$  will move to  $j$  in the second round.

It is assumed that the system is *shaky*. More precisely, all particles move with certainty in both rounds. Furthermore, in the second transition any particle has a positive probability to reach any other location. Formally,

$$(1) \quad p_{ii} = 0, \quad p_{ij} > 0 \quad \text{for some } j \neq i, \quad i = 1, \dots, n,$$

$$(2) \quad x_{ii} = 0, \quad x_{ij} > 0 \quad \text{for all } j \neq i, \quad i = 1, \dots, n.$$

(It is not necessary to assume that transition probabilities add up to unity.)

Consider the particle which originates from any location  $i$ . The probability that it

ends up at location  $j$  is  $\sum_{k=1}^n p_{ik}x_{kj}$ . Now *suppose* that the system is so shaky that our particle is at least as likely to end up in any other location  $j$  as to return to its origin  $i$ :

$$(3) \quad \sum_{k=1}^n p_{ik}x_{kj} \geq \sum_{k=1}^n p_{ik}x_{ki} \quad \text{for all } j \neq 1, \quad i = 1, \dots, n.$$

It is *conjectured* that this is impossible: the real nonnegative system (1), (2) and (3) is *inconsistent*. In other words, some particle must be more likely to return than to end up at some other location.

This problem arose in an attempt to extend a generalization of Kakutani's fixed point theorem to certain unbounded regions.

### An Infinite System of Differential Equations

*Problem 83-18\**, by T. D. ROGERS (University of Alberta).

M. von Smoluchowski [1] obtained the differential system

$$\frac{dN_k}{dt} = \sum_{i+j=k} a_{ij}N_iN_j - N_k \sum_{j=1}^{\infty} a_{kj}N_j$$

in the analysis of the coagulation of colloidal particles as a consequence of Brownian motion. The physically implausible assumption that  $a_{ij} = \text{constant}$  enables one to derive a solution in closed form. Can anything be said about the form of the solutions in the more general case? Even the cases  $a_{ij} = a_i$  or  $a_{ij} = a_j$  would be of interest. This equation has had recent application as a model for cell aggregation kinetics, and related numerical work has been carried out by T. D. Rogers and J. R. Sampson [2], [3].

### REFERENCES

- [1] S. CHANDRASEKHAR, *Stochastic problems in physics and astronomy*, in Papers on Noise and Stochastic Processes, Nelson Wax, ed., Dover, New York, 1954, pp. 1-89.
- [2] T. D. ROGERS, *Local models of cell aggregation kinetics*, Bull. Math. Biol., 39 (1977), pp. 23-42.
- [3] T. D. ROGERS AND J. R. SAMPSON, *A random walk model of cellular kinetics*, Int. J. Biomed. Comp., 1, 8 (1977), pp. 45-60.

### Properties of an Operator

*Problem 83-19\**, by P. SCHWEITZER (University of Rochester).

Determine whether or not the operator  $T: E^{NR+} \rightarrow E^{NR+}$ ,

$$(Tx)_{ir} \equiv \frac{a_{ir} + \sum_{j=1}^N M(r)_{ij}x_{jr}}{\max \left[ 1, \sum_{k=1}^R b_{ik} \left[ a_{ik} + \sum_{l=1}^N M(k)_{il}x_{lk} \right] \right]}, \quad x_{ir} \geq 0, 1 \leq r \leq R, \quad 1 \leq i \leq N$$

(where  $a_{ir} \geq 0, b_{ir} > 0, M(r)_{ij} \geq 0, \text{Spr } M(r) < 1, \forall r, i, j$ ) (Spr = Spectral radius) is a contraction operator in some norm, or a  $n$ -step contraction operator for some  $n$ . (Empirically,  $T$  has a unique fixed point  $x^*$  and  $T^m x \rightarrow x^*$  geometrically for any  $x$ .)

*Remark.* for the special case  $R = 1, T$  is a contraction operator.  $T$  reduces to (suppressing the index  $r$  or  $k$ )

$$(Tx)_i = \min \left[ \frac{1}{b_i}, a_i + \sum_{j=1}^N M_{ij}x_j \right], \quad 1 \leq i \leq N$$