Tilburg University

Comparison of bias-reducing methods for estimating the parameter in dilution series

Strijbosch, L.W.G.; Does, R.J.M.M.

Publication date:
1988

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Strijbosch, L. W. G., & Does, R. J. M. M. (1988). *Comparison of bias-reducing methods for estimating the parameter in dilution series*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 304). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM
R

UNIVERSITY
JEKE
UNIVERSITEIT
BRABANT

7626
1988
304

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS

DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM

26

04

COMPARISON OF BIAS-REDUCING METHODS FOR
ESTIMATING THE PARAMETER IN DILUTION
SERIES

Leo W.G. Strijbosch

Ronald J.M.M. Does

R. 46
woc. 510.922

FEW 304

COMPARISON OF BIAS-REDUCING METHODS FOR ESTIMATING THE PARAMETER IN DILUTION SERIES

Leo W.G. Strijbosch

Ronald J.M.M. Does

Computer Applications Group, School of Economics and Business, Tilburg University, P.O.Box 90153, 5000 LE Tilburg, The Netherlands	Department of Medical Informatics and Statistics, University of Limburg, P.O.Box 616, 6200 MD Maastricht, The Netherlands
---	--

Key Words and Phrases : limiting and serial dilution assays, maximum likelihood, jackknife methods, bootstrap methods, minimum chi-square, Monte Carlo experiments, experimental design.

ABSTRACT

Ten different estimators of the parameter in a limiting or serial dilution assay are compared. Eight of them are constructed to reduce the bias of the commonly used maximum likelihood estimator. Extensive Monte Carlo experiments using various designs, suggest that a particular jackknife version of the maximum likelihood estimator is preferred, provided that the design is not too small.

1. INTRODUCTION

Limiting and Serial Dilution Assays (LDA and SDA) are widely used in many areas, including public hygiene, bacteriology, biology and immunology; see Taswell (1987). In general these assays are primarily intended to estimate the relative frequency of a well-defined cell in a population of cells or the average number of organisms per unit volume of solution. In both LDA and SDA this

parameter is commonly estimated by using the "single-hit Poisson model" with quantal data yielded by samples taken from different dilutions. The assumptions underlying this model are well-known (see Finney (1978), Taswell (1981)) and will be described briefly, using the terminology of LDA. A test preparation contains numerous cells of which an unknown proportion ϕ has a certain property, for example immuno-competency. From this test preparation, m different dilutions are prepared. Then, from dilution j , n_j replicate cultures are taken. The number of cells in the k -th replicate culture of dilution j is a Poisson distributed variable with mean x_{jk} . A fraction ϕ of these cells has the intended property. A further assumption is that a positive response is obtained for a replicate culture, if and only if at least one cell of the specific type is present.

Statisticians can contribute to the execution of a LDA or a SDA in at least two ways. They can help the experimenter to construct an experimental design which will take advantage of existing a priori information. This hopefully precludes experimentation yielding useless data, and it enables adjusting the precision of an estimator. Furthermore they can advise on the statistical techniques to be used. In many applications of dilution analysis, the assays are very expensive and time consuming, while in some circumstances they are not repeatable either. In these cases it is of vital interest to carefully choose an experimental design and a statistical estimator minimizing bias and standard error. Recent research has been done on design problems (Loyer (1981), Taswell (1987) and Strijbosch et al. (1987)), and Monte Carlo studies have been made on the choice of the statistical procedure to be used (Salama et al. (1978), Loyer (1981), Taswell (1981), Strijbosch et al. (1987) and Does et al. (1988)). The results of these Monte Carlo studies cannot be compared properly because of the absence of generally accepted design methods: most authors used different experimental designs, when generating the simulation results. It is obvious that the statistical properties of the possible estimators

are dependent on the design used. If these properties do not hold over other possible designs it could easily occur, that one author finds that estimator 1 is better than estimator 2 while another author finds conflicting results.

This paper is organized as follows. Section 2 describes the experimental design, which has been used to compare different estimators. Section 3 discusses ten different estimators for the parameter in dilution series. The last two Sections are devoted to the Monte Carlo experiments and the results, respectively.

2. EXPERIMENTAL DESIGN

It is a very important issue, when comparing statistical estimators, to use a design which can be considered as a reference and a frame. The design method proposed by Strijbosch et al. (1987) seems to be a good candidate for use in general dilution assays and in Monte Carlo comparisons. First some notation will be introduced. Let the number of groups of replicate cultures be denoted by m , the (mean) number of cells tested in a replicate culture of group j by x_j , and the number of replicate cultures for group j by n_j , $j=1, \dots, m$. Furthermore let φ denote the unknown frequency.

It is convenient to split the total design problem into two parts. Firstly, the design parameters m , and $x_1, \dots, x_m$ are determined; secondly, the numbers of replicate cultures $n_1, \dots, n_m$ are chosen such that the expected bias and the expected standard error of an estimator are within certain bounds. The efficacy of Strijbosch et al. (1987)'s method for determining the design parameters m and $x_1, \dots, x_m$ can be explained as follows. A researcher setting up a dilution assay, will in general have some prior information about the value of φ . It is natural to think of a lower bound φ_1 and an upper bound φ_2 for φ . He wants to plan his assay such that not too many fractions of negatively responding cultures are too close to either zero or one. There must be enough dilutions which yield fractions of negatively responding cultures, somewhere

in the middle between zero and one. This must be true for every value of φ which could be the true value according to the prior information of the experimenter, that is, for every φ satisfying $\varphi_1 \leq \varphi \leq \varphi_2$. We must be more specific in order to deduct design formulae from these general set-up considerations. If only fractions between certain values P_1 and P_2 are called sufficiently informative and if we want to have (on the average) d fractions between these values for each possible φ in the range $[\varphi_1, \varphi_2]$, then the design parameters m and $x_1, \dots, x_m$ can be chosen according the formulae (i) through (iv) in Strijbosch et al. (1987). The advantages of this design method are: it incorporates researcher's criteria, it has suitable properties, and it can be easily used in Monte Carlo experiments aimed at a meaningful comparison of statistical estimators.

The most interesting statistical estimators for which comparisons are made in the various studies mentioned before, will be compared in this study, namely the minimum chi-square method (MC), the maximum likelihood method (ML) (see Taswell (1981)), three jackknife versions (Jr, Jc, and Je) of the ML estimator (see Does et al. (1988)), two methods (S1 and S2) invented by Salama et al. (1978), and three bootstrap versions (Br, Bc, and Be) of the ML estimator. In the next Section these estimators will be described briefly.

3. STATISTICAL METHODS

3.1 Notation

In Section 2 we introduced the following notation: m equals the number of groups of replicate cultures, x_j equals the (mean) number of cells tested in a replicate culture of group j , n_j equals the number of replicate cultures for group j , $j=1, \dots, m$, and φ denotes the unknown frequency. Then the data of the biometrical model can be represented as follows :

$$\{Y_{jk}\} \quad j=1, \dots, m; \quad k=1, \dots, n_j, \quad (1)$$

where Y_{jk} are independent Bernoulli distributed variables, with $P(Y_{jk} = 0) = 1 - P(Y_{jk} = 1) = \exp(-\varphi x_j)$. A negative response for a replicate culture is thus denoted by zero.

3.2 The Maximum Likelihood Method

From (1) it follows that the logarithm of the likelihood function $L(\varphi)$ is given by

$$\log L(\varphi) = \sum_{j=1}^m \sum_{k=1}^{n_j} \{-(1-Y_{jk})\varphi x_j + Y_{jk} \log(1-\exp(-\varphi x_j))\}. \quad (2)$$

The ML estimator ($\hat{\varphi}_{ML}$) is the value of φ that maximizes (2). As is pointed out in Does et al. (1988), this estimator must be slightly adapted in order to obtain an estimator with finite bias. If all Y_{jk} equal 1 (this event occurs with a small but positive probability), then the ML estimate equals infinity and hence $E(\hat{\varphi}_{ML}) = \infty$, thus leading to an infinite, rather than an asymptotically negligible bias. Does et al. (1988) proposed the following modification: whenever $Y_{jk} = 1$ for $j=1, \dots, m$, $k=1, \dots, n_j$ replace one Y_{jk} , for example the most suitable such as Y_{11} , by 0. It is shown that this simple modification suffices to reduce the bias from infinity to the desired order. This adaptation of $\hat{\varphi}_{ML}$ will be assumed throughout the paper.

Sufficiency implies that the relevant observations from a LDA consist of the independent binomial random variables R_j defined by $R_j = \sum_{k=1}^{n_j} (1-Y_{jk})$. The vector $(R_1, \dots, R_m)$ will be denoted by $\underline{R}$. Furthermore let $n_j - R_j$ be denoted by Q_j and $(Q_1, \dots, Q_m)$ by $\underline{Q}$.

3.3 Three Jackknife Versions of the ML Estimator

In general the jackknife is defined as follows. Suppose a parameter Θ is estimated on the basis of the stochastic variables

$X_1, \dots, X_N$. Consider an estimate $T_N = T_N(X_1, \dots, X_N)$. Then the i -th jackknifed pseudo-value is defined as

$$T_{Ni}^J = NT_N - (N-1)T_{N-1}(X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_N), \quad i=1, \dots, N. \quad (3)$$

The jackknife estimator T_N^J defined by

$$T_N^J = N^{-1} \sum_{i=1}^N T_{Ni}^J \quad (4)$$

reduces the bias in many cases (see Efron (1982)). In our case, the pseudo-values in (3) can be obtained in three different ways. When $n_j = n$ the biometrical model (1) is a matrix with columns that are independent, identically distributed (iid) random vectors. As jackknife estimates are in general determined from iid variables, the natural way to jackknife is to drop one column from (1) at a time (see Strijbosch et al. (1987), and Kleijnen et al. (1987)); this yields the jackknife estimate $\hat{\varphi}_{Jc}$. The two other versions are non-iid cases and are obtained by deleting one row at a time (yielding $\hat{\varphi}_{Jr}$) or one element at a time (yielding $\hat{\varphi}_{Je}$), respectively (see Does et al. (1988)). Note that these last two estimates can be obtained from R .

3.4 Three Bootstrap Versions of the ML Estimator

Analogous to the three jackknife versions there are three bootstrap versions $\hat{\varphi}_{Bc}$, $\hat{\varphi}_{Br}$, and $\hat{\varphi}_{Be}$. Suppose again that $T_N = T_N(X_1, \dots, X_N)$ is an estimate of the parameter Θ . The ordinary bootstrap approach consists of drawing M random samples $(X_1^i, \dots, X_N^i)_{i=1, \dots, M}$ of size N from $X_1, \dots, X_N$ (see Efron (1982)). Such a sample is obtained by computer generated random sampling with replacement. Although there exist more elaborated bootstrap methods (see Davison et al. (1986)) the simple bootstrap will be used. The i -th bootstrapped pseudo-value is defined as

$$T_{Ni}^B = T_N(X_1^i, \dots, X_N^i), \quad i=1, \dots, M. \quad (5)$$

The bootstrap estimator T_N^B defined by

$$T_N^B = M^{-1} \sum_{i=1}^M T_{Ni}^B \quad (6)$$

could potentially reduce the bias when estimating the parameter in LDA. As discussed in the previous subsection (§3.3), there are three different ways to define a set $(X_1, \dots, X_N)$ from the biometrical model (1), thus resulting in the bootstrap estimators $\hat{\varphi}_{Bc}$, $\hat{\varphi}_{Br}$ and $\hat{\varphi}_{Be}$. As before the c-variant represents the iid-case, and is only applicable when all $n_j = n$, $j=1, \dots, m$.

3.5 Salama et al. (1978) Bias-Reducing Methods

Using Taylor expansions and implicit function theorems, Salama et al. (1978) showed the existence of functions $H(Q, \hat{\varphi}_{ML})$ such that the estimator $\hat{\varphi}_S$ defined by

$$\hat{\varphi}_S = \hat{\varphi}_{ML} - H(Q, \hat{\varphi}_{ML}) \quad (7)$$

satisfies

$$E(\hat{\varphi}_S) = \varphi + \sum_{j=1}^m O(n_j^{-2}), \quad (8)$$

thus removing the first order bias term. The following two alternatives for the function H are given (yielding the estimators $\hat{\varphi}_{S1}$ and $\hat{\varphi}_{S2}$):

$$G_1(Q, \hat{\varphi}_{ML}) = \frac{1}{2} \sum_{j=1}^m \left(\frac{\partial^2 \hat{\varphi}_{ML}}{\partial Q_j^2} \right) n_j e^{-\hat{\varphi}_{ML} x_j} (1 - e^{-\hat{\varphi}_{ML} x_j}) \quad (9)$$

and

$$G_2(Q, \hat{\varphi}_{ML}) = \frac{1}{2} \sum_{j=1}^m \left(\frac{\partial^2 \hat{\varphi}_{ML}}{\partial Q_j^2} \right) \underline{u}_j n_j e^{-\hat{\varphi}_{ML} x_j} (1 - e^{-\hat{\varphi}_{ML} x_j}). \quad (10)$$

The second order derivatives in the right-hand side of (10) are evaluated at $\underline{u}$, defined by $\underline{u} = E(Q)$. Unfortunately the formula

given by Salama et al. (1978) for $\frac{\partial^2 \hat{\varphi}_{ML}}{\partial Q_j^2}$ shows typing errors. The

correct formula is :

$$\frac{\partial^2 \hat{\varphi}_{ML}}{\partial Q_j^2} = \frac{F_j^2}{D^3} \left\{ \sum_{i=1}^m Q_i e^{-\hat{\varphi}_{ML} x_i} F_i^2(x_i + 2F_i e^{-\hat{\varphi}_{ML} x_i}) \right. \\ \left. - 2 \frac{F_j^3}{D^2} e^{-\hat{\varphi}_{ML} x_j} \right\}, \quad (11)$$

where F_i and D are defined by, respectively,

$$F_i = F_i(\hat{\varphi}_{ML}) = x_i / (1 - e^{-\hat{\varphi}_{ML} x_i}) \quad (12)$$

and

$$D = D(Q, \hat{\varphi}_{ML}) = \sum_{j=1}^m Q_j e^{-\hat{\varphi}_{ML} x_j} F_j^2. \quad (13)$$

Salama et al. (1978) used a special modification of the ML estimator, in case all Q_j equal n_j , $j=1, \dots, m$. That modification is essentially the same as described earlier in the subsection of the ML method (§3.2).

3.6 The Minimum Chi-Square Method

The Minimum Chi-Square or MC estimator ($\hat{\varphi}_{MC}$) is determined as the value of φ that minimizes

$$\chi^2(\varphi) = \sum_{j=1}^m \frac{(R_j - n_j \exp(-\varphi x_j))^2}{(n_j \exp(-\varphi x_j) (1 - \exp(-\varphi x_j)))}. \quad (14)$$

Since the expected value of the MC estimator is infinite, $\hat{\varphi}_{MC}$ has been adapted in the same manner as $\hat{\varphi}_{ML}$ has; see §3.2.

4. MONTE CARLO EXPERIMENTS

An extended version of the simulation program described in Does et al. (1988) has been used for our Monte Carlo experiments. The modification consisted of the addition of the three bootstrap estimators, the two methods of Salama et al. (1978), and the MC

estimator. Much care has been taken to test the program. For example, the results of Salama et al. (1978) which have been based on the exact distributions of $\hat{\varphi}_{S1}$ and $\hat{\varphi}_{S2}$ could be reproduced quite satisfactorily in the simulations. In this paper the statistical estimators are compared for two different designs with three and two different values, respectively, for the number of replicates (n). Thus, using the design method described in Section 2, the following values of $\varphi_1, \varphi_2, P_1, P_2$ and d have been chosen : $\varphi_1=0.001$, $\varphi_2=0.01$, $P_1=0.15$, $P_2=0.70$, $d=2$ (yielding $m=4$) and $d=3$ (yielding $m=7$). For the case $d=2$ the simulation program has been executed with $n=6, 12$ and 18 and in the case $d=3$, with $n=6$ and 12 .

Simulation results have been obtained for 19 equidistant values of φ within the interval $[\varphi_1, \varphi_2]$. The number of generated samples for each combination of $\varphi_1, \varphi_2, P_1, P_2, d, n$ and φ was 1,000. The number of bootstrap samples (M) was 100.

The simulation program has been written in PASCAL and uses the NAG (Fortran) subroutines G05DZF and G05DYF for the generation of the Bernoulli variables $\{Y_{jk}\}$ and the bootstrap samples, respectively. The structure of the program will be described briefly for the case $d=2$. For each of the 19 values of φ , a matrix $\{y_{jk}\}$, $j=1, \dots, 4$; $k=1, \dots, 18$ is generated 1,000 times. These numbers are used twice : one time for $n=18$ and one time for either $n=6$ or $n=12$. This concession has been made in order to curtail the required CPU-time. Depending on the value of n , the numbers r_j are determined by $r_j = n - \frac{1}{K} \sum y_{jk}$, where $k=1, \dots, 6$ for $n=6$, $k=7, \dots, 18$ for $n=12$ and $k=1, \dots, 18$ for $n=18$. Thus 1,000 datasets (x_j, r_j, n) , $j=1, \dots, 4$ result for each combination of φ and n . For each dataset the program calculates, if possible, the weighted-mean estimate φ_0 (see Taswell (1981)). When all $r_j=0$ or n , this estimate cannot be determined. In that case $\varphi_0=(\varphi_1+\varphi_2)/2$ has been taken. φ_0 served as an initial estimate for the iterative determination of $\hat{\varphi}_{ML}$. $\hat{\varphi}_{ML}$ served as an initial estimate for the determination of $\hat{\varphi}_{MC}$, the jackknife, and the bootstrap pseudo-estimates. Comparison of the

statistical estimators has been based on the mean relative bias (MRB) and the coefficient of variation (CV), defined as follows :

$$\text{MRB} = \varphi^{-1} \sum_{t=1}^{1,000} (\hat{\varphi}_t - \varphi)/1,000 \quad (15)$$

and

$$\text{CV} = \varphi^{-1} \left\{ \sum_{t=1}^{1,000} (\hat{\varphi}_t - \varphi)^2/1,000 \right\}^{1/2}. \quad (16)$$

The simulations used 59 hours of CPU-time on a VAX 8700 computer.

5. RESULTS AND CONCLUSIONS

The uncorrected ML estimator exhibits a positive bias. Theoretically this is due to the fact that there is a positive probability that all dilutions will produce growth so that the ML estimate becomes infinite. In practice, when m and $n_j, j=1, \dots, m$ are large enough, this situation will not occur, and Monte Carlo simulations reveal the positive bias in that case. Resampling plans such as the jackknife and the bootstrap are designed to reduce the bias (see Section 3). The Monte Carlo results in Strijbosch et al. (1987) showed that jackknifing in LDA reduces both the bias and the mean squared error of the estimator. Does et al. (1988) explained that there are three different ways of constructing a jackknife estimator in dilution analysis. It is made plausible that all three jackknife estimators reduce the bias by eliminating the first order bias term, and Monte Carlo simulations showed the relative properties of these estimators with respect to the ML estimator for various designs. The three bootstrap versions of the ML estimator can be obtained in a similar way. The two bias-reducing methods in Salama et al. (1978) eliminate the first-order bias term in a way different from jackknifing and bootstrapping, and therefore it is interesting to compare these estimators. From former results (see Strijbosch et al. (1987) and Fazekas de St. Groth (1982)) it is clear that the MC estimator is not attractive. Nevertheless this

method is also considered in the Monte Carlo comparisons, especially because of the curious and inconsistent behaviour of its bias and mean squared error.

In the Figures 1a through 5b not all methods are included, in order to prevent confusion resulting from too many curves. For each category only the "best" one is shown. Does et al. (1988) concluded that, among the three jackknife estimators, $\hat{\varphi}_{Je}$ is the best. The present results confirm this conclusion. Thus $\hat{\varphi}_{Jr}$ and $\hat{\varphi}_{Jc}$ are not shown in the Figures. Comparing the simulation results for the three bootstrap estimates, $\hat{\varphi}_{Br}$ is obviously the best estimator. In general, the results tend to be such, that $MRB(\hat{\varphi}_{Be}) \sim 2 * MRB(\hat{\varphi}_{ML})$ and $MRB(\hat{\varphi}_{Br}) < MRB(\hat{\varphi}_{Bc}) \leq MRB(\hat{\varphi}_{Be})$. The coefficients of variation for the three bootstrap estimators are more comparable. Thus $\hat{\varphi}_{Bc}$ and $\hat{\varphi}_{Be}$ are not presented in the Figures. The estimates of Salama et al. (1978) have nearly equal MRB and CV. However, there might be a slight preference for $\hat{\varphi}_{S2}$. Thus $\hat{\varphi}_{S1}$ is not included in the Figures. When comparing the remaining five estimators in the Figures 1a through 5b, it becomes clear that in general the estimators $\hat{\varphi}_{Je}$ and $\hat{\varphi}_{S2}$ should be preferred. In small designs, however, the jackknife estimator has the undesired property of a strongly increasing CV for values of φ near φ_2 which can be explained by the frequent occurrence of the situation that all Y_{jk} equal 1 when calculating the pseudo-estimates (see Figure 1b). An attractive property of the jackknife is that it also yields the variance estimate used to determine proper confidence bounds for φ . A major disadvantage of the estimating methods of Salama et al. (1978) is the lack of an estimator for the variance.

Provided that an experimenter works with designs, which are not too small (the design of Figure 4 seems to be large enough), it is clear that the jackknife version of the ML estimator - obtained by leaving out one element at a time - is the statistical procedure of choice.

ACKNOWLEDGMENTS

The authors thank Jack Kleijnen for helpful suggestions.

BIBLIOGRAPHY

- Davison, A.C., Hinkley, D.V. and Schechtman, E (1986). Efficient bootstrap simulation. *Biometrika* 73, 555-566.
- Does, R.J.M.M., Strijbosch, L.W.G. and Albers, W. (1988). Using jackknife methods for estimating the parameter in dilution series. *Medical Informatics and Statistics Report* 13, University of Limburg, Maastricht. Submitted for publication.
- Efron, B. (1982). The jackknife, the bootstrap and other resampling plans. Philadelphia : SIAM.
- Fazekas de St. Groth, S. (1982). The evaluation of limiting dilution assays. *Journal of Immunological Methods* 49, R11-R23.
- Finney, D.J. (1978). *Statistical Method in Biological Assay* . 3rd Edition, New York : Academic Press.
- Kleijnen, J.P.C., Karremans, P.C.A., Oortwijn, W.K. and Van Groenendaal, W.J.H. (1987). Jackknifing estimated weighted least squares: JEWS. *Commun. Statist.-Theor. Meth.* , 16(3), 747-764.
- Loyer, M.W. (1981). Using a serial dilution experiment to estimate the density of organisms. *Unpublished PH. D. Thesis, Montana State University*.
- Salama, I.A., Koch, G.G. and Tolley, H.D. (1978). On the estimation of the most probable number in a serial dilution assay. *Commun. Statist.-Theor. Meth.* , A7(13), 1267-1281.
- Strijbosch, L.W.G., Buurman, W.A., Does, R.J.M.M., Zinken, P.H. and Groenewegen, G. (1987). Limiting dilution assays : experimental design and statistical analysis. *Journal of Immunological Methods* 97, 133-140.
- Taswell, C. (1981). Limiting dilution assays for the determination of immunocompetent cell frequencies I : data analysis. *Journal of Immunology* 126 , 1614-1619.
- Taswell, C. (1987). Limiting dilution assays for the separation, characterization, and quantitation of biologically active particles and their clonal progeny. In *Cell Separation : Methods*

and Selected Applications 4, Pretlow, T.G. and Pretlow, T.P.
(eds), 109-145, New York : Academic Press.

Figure 1a

 $\varphi^1 = .001$ $\varphi^2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 2$ $m = 4$ $n = 6$

Figure 1b

 $\varphi^1 = .001$ $\varphi^2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 2$ $m = 4$ $n = 6$

Figure 2a

 $\varphi_1 = .001$ $\varphi_2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 2$ $m = 4$ $n = 12$

Figure 2b

 $\varphi_1 = .001$ $\varphi_2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 2$ $m = 4$ $n = 12$

Figure 3a

 $\phi^1 = .001$ $\phi^2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 2$ $m = 4$ $n = 18$

Figure 3b

 $\phi^1 = .001$ $\phi^2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 2$ $m = 4$ $n = 18$

Figure 4a

 $\varphi_1 = .001$ $\varphi_2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 3$ $m = 7$ $n = 6$

Figure 4b

 $\varphi_1 = .001$ $\varphi_2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 3$ $m = 7$ $n = 6$

Figure 5a

 $\varphi_1 = .001$ $\varphi_2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 3$ $m = 7$ $n = 12$

Figure 5b

 $\varphi_1 = .001$ $\varphi_2 = .01$ $p_1 = .15$ $p_2 = .70$ $d = 3$ $m = 7$ $n = 12$

IN 1987 REEDS VERSCHENEN

- 242 Gerard van den Berg
Nonstationarity in job search theory
- 243 Annie Cuyt, Brigitte Verdonk
Block-tridiagonal linear systems and branched continued fractions
- 244 J.C. de Vos, W. Vervaat
Local Times of Bernoulli Walk
- 245 Arie Kapteyn, Peter Kooreman, Rob Willemse
Some methodological issues in the implementation
of subjective poverty definitions
- 246 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel
Sampling for Quality Inspection and Correction: AOQL Performance
Criteria
- 247 D.B.J. Schouten
Algemene theorie van de internationale conjuncturele en structurele
afhankelijkheden
- 248 F.C. Bussemaker, W.H. Haemers, J.J. Seidel, E. Spence
On (v,k,λ) graphs and designs with trivial automorphism group
- 249 Peter M. Kort
The Influence of a Stochastic Environment on the Firm's Optimal Dyna-
mic Investment Policy
- 250 R.H.J.M. Gradus
Preliminary version
The reaction of the firm on governmental policy: a game-theoretical
approach
- 251 J.G. de Gooijer, R.M.J. Heuts
Higher order moments of bilinear time series processes with symmetri-
cally distributed errors
- 252 P.H. Stevers, P.A.M. Versteijne
Evaluatie van marketing-activiteiten
- 253 H.P.A. Mulders, A.J. van Reeken
DATAAL - een hulpmiddel voor onderhoud van gegevensverzamelingen
- 254 P. Kooreman, A. Kapteyn
On the identifiability of household production functions with joint
products: A comment
- 255 B. van Riel
Was er een profit-squeeze in de Nederlandse industrie?
- 256 R.P. Gilles
Economies with coalitional structures and core-like equilibrium con-
cepts

- 257 P.H.M. Ruys, G. van der Laan
Computation of an industrial equilibrium
- 258 W.H. Haemers, A.E. Brouwer
Association schemes
- 259 G.J.M. van den Boom
Some modifications and applications of Rubinstein's perfect equilibrium model of bargaining
- 260 A.W.A. Boot, A.V. Thakor, G.F. Udell
Competition, Risk Neutrality and Loan Commitments
- 261 A.W.A. Boot, A.V. Thakor, G.F. Udell
Collateral and Borrower Risk
- 262 A. Kapteyn, I. Woittiez
Preference Interdependence and Habit Formation in Family Labor Supply
- 263 B. Bettonvil
A formal description of discrete event dynamic systems including perturbation analysis
- 264 Sylvester C.W. Eijffinger
A monthly model for the monetary policy in the Netherlands
- 265 F. van der Ploeg, A.J. de Zeeuw
Conflict over arms accumulation in market and command economies
- 266 F. van der Ploeg, A.J. de Zeeuw
Perfect equilibrium in a model of competitive arms accumulation
- 267 Aart de Zeeuw
Inflation and reputation: comment
- 268 A.J. de Zeeuw, F. van der Ploeg
Difference games and policy evaluation: a conceptual framework
- 269 Frederick van der Ploeg
Rationing in open economy and dynamic macroeconomics: a survey
- 270 G. van der Laan and A.J.J. Talman
Computing economic equilibria by variable dimension algorithms: state of the art
- 271 C.A.J.M. Dirven and A.J.J. Talman
A simplicial algorithm for finding equilibria in economies with linear production technologies
- 272 Th.E. Nijman and F.C. Palm
Consistent estimation of regression models with incompletely observed exogenous variables
- 273 Th.E. Nijman and F.C. Palm
Predictive accuracy gain from disaggregate sampling in arima - models

- 274 Raymond H.J.M. Gradus
The net present value of governmental policy: a possible way to find the Stackelberg solutions
- 275 Jack P.C. Kleijnen
A DSS for production planning: a case study including simulation and optimization
- 276 A.M.H. Gerards
A short proof of Tutte's characterization of totally unimodular matrices
- 277 Th. van de Klundert and F. van der Ploeg
Wage rigidity and capital mobility in an optimizing model of a small open economy
- 278 Peter M. Kort
The net present value in dynamic models of the firm
- 279 Th. van de Klundert
A Macroeconomic Two-Country Model with Price-Discriminating Monopolists
- 280 Arnoud Boot and Anjan V. Thakor
Dynamic equilibrium in a competitive credit market: intertemporal contracting as insurance against rationing
- 281 Arnoud Boot and Anjan V. Thakor
Appendix: "Dynamic equilibrium in a competitive credit market: intertemporal contracting as insurance against rationing"
- 282 Arnoud Boot, Anjan V. Thakor and Gregory F. Udell
Credible commitments, contract enforcement problems and banks: intermediation as credibility assurance
- 283 Eduard Ponds
Wage bargaining and business cycles a Goodwin-Nash model
- 284 Prof.Dr. hab. Stefan Mynarski
The mechanism of restoring equilibrium and stability in polish market
- 285 P. Meulendijks
An exercise in welfare economics (II)
- 286 S. Jørgensen, P.M. Kort, G.J.C.Th. van Schijndel
Optimal investment, financing and dividends: a Stackelberg differential game
- 287 E. Nijssen, W. Reijnders
Privatisering en commercialisering; een oriëntatie ten aanzien van verzelfstandiging
- 288 C.B. Mulder
Inefficiency of automatically linking unemployment benefits to private sector wage rates

- 289 M.H.C. Paardekooper
A Quadratically convergent parallel Jacobi process for almost diagonal matrices with distinct eigenvalues
- 290 Pieter H.M. Ruys
Industries with private and public enterprises
- 291 J.J.A. Moors & J.C. van Houwelingen
Estimation of linear models with inequality restrictions
- 292 Arthur van Soest, Peter Kooreman
Vakantiebestemming en -bestedingen
- 293 Rob Alessie, Raymond Gradus, Bertrand Melenberg
The problem of not observing small expenditures in a consumer expenditure survey
- 294 F. Boekema, L. Oerlemans, A.J. Hendriks
Kansrijkheid en economische potentie: Top-down en bottom-up analyses
- 295 Rob Alessie, Bertrand Melenberg, Guglielmo Weber
Consumption, Leisure and Earnings-Related Liquidity Constraints: A Note
- 296 Arthur van Soest, Peter Kooreman
Estimation of the indirect translog demand system with binding non-negativity constraints

IN 1988 REEDS VERSCHENEN

- 297 Bert Bettonvil
Factor screening by sequential bifurcation
- 298 Robert P. Gilles
On perfect competition in an economy with a coalitional structure
- 299 Willem Selen, Ruud M. Heuts
Capacitated Lot-Size Production Planning in Process Industry
- 300 J. Kriens, J.Th. van Lieshout
Notes on the Markowitz portfolio selection method
- 301 Bert Bettonvil, Jack P.C. Kleijnen
Measurement scales and resolution IV designs: a note
- 302 Theo Nijman, Marno Verbeek
Estimation of time dependent parameters in linear models
using cross sections, panels or both
- 303 Raymond H.J.M. Gradus
A differential game between government and firms: a non-cooperative
approach

Bibliotheek K. U. Brabant

17 000 01065945 7