Tilburg University

Cost allocation and communication

van den Nouweland, C.G.A.M.; Borm, P.E.M.; Owen, G.; Tijs, S.H.

Publication date:
1991

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

van den Nouweland, C. G. A. M., Borm, P. E. M., Owen, G., & Tijs, S. H. (1991). *Cost allocation and communication*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 469). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CRM

CBM
R

7626
1991
469

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS

DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM

COST ALLOCATION AND COMMUNICATION

Anne van den Nouweland

Peter Borm

Guillermo Owen

Stef Tijs

FEW 469

Handwritten notes:
R.O.B.
5/17
1988

COST ALLOCATION AND COMMUNICATION

Anne van den Nouweland¹, Peter Borm², Guillermo Owen³ and Stef Tijs¹.

Abstract. This paper considers a special class of cost allocation problems, where the communication possibilities among the agents are restricted. Integral formulas are derived for two allocation rules: the Myerson value and the position value.

Key words. Cost allocation, communication, Myerson value, position value.

¹ Department of Mathematics, University of Nijmegen, Toernooiveld 1, 6525 ED Nijmegen, The Netherlands

² Department of Econometrics, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands

³ Department of Mathematics, Naval Postgraduate School, Monterey, CA 93943, USA. Financial support of the CentER for Economic Research of the Department of Economics, Tilburg University, The Netherlands, is gratefully acknowledged.

1. INTRODUCTION

Consider a group of economic agents who all want to make use of some machines or facilities. If the agents cooperate they can possibly share some facilities and thus save costs. In section 2 it is shown that this kind of cost allocation problems give rise to concave cost games which lie in the cone generated by all dual unanimity games.

Subsequently, we investigate the consequences of a restriction in communication possibilities between the agents on this kind of cost allocation problems. Throughout this paper we assume that the communication possibilities can be modelled by means of a (communication) graph. Based on the Shapley value, two solution concepts for communication situations were introduced: the Myerson value and the position value. An axiomatic characterization of the Myerson value was given by *Myerson* (1977) and *Borm et al.* (1990) gave an axiomatic characterization of the position value in case of cycle-free communication graphs. *Van den Nouweland* and *Borm* (1990) proved that if a communication graph is cycle-complete (cycle-free) and the underlying cost game is concave, then the Myerson value (position value) is in the core of the corresponding graph-restricted game.

Owen (1986) and *Borm et al.* (1990) provided integral formulas to compute the Myerson value and the position value in situations where the communication graph is cycle-free and the underlying game is a quadratic measure game. In section 3 we derive integral formulas for the Myerson value and the position value in situations where the underlying cost game is a dual unanimity game.

2. THE MODEL

Let $N := \{1, \dots, n\}$ and $2^N := \{S \mid S \subseteq N\}$. By TU^N we denote the class of all *transferable utility games* (N, v) with player set N and *characteristic function* $v : 2^N \rightarrow \mathbb{R}$ with $v(\emptyset) = 0$. There are basically two ways to interpret a TU -game (N, v) , the amount $v(S)$ can represent the revenue/gains for a *coalition* $S \subseteq N$ or it can represent the costs for this coalition. We prefer to denote a TU -game by (N, c) if it is to be interpreted as a cost game.

In this paper we consider cost games that are generated by cost allocation problems of the form $\langle N, F, p, d \rangle$. Here, N is the set of players, F is a finite set of facilities, $p : F \rightarrow \mathbb{R}_+$ is a function that assigns to every facility its non-negative price, and $d : N \rightarrow 2^F$ is a function that assigns to every player the subset of facilities demanded by this player.

Let $\langle N, F, p, d \rangle$ be such a cost allocation problem. Then the players in a coalition $S \subseteq N$ have to purchase each facility that at least one of them demands. On the other hand,

if two (or more) players demand the same facility, then they only have to purchase it once. Hence, this allocation problem leads to a cost game (N, c) with

$$c(S) := \sum_{r \in F} p(r) u_{N_r}^*(S),$$

where, for all $r \in F$, $N_r := \{i \in N \mid r \in d(i)\}$ is the set of players who demand facility r , and $u_{N_r}^*$ is the *dual unanimity game* on N_r , defined by $u_{N_r}^*(S) = \begin{cases} 1 & \text{if } S \cap N_r \neq \emptyset \\ 0 & \text{else} \end{cases}$. The proof of theorem 1 is straightforward and therefore omitted.

THEOREM 1. The class of all cost games corresponding to cost allocation problems of the form $\langle N, F, p, d \rangle$ is the convex cone generated by all dual unanimity games u_M^* with $M \in 2^N \setminus \{\emptyset\}$.

A cost game (N, c) is called *concave* if it is more advantageous to join a larger coalition or, in formula, if

$$c(S \cup \{i\}) - c(S) \geq c(T \cup \{i\}) - c(T)$$

for all $i \in N$ and all $S \subseteq T \subseteq N \setminus \{i\}$. The class of concave games with player set N is a convex cone and (N, u_M^*) is a concave game for every $M \subseteq N$. Therefore, a direct consequence of theorem 1 is that all cost games corresponding to cost allocation problems of the form $\langle N, F, p, d \rangle$ are concave.

It may be noted that the *Shapley value* $\Phi(N, c) \in \mathbb{R}^N$ (cf. *Shapley* (1953)) of the cost game (N, c) corresponding to the cost allocation problem $\langle N, F, p, d \rangle$ has a nice interpretation. As is easily seen the Shapley value of a dual unanimity game (N, u_M^*) equals 0 for $i \notin M$ and $\frac{1}{|M|}$ for $i \in M$. Since the Shapley value is linear, it follows that

$$\Phi_i(N, c) = \sum_{r \in d(i)} \frac{p(r)}{|N_r|}$$

for all $i \in N$, which implies that the costs of each facility are equally divided among the players that make use of it.

3. INTEGRAL FORMULAS FOR THE MYERSON VALUE AND THE POSITION VALUE

So far, we implicitly assumed that all players can freely communicate with one another. Now suppose that communication between the players is restricted and that the communication possibilities are determined by an undirected (*communication*) graph

(N, A) in which the points are the players and the arcs correspond to pairs of players who can communicate directly. A triple (N, c, A) , where (N, c) is a cost game and (N, A) is a communication graph, is called a *communication situation*.

Let (N, c, A) be a communication situation. Then the players in a coalition $S \subseteq N$ can effect communication through all communication links of

$$A(S) := \{\{i, j\} \in A \mid \{i, j\} \subseteq S\}.$$

Hence a coalition S splits up into (*communication*) *components* in the following way: $T \subseteq S$ is a component within S if and only if the graph $(T, A(T))$ is connected and there is no set $\bar{T}$ such that $T \subsetneq \bar{T} \subseteq S$ and $(\bar{T}, A(\bar{T}))$ is connected. We denote the resulting partition of S by S/A .

Taking into account these communication restrictions, the costs $c_A(S)$ for a coalition $S \in 2^N$ can be defined as

$$c_A(S) := \sum_{T \in S/A} c(T).$$

(N, c_A) is called the *graph-restricted game*.

One can also focus on the communication links. The cost savings for the grand coalition induced by the presence of the communication links in $L \subseteq A$ are defined as

$$r_N^c(L) := \sum_{i \in N} c(\{i\}) - \sum_{T \in N/L} c(T).$$

(A, r_N^c) is called the *arc (cost savings) game*.

Now we are ready to formulate the definitions of the Myerson value and the position value.

The *Myerson value* $\mu(N, c, A) \in \mathbb{R}^N$ (cf. *Myerson (1977)*) is defined as the Shapley value of the corresponding graph-restricted game, i.e.

$$\mu(N, c, A) := \Phi(N, c_A).$$

The *position value* of a communication situation (cf. *Borm et al. (1990)*) is based upon the Shapley value of the corresponding arc game: the corresponding cost savings of each arc are equally divided among the players it connects. With the aid of these cost savings the corresponding cost allocation rule $\pi(N, c, A) \in \mathbb{R}^N$, defined by

$$\pi_i(N, c, A) := c(\{i\}) - \sum_{a \in A_i} \frac{1}{2} \Phi_a(A, r_N^c)$$

for all $i \in N$, is called the position value of the communication situation (N, c, A) . Here, $A_i := \{\{i, j\} \in A \mid j \in N\}$ is the set of all communication links of which player i is an end point.

Both the Myerson value and the position value are linear with respect to the underlying cost game. Therefore, in deriving integral formulas for both values we can restrict our attention to dual unanimity games, because all cost games we consider are positive combinations of dual unanimity games (cf. theorem 1).

First we derive an integral formula for the Myerson value of a communication situation (N, u_M^*, A) , $M \subseteq N$, if the communication graph (N, A) is a tree.

Let (N, c, A) be a communication situation. As is well known, the game (N, c_A) can be written as

$$c_A = \sum_{S \in 2^N \setminus \{\emptyset\}} \Delta_{c_A}(S) u_S,$$

where u_S is the *unanimity game* on S , defined by

$$u_S(T) = \begin{cases} 1 & \text{if } S \subseteq T \\ 0 & \text{else} \end{cases}$$

and the *dividends* $\Delta_{c_A}(S)$ (cf. *Harsanyi* (1959)) are given by

$$\Delta_{c_A}(S) = \sum_{T \subseteq S} (-1)^{|S|-|T|} c_A(T)$$

for all $S \in 2^N \setminus \{\emptyset\}$.

In the following we consider a fixed communication situation (N, u_M^*, A) where $|N| \geq 2$ and (N, A) is a tree.

In this case the dividends $\Delta_{(u_M^*)_A}(S)$ are given by (cf. *Owen* (1986))

$$\Delta_{(u_M^*)_A}(S) = \begin{cases} (-1)^{|S \cap M|+1} & \text{if } (S, A(S)) \text{ is connected and } \text{Ext}(S, A(S)) = S \cap M \\ 0 & \text{else} \end{cases}$$

for all $S \in 2^N \setminus \{\emptyset\}$. Here, for a connected graph $(S, A(S))$

$$\text{Ext}(S, A(S)) := \{i \in S \mid |A_i \cap A(S)| \leq 1\}$$

denotes the set of *extreme points*.

Consequently,

$$\mu_i(N, u_M^*, A) = \Phi_i(N, (u_M^*)_A) = \sum_{S \in \Sigma(A): i \in S} \frac{(-1)^{|S \cap M|+1}}{|S|} \quad (1)$$

for all $i \in N$, where

$$\Sigma(A) := \{S \subseteq N \mid (S, A(S)) \text{ is connected and } \text{Ext}(S, A(S)) = S \cap M\}.$$

In order to apply (1) we have to find all elements in $\Sigma(A)$ containing a player i . This can be done in the following way.

For every $S \subseteq C$ with $C \in N/A$, we can define the *connected hull* of S (cf. *Owen* (1986)) by

$$H(S) := \bigcap \{T \mid S \subseteq T \subseteq C, (T, A(T)) \text{ is a connected graph}\}.$$

Let $i \in N$ and let $A_i := \{\{i, i(1)\}, \dots, \{i, i(t)\}\}$. Clearly, since (N, A) does not contain a cycle, the partition $N/(A \setminus A_i)$ contains t distinct components $C(1), \dots, C(t)$ with $i(k) \in C(k)$ for all $k \in \{1, \dots, t\}$. For each $k \in \{1, \dots, t\}$ we define a connected subgraph $(T(k), A(T(k)))$ of $(C(k), A(C(k)))$ by

$$T(k) := \bigcup_{j \in M \cap C(k)} \{H(\{i(k), j\}) \mid |H(\{i(k), j\}) \cap M| = 1\}. \quad (2)$$

So, in particular, $T(k) = \emptyset$ if $C(k) \cap M = \emptyset$, $T(k) = \{i(k)\}$ if $i(k) \in M$ and in all other cases $\text{Ext}(T(k), A(T(k))) \setminus \{i(k)\} = T(k) \cap M$.

In deriving a generating function (cf. *Owen* (1972)) for the Myerson value we suppose that each player has a probability x to be 'active' or 'operational' and we compute for each subgraph $(T(k), A(T(k)))$ the probability $P_k(x)$ that at least one of the players in $M \cap T(k)$ can actually interact with player i . Using the inclusion-exclusion principle this probability $P_k(x)$ is given by

$$P_k(x) = \sum_{S \subseteq (T(k) \cap M), S \neq \emptyset} (-1)^{|S|+1} x^{|H(S \cup \{i(k)\})|} \quad (3)$$

for all $k \in \{1, \dots, t\}$. Note that $P_k(x) = 0$ if $T(k) \cap M = \emptyset$. The expected costs player i generates by linking up the components $C(1), \dots, C(t)$ are described by the generating function $\theta_i(x)$, where

$$\theta_i(x) := \begin{cases} 1 - \sum_{k=1}^t P_k(x) & \text{if } i \in M \\ - \sum_{r=2}^t \sum_{K \subseteq \{1, \dots, t\}, |K|=r} (-1)^r \prod_{k \in K} P_k(x) & \text{if } i \notin M. \end{cases} \quad (4)$$

In particular we have that $\theta_i(x) = 0$ if $i \notin M$ and $t = 1$.

Now we can formulate

THEOREM 2. Let (N, A) be a tree with $|N| \geq 2$. Then for all $M \in 2^N \setminus \{\emptyset\}$,

$$\mu_i(N, u_M^*, A) = \int_0^1 \theta_i(x) dx \quad (5)$$

for all $i \in N$, where $\theta_i(x)$ is defined as in (4).

Proof. Let $i \in M$. Then

$$\begin{aligned}
\int_0^1 \theta_i(x) dx &= 1 - \sum_{k=1}^t \left(\int_0^1 P_k(x) dx \right) \\
&= 1 - \sum_{k=1}^t \left(\int_0^1 \sum_{S \subseteq (T(k) \cap M), S \neq \emptyset} (-1)^{|S|+1} x^{|H(S \cup \{i(k)\})|} dx \right) \\
&= 1 - \sum_{k=1}^t \sum_{S \subseteq (T(k) \cap M), S \neq \emptyset} \frac{(-1)^{|S|+1}}{|H(S \cup \{i(k)\})| + 1}. \tag{6}
\end{aligned}$$

Note that, since $i \in M$, for each $k \in \{1, \dots, t\}$ a coalition $S \subseteq T(k) \cap M$ with $S \neq \emptyset$ uniquely determines a set $T = H(S \cup \{i\}) \in \Sigma(A)$, which satisfies $\text{Ext}(T, A(T)) \setminus \{i\} = S$. Hence, (6) equals

$$1 - \sum_{k=1}^t \sum_{S \in \Sigma_k(A): i \in S, |S| \geq 2} \frac{(-1)^{|S \cap M|}}{|S|}, \tag{7}$$

where $\Sigma_k(A) := \{S \in \Sigma(A) \mid S \subseteq (T(k) \cup \{i\})\}$ for all $k \in \{1, \dots, t\}$.

The fact that $i \in M$ implies that for all $S \in \Sigma(A)$ with $i \in S$ and $|S| \geq 2$ there is a $k \in \{1, \dots, t\}$ such that $S \subseteq T(k) \cup \{i\}$. Therefore, (7) equals

$$1 - \sum_{S \in \Sigma(A): i \in S, |S| \geq 2} \frac{(-1)^{|S \cap M|}}{|S|}. \tag{8}$$

Since $i \in M$, we have $\{i\} \in \Sigma(A)$, so (8) equals

$$\sum_{S \in \Sigma(A): i \in S} \frac{(-1)^{|S \cap M|+1}}{|S|} = \mu_i(N, u_M^*, A).$$

Let $i \in N \setminus M$. First note that we may assume that $|\{k \in \{1, \dots, t\} \mid T(k) \neq \emptyset\}| \geq 2$, for otherwise $\theta_i(x) = 0$ and $\{S \in \Sigma(A) \mid i \in S\} = \emptyset$, so trivially (5) is satisfied.

Since for all $K \subseteq \{1, \dots, t\}$ with $|K| \geq 2$

$$\begin{aligned}
\prod_{k \in K} P_k(x) &= \prod_{k \in K} \left(\sum_{S \subseteq (T(k) \cap M): S \neq \emptyset} (-1)^{|S|+1} x^{|H(S \cup \{i(k)\})|} \right) \\
&= \sum_{(S(k))_{k \in K} \in \Gamma(K)} \left(\prod_{k \in K} (-1)^{|S(k)|+1} x^{|H(S(k) \cup \{i(k)\})|} \right), \tag{9}
\end{aligned}$$

where $\Gamma(K) := \{(S(k))_{k \in K} \mid S(k) \subseteq (T(k) \cap M), S(k) \neq \emptyset \text{ for all } k \in K\}$, it follows that

$$\begin{aligned}
\int_0^1 \theta_i(x) dx &= \sum_{r=2}^t \sum_{K \subseteq \{1, \dots, t\}, |K|=r} (-1)^{r+1} \sum_{(S(k))_{k \in K} \in \Gamma(K)} \frac{(-1)^{\sum_{k \in K} (|S(k)|+1)}}{1 + \sum_{k \in K} |H(S(k) \cup \{i(k)\})|} \\
&= \sum_{r=2}^t \sum_{K \subseteq \{1, \dots, t\}, |K|=r} \sum_{(S(k))_{k \in K} \in \Gamma(K)} \frac{(-1)^{1+\sum_{k \in K} |S(k)|}}{|H(\bigcup_{k \in K} S(k))|}. \tag{10}
\end{aligned}$$

Since $i \notin M$, for each $K \subseteq \{1, \dots, t\}$ with $|K| \geq 2$ a set $(S(k))_{k \in K} \in \Gamma(K)$ uniquely determines a set $T = H(\bigcup_{k \in K} S(k)) \in \Sigma(A)$, which satisfies $\text{Ext}(T, A(T)) = \bigcup_{k \in K} S(k)$ and $i \in T$. Hence, (10) equals

$$\sum_{S \in \Sigma(A): i \in S} \frac{(-1)^{|S \cap M|+1}}{|S|} = \mu_i(N, u_M^*, A). \quad \square$$

We illustrate the actual computation of the Myerson value in

EXAMPLE 1. Let $N = \{1, \dots, 10\}$ and $M = \{1, 3, 4, 7, 10\}$. The graph (N, A) is represented in figure 1.

figure 1

Consider player $2 \notin M$. Following (2) we obtain four subgraphs corresponding to $T(1) = \{1\}$, $T(2) = \{3\}$, $T(3) = \emptyset$ and $T(4) = \{6, 7, 8, 10\}$. The corresponding polynomials (cf. (3)) are $P_1(x) = P_2(x) = x$, $P_3(x) = 0$ and $P_4(x) = x^2 + x^3 - x^4$. Hence, according to (4),

$$\begin{aligned}
\theta_2(x) &= -P_1(x)P_2(x) - P_1(x)P_4(x) - P_2(x)P_4(x) + P_1(x)P_2(x)P_4(x) = \\
&= -x^2 - 2x^3 - x^4 + 3x^5 - x^6. \text{ So, } \mu_2(N, u_M^*, A) = \int_0^1 \theta_2(x) dx = -\frac{71}{105}.
\end{aligned}$$

Now consider player $3 \in M$. We obtain two subgraphs corresponding to $T(1) = \{4\}$ and $T(2) = \{1, 2, 6, 7, 8, 10\}$. The corresponding polynomials are $P_1(x) = x$ and $P_2(x) = x^2 + x^3 + x^4 - x^4 - x^5 - x^5 + x^6 = x^2 + x^3 - 2x^5 + x^6$. So, according to (4), $\theta_3(x) = 1 - P_1(x) - P_2(x) = 1 - x - x^2 - x^3 + 2x^5 - x^6$ and $\mu_3(N, u_M^*, A) = \int_0^1 \theta_3(x) dx = \frac{3}{28}$. For the sake of completeness we note that

$$\mu(N, u_M^*, A) = \frac{1}{420}(255, -284, 45, 210, 0, -249, 269, -123, 0, 297).$$

It may seem that the calculation of the polynomials $P_k(x)$ (cf. (3)) may be quite lengthy, especially if the sets $T(k) \cap M$ have a large number of elements. An alternative way to obtain these polynomials is described below.

Let $i \in N$ and let $(T(k), A(T(k)))$ be one of the connected subgraphs corresponding to player i as described in (2). Suppose each player p has a probability x_p to be 'active'. Consider the polynomial

$$1 - \prod_{j \in T(k) \cap M} (1 - \prod_{p \in H(\{i(k), j\})} x_p) \quad (11)$$

and expand it. Now reduce the obtained polynomial to a multilinear polynomial by the simple recourse of reducing each higher exponent to a 1. Finally, by replacing the probabilities x_p by the probability $x \in [0, 1]$ we obtain the polynomial $P_k(x)$.

In this, we are effectively using the multilinear extension (cf. *Owen (1972)*). As we know, the partial derivative $\frac{\partial f}{\partial x_i}$ of this extension corresponds to the expectation that other players will collaborate with player i . Typically, for $S \subseteq N \setminus \{i\}$, the term

$$u(S) := \prod_{j \in S} x_j \quad (12)$$

corresponds to the probability that all members of S collaborate, given that each $j \in S$ has probability x_j of collaboration and assuming independence. Now, if S and T are disjoint, then

$$u(S \cup T) = u(S) \cdot u(T), \quad (13)$$

since the players in S and T are independent. When $S \cap T \neq \emptyset$, however, (13) is not quite correct. Rather,

$$u(S \cup T) = \overline{u(S) \cdot u(T)}, \quad (14)$$

where the bar corresponds to a reduction operation: each exponent larger than 1 is reduced to 1. If, for example, $S = \{1, 2\}$ and $T = \{2, 3\}$, then $u(S) = x_1 x_2$, $u(T) = x_2 x_3$ and $u(S \cup T) = \overline{x_1 x_2^2 x_3} = x_1 x_2 x_3$.

EXAMPLE 2. Let (N, u_M^*, A) be the communication situation as described in example 1. Again consider player 2. For the subgraph corresponding to $T(1) = \{1\}$, expression (11) yields

$$1 - (1 - x_1),$$

which results in the polynomial $P_1(x) = x$.

In the same way we obtain $P_2(x) = x$ for the polynomial corresponding to $T(2) = \{3\}$. Defining the empty product to be 1, we easily see $P_3(x) = 0$.

Finally, for the subgraph corresponding to $T(4) = \{6, 7, 8, 10\}$, expression (11) yields

$$1 - (1 - x_6 x_7)(1 - x_6 x_8 x_{10}).$$

Expanding this, we obtain

$$x_6 x_7 + x_6 x_8 x_{10} - x_6^2 x_7 x_8 x_{10}.$$

Reducing this polynomial and then replacing all x_p 's by x , we obtain

$$x^2 + x^3 - x^4,$$

which is exactly the desired polynomial $P_4(x)$.

We now concentrate on the position value. The arc game corresponding to the communication situation (N, u_M^*, A) can be expressed as

$$r_N^{u_M^*} = \sum_{L \in 2^A \setminus \{\emptyset\}} \Delta_{r_N^{u_M^*}}(L) u_L,$$

where, for all $L \in 2^A \setminus \{\emptyset\}$, (A, u_L) is the (arc) unanimity game on L . The results of *Borm et al.* (1990) imply that for all $L \in 2^A \setminus \{\emptyset\}$

$$\Delta_{r_N^{u_M^*}}(L) = \begin{cases} (-1)^{|N(L) \cap M|} & \text{if } (N(L), L) \text{ is a tree and } \text{Ext}(N(L), L) = N(L) \cap M \\ 0 & \text{else,} \end{cases}$$

where $N(L)$ is the set of players who are end points of an arc in L . Hence, for all $a \in A$

$$\Phi_a(A, r_N^{u_M^*}) = \sum_{L \in \Lambda(A): a \in L} \frac{(-1)^{|N(L) \cap M|}}{|L|}, \quad (15)$$

where $\Lambda(A) := \{L \subseteq A \mid (N(L), L) \text{ is a tree, } \text{Ext}(N(L), L) = N(L) \cap M\}$.

In order to apply (15) we have to find all elements in $\Lambda(A)$ containing an arc a . This can be done in the following way.

Let $a = \{i(1), i(2)\} \in A$. Clearly, since (N, A) is a tree, the partition $N/(A \setminus \{a\})$ consists of two distinct components $C(1)$ and $C(2)$ with $i(1) \in C(1)$ and $i(2) \in C(2)$. For each $k \in \{1, 2\}$ we define the connected subgraph $(T(k), A(T(k)))$ of $(C(k), A(C(k)))$ as in (2). In deriving a generating function for $\Phi(A, r_N^{u, \star M})$ we suppose that each arc has a probability x to be 'available' and we compute for each subgraph $(T(k), A(T(k)))$ the probability $P_k(x)$ that at least one of the players in $M \cap T(k)$ can actually interact with player $i(k)$. Using the inclusion-exclusion principle this probability $P_k(x)$ is given by

$$P_k(x) = \sum_{S \subseteq (T(k) \cap M), S \neq \emptyset} (-1)^{|S|+1} x^{|A(H(S \cup \{i(k)\}))|} \quad (16)$$

for $k \in \{1, 2\}$. The expected cost savings an arc generates by linking up the components $C(1)$ and $C(2)$ are described by the generating function $\theta_a(x)$,

$$\theta_a(x) := P_1(x) \cdot P_2(x). \quad (17)$$

Now we can formulate

THEOREM 3. Let (N, A) be a tree with $|N| \geq 2$. Then for all $M \in 2^N \setminus \{\emptyset\}$,

$$\Phi_a(A, r_N^{u, \star M}) = \int_0^1 \theta_a(x) dx \quad (18)$$

for all $a \in A$, where $\theta_a(x)$ is defined as in (17).

Proof. Let $a \in A$. We may assume that $T(1)$ and $T(2)$ are both non-empty, for otherwise $\theta_a(x) = 0$ and $\{L \in \Lambda(A) \mid a \in L\} = \emptyset$, so, trivially (18) is satisfied. Now

$$\begin{aligned} \int_0^1 \theta_a(x) dx &= \int_0^1 P_1(x) P_2(x) dx \\ &= \int_0^1 \prod_{k=1}^2 \left(\sum_{S \subseteq (T(k) \cap M): S \neq \emptyset} (-1)^{|S|+1} x^{|A(H(S \cup \{i(k)\}))|} \right) dx \\ &= \int_0^1 \sum_{(S(1), S(2)) \in \Gamma(\{1, 2\})} \left(\prod_{k=1}^2 (-1)^{|S(k)|+1} x^{|A(H(S(k) \cup \{i(k)\}))|} \right) dx, \quad (19) \end{aligned}$$

where $\Gamma(\{1, 2\}) := \{(S(1), S(2)) \mid S(k) \subseteq (T(k) \cap M), S(k) \neq \emptyset \text{ for } k \in \{1, 2\}\}$.

Carrying out the integration in expression (19) we obtain

$$\begin{aligned}
& \sum_{(S(1), S(2)) \in \Gamma(\{1,2\})} \frac{(-1)^{\sum_{k=1}^2 (|S(k)|+1)}}{1 + \sum_{k=1}^2 |A(H(S(k) \cup \{i(k)\}))|} \\
&= \sum_{(S(1), S(2)) \in \Gamma(\{1,2\})} \frac{(-1)^{|S(1)|+|S(2)|}}{|A(H(S(1) \cup S(2)))|}. \tag{20}
\end{aligned}$$

Note that each pair $(S(1), S(2)) \in \Gamma(\{1,2\})$ uniquely determines a set $L = A(H(S(1) \cup S(2))) \in \Lambda(A)$, which satisfies $a \in L$ and $\text{Ext}(N(L), L) = S(1) \cup S(2)$. Hence, (20) equals

$$\sum_{L \in \Lambda(A): a \in L} \frac{(-1)^{|N(L) \cap M|}}{|L|} = \Phi_a(A, r_N^{\star M}). \quad \square$$

We illustrate the actual computation of the position value in

EXAMPLE 3. Let $(N, u_M^{\star}, A)$ be the communication situation as described in example 1. Consider player 6. To obtain the position value of player 6 we have to compute

$\Phi_{\{2,6\}}(A, r_N^{\star M})$, $\Phi_{\{6,7\}}(A, r_N^{\star M})$ and $\Phi_{\{6,8\}}(A, r_N^{\star M})$.

For $a := \{2,6\}$ we obtain (cf. (2)) the subgraphs corresponding to $T(1) = \{1,2,3\}$ and $T(2) = \{6,7,8,10\}$ with corresponding polynomials (cf. (16)) $P_1(x) = 2x - x^2$ and $P_2(x) = x + x^2 - x^3$. Hence, according to (17), $\theta_a(x) = 2x^2 + x^3 - 3x^4 + x^5$ and $\Phi_a(A, r_N^{\star M}) = \int_0^1 \theta_a(x) dx = \frac{29}{60}$.

For $b := \{6,7\}$ we obtain $T(1) = \{1,2,3,6,8,10\}$ and $T(2) = \{7\}$ and the polynomials $P_1(x) = 3x^2 - x^3 - 2x^4 + x^5$ and $P_2(x) = 1$. Hence, $\theta_b(x) = 3x^2 - x^3 - 2x^4 + x^5$ and $\Phi_b(A, r_N^{\star M}) = \int_0^1 \theta_b(x) dx = \frac{31}{60}$.

Finally, for $c := \{6,8\}$ we obtain $T(1) = \{1,2,3,6,7\}$ and $T(2) = \{8,10\}$ with corresponding polynomials $P_1(x) = x + 2x^2 - 3x^3 + x^4$ and $P_2(x) = x$. Hence, we have

$\theta_c(x) = x^2 + 2x^3 - 3x^4 + x^5$ and $\Phi_c(A, r_N^{\star M}) = \int_0^1 \theta_c(x) dx = \frac{2}{5}$.

We now compute

$$\pi_6(N, u_M^{\star}, A) = u_M^{\star}(\{6\}) - \sum_{\ell \in A_6} \frac{1}{2} \Phi_{\ell}(A, r_N^{\star M}) = 0 - \frac{1}{2} \left(\frac{29}{60} + \frac{31}{60} + \frac{24}{60} \right) = -\frac{7}{10}.$$

For the sake of completeness we note that

$$\pi(N, u_M^{\star}, A) = \frac{1}{120} (84, -101, 24, 60, 0, -84, 89, -48, 0, 96).$$

Similarly to the method described for the polynomials with respect to the Myerson value, there is an alternative way to compute the polynomials $P_k(x)$ as described in (16). The desired polynomial is obtained by considering the polynomial

$$1 - \prod_{j \in T(k) \cap M} (1 - \prod_{a \in A(H(\{i(k), j\}))} x_a)$$

and then following a reduction procedure similar to the one described before.

In deriving integral formulas for the Myerson value and the position value we restricted our attention to communication graphs that are trees. However, since for all communication situations (N, c, A) and all components $T \in N/A$ we have that

$$\mu_i(N, c, A) = \mu_i(T, c|_T, A(T))$$

and

$$\pi_i(N, c, A) = \pi_i(T, c|_T, A(T))$$

for all $i \in T$ (where $c|_T$ denotes the restriction of c to T), the integral formulas can be used to compute both values for communication situations with cycle-free communication graphs.

REFERENCES

- Borm, P.E.M., Owen, G., and Tijs, S. (1990). 'Values of points and arcs in communication situations'. Report 9004, Dept. of Mathematics, University of Nijmegen, The Netherlands.
- Harsanyi, J.C. (1959). 'A bargaining model for the cooperative n -person game'. *Annals of Math. Studies* 40, 325-356.
- Myerson, R.B. (1977). 'Graphs and cooperation in games'. *Math. Oper. Res.* 2, 225-229.
- Nouweland, A. van den, and Borm, P. (1990). 'On the convexity of communication games'. Report 9047, Dept. of Mathematics, University of Nijmegen, The Netherlands.
- Owen, G. (1972). 'Multilinear extensions of games'. *Management Science* 18, supplementary issue, P64-P79.
- Owen, G. (1986). 'Values of graph-restricted games'. *SIAM. J. Alg. Disc. Meth.* 7, 210-220.
- Shapley, L.S. (1953). 'A value for n -person games'. *Annals of Math. Studies* 28, 307-317.

IN 1990 REEDS VERSCHENEN

- 419 Bertrand Melenberg, Rob Alessie
A method to construct moments in the multi-good life cycle consumption model
- 420 J. Kriens
On the differentiability of the set of efficient (μ, σ^2) combinations in the Markowitz portfolio selection method
- 421 Steffen Jørgensen, Peter M. Kort
Optimal dynamic investment policies under concave-convex adjustment costs
- 422 J.P.C. Blanc
Cyclic polling systems: limited service versus Bernoulli schedules
- 423 M.H.C. Paardekooper
Parallel normreducing transformations for the algebraic eigenvalue problem
- 424 Hans Gremmen
On the political (ir)relevance of classical customs union theory
- 425 Ed Nijssen
Marketingstrategie in Machtsperspectief
- 426 Jack P.C. Kleijnen
Regression Metamodels for Simulation with Common Random Numbers: Comparison of Techniques
- 427 Harry H. Tigelaar
The correlation structure of stationary bilinear processes
- 428 Drs. C.H. Veld en Drs. A.H.F. Verboven
De waardering van aandelenwarrants en langlopende call-opties
- 429 Theo van de Klundert en Anton B. van Schaik
Liquidity Constraints and the Keynesian Corridor
- 430 Gert Nieuwenhuis
Central limit theorems for sequences with $m(n)$ -dependent main part
- 431 Hans J. Gremmen
Macro-Economic Implications of Profit Optimizing Investment Behaviour
- 432 J.M. Schumacher
System-Theoretic Trends in Econometrics
- 433 Peter M. Kort, Paul M.J.J. van Loon, Mikuláš Luptacik
Optimal Dynamic Environmental Policies of a Profit Maximizing Firm
- 434 Raymond Gradus
Optimal Dynamic Profit Taxation: The Derivation of Feedback Stackelberg Equilibria

- 435 Jack P.C. Kleijnen
Statistics and Deterministic Simulation Models: Why Not?
- 436 M.J.G. van Eijs, R.J.M. Heuts, J.P.C. Kleijnen
Analysis and comparison of two strategies for multi-item inventory systems with joint replenishment costs
- 437 Jan A. Weststrate
Waiting times in a two-queue model with exhaustive and Bernoulli service
- 438 Alfons Daems
Typologie van non-profit organisaties
- 439 Drs. C.H. Veld en Drs. J. Grazell
Motieven voor de uitgifte van converteerbare obligatieleningen en warrantobligatieleningen
- 440 Jack P.C. Kleijnen
Sensitivity analysis of simulation experiments: regression analysis and statistical design
- 441 C.H. Veld en A.H.F. Verboven
De waardering van conversierechten van Nederlandse converteerbare obligaties
- 442 Drs. C.H. Veld en Drs. P.J.W. Duffhues
Verslaggevingsaspecten van aandelenwarrants
- 443 Jack P.C. Kleijnen and Ben Annink
Vector computers, Monte Carlo simulation, and regression analysis: an introduction
- 444 Alfons Daems
"Non-market failures": Imperfecties in de budgetsector
- 445 J.P.C. Blanc
The power-series algorithm applied to cyclic polling systems
- 446 L.W.G. Strijbosch and R.M.J. Heuts
Modelling (s,Q) inventory systems: parametric versus non-parametric approximations for the lead time demand distribution
- 447 Jack P.C. Kleijnen
Supercomputers for Monte Carlo simulation: cross-validation versus Rao's test in multivariate regression
- 448 Jack P.C. Kleijnen, Greet van Ham and Jan Rotmans
Techniques for sensitivity analysis of simulation models: a case study of the CO₂ greenhouse effect
- 449 Harrie A.A. Verbon and Marijn J.M. Verhoeven
Decision-making on pension schemes: expectation-formation under demographic change

- 450 Drs. W. Reijnders en Drs. P. Verstappen
Logistiek management marketinginstrument van de jaren negentig
- 451 Alfons J. Daems
Budgeting the non-profit organization
An agency theoretic approach
- 452 W.H. Haemers, D.G. Higman, S.A. Hobart
Strongly regular graphs induced by polarities of symmetric designs
- 453 M.J.G. van Eijs
Two notes on the joint replenishment problem under constant demand
- 454 B.B. van der Genugten
Iterated WLS using residuals for improved efficiency in the linear model with completely unknown heteroskedasticity
- 455 F.A. van der Duyn Schouten and S.G. Vanneste
Two Simple Control Policies for a Multicomponent Maintenance System
- 456 Geert J. Almekinders and Sylvester C.W. Eijffinger
Objectives and effectiveness of foreign exchange market intervention
A survey of the empirical literature
- 457 Saskia Oortwijn, Peter Borm, Hans Keiding and Stef Tijs
Extensions of the τ -value to NTU-games
- 458 Willem H. Haemers, Christopher Parker, Vera Pless and Vladimir D. Tonchev
A design and a code invariant under the simple group Co_3
- 459 J.P.C. Blanc
Performance evaluation of polling systems by means of the power-series algorithm
- 460 Leo W.G. Strijbosch, Arno G.M. van Doorne, Willem J. Selen
A simplified MOLP algorithm: The MOLP-S procedure
- 461 Arie Kapteyn and Aart de Zeeuw
Changing incentives for economic research in The Netherlands
- 462 W. Spanjers
Equilibrium with co-ordination and exchange institutions: A comment
- 463 Sylvester Eijffinger and Adrian van Rixtel
The Japanese financial system and monetary policy: A descriptive review
- 464 Hans Kremers and Dolf Talman
A new algorithm for the linear complementarity problem allowing for an arbitrary starting point
- 465 René van den Brink, Robert P. Gilles
A social power index for hierarchically structured populations of economic agents

IN 1991 REEDS VERSCHENEN

- 466 Prof.Dr. Th.C.M.J. van de Klundert - Prof.Dr. A.B.T.M. van Schaik
Economische groei in Nederland in een internationaal perspectief
- 467 Dr. Sylvester C.W. Eijffinger
The convergence of monetary policy - Germany and France as an example
- 468 E. Nijssen
Strategisch gedrag, planning en prestatie. Een inductieve studie
binnen de computerbranche

Bibliotheek K. U. Brabant

17 000 01066358 2