Tilburg University

Analysis and computation of (n,N)

van der Duyn Schouten, F.A.; Vanneste, S.G.

Publication date:
1989

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

van der Duyn Schouten, F. A., & Vanneste, S. G. (1989). *Analysis and computation of (n,N) : Strategies for maintenance of a two-component system*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 372). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM

CBM
R

UNIVERSITY

LE

UNIVERSITEIT

BRABANT

7626
1989
372

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS

* C I N O 1 3 6 9 *

CBM

~~ER~~

7626

1989

nr. 372

DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM

**ANALYSIS AND COMPUTATION OF (n,N) -
STRATEGIES FOR MAINTENANCE OF A
TWO-COMPONENT SYSTEM**

F.A. van der Duyn Schouten
S.G. Vanneste

FEW 372

Tilburg University
Faculty of Economics
P.O. Box 90153
5000 LE Tilburg
The Netherlands

R61

658.58

69

Abstract

In this paper we investigate (n,N) -strategies for the maintenance of a two component series system. Vergin and Scriabin [9] provided some numerical evidence on the near-optimality of this type of policies. In a recent paper Özekici [6] gave a characterization of the structure of the optimal policy and its possible deviations from the (n,N) -structure for the discounted cost criterion. The same kind of structure is shown to hold for the average cost criterion. In view of the complicated form of the optimal policy and the near-optimality of the (n,N) -policies we present a fast computational method to compute the average costs under a given (n,N) -policy. This method is based on a well-know embedding technique. Moreover, a heuristic based on this computational method, is presented by which the optimal values of n and N can be determined.

Key words and phrases: Multicomponent reliability system; preventive and opportunistic replacement; Markov decision processes; control limit rules; computational aspects.

1. Introduction

In this paper we investigate the two component maintenance scheduling problem introduced by Vergin and Scriabin [9]. This problem concerns the maintenance of a series system, consisting of two (identical) components, which are subject to stochastic failures. The components are functioning independent of each other. When one of the components breaks down the system fails which involves break-down costs. To avoid a frequent occurrence of this cost preventive replacement is allowed. Additionally we have the option of opportunistic replacement, which refers to the possibility to replace both components, when only one of them has failed or reached its preventive age. This makes sense not only because it usually is cheaper to replace two components simultaneously rather than separately but also due to the series structure of the system. Replacements can only take place at inspection epochs, which are assumed to be equidistant discrete time epochs.

To find an optimal maintenance schedule one might use standard policy-improvement or value iteration techniques from Markov decision theory. However, there are two drawbacks. First of all due to the two dimensional state space (denoting the ages of both components) these standard techniques are in general very time-consuming and even prohibitive for moderate size problems. In the second place the actual optimal policy, found by one of these procedures, may have a rather irregular structure, which makes it necessary to explicitly specify the optimal decision for all possible age-combinations. From a practical point of view one might be interested in policies which are easy to characterize and, on the other hand, close to optimality. Moreover, it is important that for the proposed policy the actual average expected costs per unit time can be computed rather easily. A class of policies satisfying these conditions are the so-called (n, N) -policies. An (n, N) -policy prescribes to replace a component when it has failed or when its age has reached the value N and, if one of the components is replaced, to replace the other simultaneously when its age is greater than or equal to n ($\leq N$). In this sense an (n, N) -policy seems to be the natural two-dimensional generalization of the well-known one-dimensional control-limit rules.

Vergin and Scriabin [9] have shown by numerical comparisons that (n,N) -policies are close to optimality for a wide range of cost parameters. Özekici [6] provided some additional support for this conclusion based on the analysis of the optimal policy. These results are mentioned in more detail in section 3. The main result of this paper is dealt with in section 4. There we present an exact computational method for the average costs under a given (n,N) -policy. This method is based on a well-known embedding technique (see e.g. Tijms and Van der Duyn Schouten [8]). Finally we use this computational method in section 5 as a building block in a heuristic to determine the best policy within the class of (n,N) -policies.

Recently various authors studied models which are closely related to ours. Epstein and Wilamowsky [2] investigate a system consisting of two components one of which has an exponential lifetime, while the other has a fixed finite lifelength with failure rate equal to zero. A characterization of the optimal replacement policy is given. Opportunistic maintenance is also the subject of a paper by Bäckert and Rippin [1]. They consider a three component system. At failure epochs of one of these components opportunistic replacement of the other components is considered. The performance of three different solution techniques, among which the Markov decision approach, are compared. Finally we mention a paper by Haurie and l'Ecuyer [3]. They analyse the optimal preventive replacement rule over a finite horizon of a parallel system comprised of m identical independent components. There are fixed as well as linear replacement costs but no break-down costs. It is shown that the optimal policy may have some counterintuitive properties with respect to monotonicity in its state vector.

2. Mathematical model and its solution techniques

Consider a series system with two identical and independently operating components 1 and 2. Both components are subject to stochastic failures. The times until failure are stochastic variables with known probability distribution function. The system is inspected only at discrete equidistant time epochs. When upon inspection a component turns out to have failed during the last period it has to be replaced immediately and a breakdown cost is incurred. The time needed for replacement is negligible.

On the other hand when at an inspection epoch both components are still working there are four possible decisions to make: replace none component, replace only component 1, replace only component 2 or replace both components. A replacement in this situation is called a preventive replacement or an opportunistic replacement. The former refers to the opportunity to prevent breakdown cost by timely replacement, the latter refers to the opportunity to prevent cumulation of replacement costs by joint replacement instead of separate replacements. After replacement of a component a new identical component starts with age equal to zero. The lifetime distributions are supposed to have finite support i.e. when a component has reached its maximal lifelenght it certainly will fail during the next period.

The following non-negative costs are incurred:

- b := system breakdown cost.
- r_1 := replacement cost for a single component (either 1 or 2).
- r_{12} := cost of joint replacement of both components 1 and 2.

Note that due to the series structure the same breakdown cost b is incurred when either one or both components fail.

Assumption 1. $r_1 \leq r_{12} \leq 2r_1$ □

Next we consider the aging process of the system. As stated before we assume that both components are independent. For a relaxation of this assumption we refer to the recent paper of Özekici [6]. (see also section 3).

Define

L_i := the lifetime of component i expressed in inspection periods, $i=1,2$.

Let

$\mu := E(L_i), i=1,2$ and

$$q_n := P(L_i = n | L_i \geq n), \quad n=0, \dots, m; \quad i=1,2$$

where

$$m := \min \{n: P(L_i \geq n+1) = 0\}$$

and

$$p_n := 1 - q_n, \quad n=0, \dots, m.$$

Note that q_n denotes the probability that a component will fail during the next period given that its age equals n at the beginning of this period. We say that a component at an inspection epoch is found in state n when it has not failed and its age equals n . When the component has failed we say that it is in state $m+1$.

To model this system as a Markov decision process we introduce as a state space

$$S = \{1, 2, \dots, m+1\}^2$$

where for $(i, j) \in S$ we denote by $i(j)$ the state of the first (second) component.

The set of possible actions A consists of:

- i := replace only component i , $i=1,2$
- 12 := replace both components
- 0 := replace no component

The set of feasible actions in state $s \in S$ is denoted by $A(s)$.

The one-step cost function $c(s, a)$, $s \in S$, $a \in A$ is easily composed from the given cost parameters b , r_1 and r_{12} . For example

$$c((m+1, j), 12) = b + r_{12}, \quad 1 \leq j \leq m+1$$

$$c((i, j), 12) = r_{12}, \quad 1 \leq i, j \leq m$$

$$c((i,j),1) = r_1, \quad 1 \leq i,j \leq m$$

Finally the one-step transition probabilities $p_{st}(a)$, $s,t \in S$, $a \in A$ are completely determined by the survival probabilities p_n . For example

$$p_{(i,j),(i+1,j+1)}^{(0)} = p_i p_j, \quad 1 \leq i,j \leq m-1$$

$$p_{(i,j),(m+1,j+1)}^{(0)} = q_i p_j, \quad 1 \leq i \leq m, \quad 1 \leq j \leq m-1$$

$$p_{(i,j),(1,1)}^{(12)} = p_0^2, \quad 1 \leq i,j \leq m+1.$$

Assumption 2. $0 = p_m < p_{m-1} \leq \dots \leq p_1 \leq p_0 < 1$. □

As far as the lifetime distribution of the components is concerned knowledge of the sequence $(p_k)_{k=0}^m$ suffices. It is irrelevant whether the lifetimes themselves have a discrete or continuous probability distribution. □

Denote by $X(t) = (X_1(t), X_2(t))$ the states of both components at the t -th inspection epoch. For every stationary policy R the process $\{X(t), t = 0, 1, 2, \dots\}$ constitutes a discrete-time Markov chain on S . It is easy to see that $(m+1, m+1)$ is a positive recurrent state under every stationary policy R . Due to assumption 2 it follows that $(m+1, m+1)$ can be reached from every starting state (i, j) . This implies that there exists only one recurrent class under every policy R . Let $C(R)$ denote this recurrent class and let $D(R)$ be the set of transient states under policy R . Note that state $(i, j) \in D(R)$ iff it cannot be reached from $(m+1, m+1)$. Finally we note that $X(t)$ is aperiodic on $C(R)$ for every policy R since we can return to $(m+1, m+1)$ in one step. We are now interested in that stationary policy R that minimizes

$$g(R) := \lim_{n \rightarrow \infty} E_R \frac{1}{n+1} \sum_{t=0}^n c(X(t), R(X(t)))$$

The standard tools to solve this Markov decision problem numerically are the policy-iteration and the value-iteration algorithm (see e.g. Tijms [7]). From a numerical point of view the value-iteration algorithm is

preferred in particular for big state spaces. This algorithm recursively solves the finite horizon Bellman optimality equation

$$(2.1) \quad V_n(s) = \min_{a \in A(s)} \{c(s,a) + \sum_{t \in S} p_{st}(a) V_{n-1}(t)\}, \quad s \in S, n \geq 1.$$

On the other hand we know from Markov decision theory (see e.g. Howard [4] or Tijms [7]) that for any stationary policy R the average costs $g(R)$ and the so-called relative values $v_R(s)$, $s \in S$ are the unique solution to the set of linear equations

$$(2.2) \quad \begin{cases} v_R(s) = c(s, R(s)) - g(R) + \sum_{t \in S} p_{st}(R(s)) v_R(t), & s \in S \\ v_R(m+1, m+1) = 0. \end{cases}$$

A step in the general policy improvement algorithm proceeds as follows. First we define for a given policy R the policy improvement quantity

$$(2.3) \quad T_R(s, a) := c(s, a) - g(R) + \sum_{t \in S} p_{st}(a) v_R(t)$$

For any stationary policy $\bar{R}$ we have (see Tijms [7])

$$g(\bar{R}) \leq g(R)$$

whenever

$$T_R(s, \bar{R}(s)) \leq v_R(s) \quad \text{for all } s \in S.$$

In section 5 we use this result to derive a policy improvement procedure within the class of (n, N) -policies.

To conclude this section we note that the average optimal policy R^* can be found as the minimizing action in the average optimality equation (see e.g. Tijms [7])

$$(2.4) \quad \begin{cases} v(s) = \min_{a \in A(s)} \{c(s,a) - g + \sum_{t \in S} p_{st}(a) v(t)\} \\ v(m+1, m+1) = 0 \end{cases}$$

3. The structure of the optimal policy

Up to now the most complete characterization of the structure of the optimal policy has been given by Özekici [6]. Özekici's model is more general than ours in the followings aspects. The system considered by Özekici consists of n different components whose ages develop over time as an increasing Hunt process, which implies amongst others that the ages of the components are not necessarily independent and may increase at different rates as well as by jumps. Our cost structure is also a special case of the cost structure assumed by Özekici: a general maintenance/replacement cost function depending on the components to be replaced (but not on their ages) and an operating cost function of the age-vector. Under fairly general assumptions on both cost functions and aging process Özekici shows that for the optimal replacement policy the four regions of the state space where the four different actions 0, 1, 2 and 12 are optimal are connected subsets of the state space with certain regularity conditions for the boundaries. A typical example of the optimal policy for two components in Özekici's model is shown in figure 1

Figure 1 The general form of the optimal policy in Özekici's model.

Let us denote the region where the optimal policy prescribes action a by R_a , $a \in A$.

The special characteristics of the optimal policy are:

- (i) the boundaries between R_1 and R_{12} and between R_2 and R_{12} are straight lines.
- (ii) the other boundaries do not necessarily have a convex or concave shape
- (iii) for fixed age i of one component the replacement of the other component is controlled by a control limit rule i.e.

$$(i, j) \in R_2 \cup R_{12} \text{ implies } (i, j+k) \in R_2 \cup R_{12} \text{ for } k \geq 0.$$
- (iv) If $(i, j) \in R_2$ then $(i, k) \in R_2$ for $m+1 \geq k \geq j$
- (v) If $(i, j) \in R_{12}$ then $(k, l) \in R_{12}$ for all $m+1 \geq k \geq i$ and all $m+1 \geq l \geq j$
- (vi) it may occur that $(i, j) \in R_2$ and $(i+k, j) \in R_0$ for some $k \geq 0$ (similarly for R_1 and R_0)
- (vii) it may occur that $(i, j) \in R_0$ and $(i, j+k) \in R_1$ for some $k \geq 0$ (similarly for R_0 and R_2)

The characteristics (i) up to (v) are proven by Özekici for the discounted cost criterion and below we will show that they hold for the average cost criterion. From extensive numerical experiments we have found situations where (ii) and (vi) occur for the average cost criterion (see figure 2). The situation mentioned under (vii) did not occur. We conjecture that it cannot occur, although we were not able to prove it.

Figure 2 Some specific forms of the optimal maintenance policy.

Next we derive the properties mentioned above of the average optimal policy. Let us specify the Bellman optimality equation (2.1) for our model. Define for $0 \leq i, j \leq m$

$$(3.1) \quad W_{n+1}(i, j) := p_i p_j V_n(i+1, j+1) + (1-p_i) p_j V_n(m+1, j+1) + \\ + p_i (1-p_j) V_n(i+1, m+1) + (1-p_i) (1-p_j) V_n(m+1, m+1)$$

Now relation (2.1) can be rewritten as follows

$$(3.2) \quad V_{n+1}(i, j) = \min \{W_{n+1}(i, j), r_1 + W_{n+1}(0, j), r_1 + W_{n+1}(i, 0), \\ r_{12} + W_{n+1}(0, 0)\}$$

$$(3.3) \quad V_{n+1}(i, m+1) = \min \{b + r_1 + W_{n+1}(i, 0), b + r_{12} + W_{n+1}(0, 0)\}$$

$$(3.4) \quad V_{n+1}(m+1, j) = \min \{b + r_1 + W_{n+1}(0, j), b + r_{12} + W_{n+1}(0, 0)\}$$

$$(3.5) \quad V_{n+1}(m+1, m+1) = b + r_{12} + W_{n+1}(0, 0).$$

Similarly the average optimality equation (2.4) can be rewritten into

$$(3.6) \quad v(i, j) = \min \{-g + w(i, j), r_1 - g + w(0, j), r_1 - g + w(i, 0), \\ r_{12} - g + w(0, 0)\}$$

$$(3.7) \quad v(i, m+1) = \min \{b + r_1 - g + w(i, 0), b + r_{12} - g + w(0, 0)\}$$

$$(3.8) \quad v(m+1, j) = \min \{b + r_1 - g + w(0, j), b + r_{12} - g + w(0, 0)\}$$

$$(3.9) \quad v(m+1, m+1) = b + r_{12} - g + w(0, 0)$$

where for $0 \leq i, j \leq m$

$$(3.10) \quad w(i,j) = p_i p_j v(i+1,j+1) + (1-p_i) p_j v(m+1,j+1) + \\ + p_i (1-p_j) v(i+1,m+1) + (1-p_i)(1-p_j) v(m+1,m+1)$$

Theorem 1 The average optimal policy R^* is symmetric.

Proof From (3.2) unto (3.5) it easily follows by induction on n that $V_n(\dots)$ is symmetric starting with $V_0(\dots) \equiv 0$. Since the solution $v(\dots)$ of the average optimality equations (3.6) unto (3.9) satisfy (see Tijms [7])

$$v(i,j) - v(k,\ell) = \lim_{n \rightarrow \infty} \{V_n(i,j) - V_n(k,\ell)\}$$

it follows that $v(\dots)$ is symmetric, which in turn implies that the minimizing action in the right hand side of (3.6) unto (3.9) is symmetric. The observation that the minimizing action in the right hand side of the average optimality equation constitutes an average optimal policy (see Tijms [7]) completes the proof. $\square$

Theorem 2 Any solution of the equations (3.6) unto (3.9) satisfy

$$(3.11) \quad v(i,j) \leq v(i,j+1) , \quad 1 \leq i \leq m+1 , \quad 1 \leq j \leq m$$

and

$$(3.12) \quad v(i,j) \leq v(i+1,j) , \quad 1 \leq i \leq m , \quad 1 \leq j \leq m+1$$

Proof By theorem 1 it is sufficient to show that (3.11) holds. From (3.6) unto (3.9) it follows that for $1 \leq i,j \leq m+1$

$$(3.13) \quad v(i,j) \leq b + r_{12} - g + w(0,0) = v(m+1,m+1)$$

Rewrite (3.10) into

$$\begin{aligned}
(3.14) \quad w(i, j) &= v(m+1, m+1) - p_i p_j (v(m+1, m+1) - v(i+1, j+1)) - \\
&- (1-p_i) p_j (v(m+1, m+1) - v(m+1, j+1)) - p_i (1-p_j) (v(m+1, m+1) - \\
&- v(i+1, m+1))
\end{aligned}$$

From (3.6) unto (3.9) it follows that (3.11) trivially holds for $j=m$ and all $1 \leq i \leq m+1$. Now suppose that (3.11) holds for all $k+1 \leq j \leq m$ and all $1 \leq i \leq m+1$.

To show that (3.11) also holds for $j=k$ and all $1 \leq i \leq m+1$ it suffices to prove that for all $0 \leq i \leq m$

$$(3.15) \quad w(i, k) \leq w(i, k+1)$$

From (3.14) it follows that

$$w(i, k) - w(i, k+1) = -\Delta_1 - \Delta_2 + \Delta_3$$

where

$$\begin{aligned}
\Delta_1 &:= p_i p_k (v(m+1, m+1) - v(i+1, k+1)) - p_i p_{k+1} (v(m+1, m+1) - \\
&- v(i+1, k+2))
\end{aligned}$$

$$\begin{aligned}
\Delta_2 &:= (1-p_i) p_k (v(m+1, m+1) - v(m+1, k+1)) - (1-p_i) p_{k+1} (v(m+1, m+1) - \\
&- v(m+1, k+2))
\end{aligned}$$

$$\Delta_3 := p_i (p_k - p_{k+1}) (v(m+1, m+1) - v(i+1, m+1))$$

From assumption 2 and the induction hypothesis it follows that $\Delta_1 \geq 0$ and $\Delta_2 \geq 0$. Next $\Delta_3 \geq 0$ follows from assumption 2 and relation (3.13). Finally we note that by the induction hypothesis

$$\begin{aligned}
\Delta_1 &\geq p_i p_k (v(m+1, m+1) - v(i+1, k+2)) - p_i p_{k+1} (v(m+1, m+1) - \\
&\quad - v(i+1, k+2)) = p_i (p_k - p_{k+1}) (v(m+1, m+1) - v(i+1, k+2)) \\
&\geq p_i (p_k - p_{k+1}) (v(m+1, m+1) - v(i+1, m+1)) \\
&= \Delta_3
\end{aligned}$$

which yields (3.15) □

Corollary 1 The average optimal policy R^* with regions R_a^* , $a \in A$ has the following properties:

- (i) If $j > i$ than the optimal policy can be chosen such that $R^*(i, j) \neq 1$
- (ii) If $(i, j) \in R_2^*$ than $(i, j+1) \in R_2^*$
- (iii) If $(i, j) \in R_{12}^*$ than $(k, l) \in R_{12}^*$ for all $k \geq i$ and all $l \geq j$.

Proof

- (i) For $j > i$ we have

$$w(i, 0) = w(0, i) \leq w(0, j).$$

Hence

$$r_1 - g + w(i, 0) \leq r_1 - g + w(0, j)$$

which implies that action 2 is always as good as action 1 in the optimization of the right hand side of (3.6)

(ii) For $(i,j) \in R_2^*$ it follows that

$$\begin{aligned} r_1 + w(i,0) &\leq \min \{w(i,j), r_1 + w(0,j), r_{12} + w(0,0)\} \\ &\leq \min \{w(i,j+1), r_1 + w(0,j+1), r_{12} + w(0,0)\} \end{aligned}$$

Hence $(i,j+1) \in R_2^*$.

(iii) The proof of (iii) proceeds similarly as (ii). □

Note that corollary 1 implies that the boundary between R_2^* and R_{12}^* is a straight line.

Besides the theoretical results contained in corollary 1 additional insight can be obtained by numerical experiments. By the value iteration algorithm we generated optimal policies under a wide variety of cost parameters and lifetime distributions. This general purpose algorithm works satisfactorily on a personal computer for a state space upto $m = 15$. The number of iterations required varied between 20 and 100.

In the appendix we present the numerical results. Apart from the optimal value of the average costs g^* we also give the average costs $g_{(n,N)}^*$ of the best policy of (n,N) -type. A policy R is called of (n,N) -type if there exist natural numbers $n \leq N$ such that $R(i,j) = 0$ if $0 \leq i,j \leq N$, $R(i,j) = 1$ if $i \geq N, j < n$; $R(i,j) = 2$ if $j \geq N, i < n$ and $R(i,j) = 12$ otherwise (see figure 2d).

A summary of the numerical results is presented in table 1 below where the frequency of occurrence of different values of $g_{(n,N)}^* - g^*$ as a percentage of g^* is tabulated.

%	0	0-0.1	0.1-0.2	0.2-0.3	0.3-0.4	0.4-0.5	0.5-0.6	>0.6
#	30	6	5	2	0	1	1	0

Table 1 Frequency of relative differences between $g_{(n,N)}^*$ and g^* .

In those cases where the difference between $g_{(n,N)}^*$ and g^* was equal to zero, the optimal policy was not necessarily of (n,N) -type. It sometimes occurs that the optimal policy itself (as it is generated by the value iteration algorithm) is not of (n,N) -type, but can be put into this form by modification of the actions on the set of transient states. It is not hard to see that such a modification only affects the value of the relative values on the transient states but not of the average costs itself. Moreover such a modification does not change the set of recurrent states. We conclude this section with a remark concerning another experimental finding. We generated a number of survival distributions by a discretization of a given Weibull distribution. We investigated the effect of a decreasing grid size (increasing number of possible states) on the form of the optimal policy. It turned out that the value of $g_{(n,N)}^*$ approached the value of g^* , although we could not conclude that the form of the optimal policy converged to a (n,N) -form.

Both the theoretical and numerical results reported in this section justify a special attention to policies of (n,N) -type, in particular because these kind of policies are easy to handle in practice. Therefore we present in the next section an embedding technique by which the average costs under a given (n,N) -policy can be efficiently computed. This technique is used as a building block in a heuristical algorithm to compute the best (n,N) -policy. This algorithm will be presented in section 5.

4. Analysis of a (n,N) -policy via embedding

In this section we present an efficient numerical method to compute the average costs $g(n,N)$ under a fixed policy of (n,N) -type. This method turns out to be faster than the application of the value iteration method for a fixed policy. Vergin and Scriabin [9] suggest simulation as an effective method to compute $g(n,N)$.

In this section we consider a fixed (n,N) -policy with average costs $g(n,N)$.

Let $X(t) = (X_1(t), X_2(t))$ denote the Markov chain on S describing the states of both components under the given (n,N) -policy and suppose that $X(0) = (m+1, m+1)$. Denote by T_n , $n \geq 1$ the epoch of the n -th replacement

(either a single replacement of one of both types or a joint replacement) and define

$$Z_0 := X(0)$$

and

$$Z_n := X(T_n), n \geq 1$$

Then $(Z_n)_{n=0}^{\infty}$ constitutes a Markov chain on state space

$$\begin{aligned} \tilde{S} := & \{(i, N) : 1 \leq i \leq N\} \cup \{(N, j) : 1 \leq j \leq N\} \cup \{(i, m+1) : 1 \leq i \leq N\} \cup \\ & \cup \{(m+1, j) : 1 \leq j \leq N\} \cup \{(m+1, m+1)\} \end{aligned}$$

Note that state (i, N) with $i < N-n+1$ can only be entered by $(Z_n)_{n=0}^{\infty}$ from $(N, N-i)$ or $(m+1, N-i)$. However, since $N-i > n-1$ the policy prescribes action 12 in $(N, N-i)$ and $(m+1, N-i)$. Hence (i, N) is not accessible under the given policy, so that the state space can be reduced to

$$\begin{aligned} E := & \{(i, N) : N-n+1 \leq i \leq N\} \cup \{(N, j) : N-n+1 \leq j \leq N\} \cup \\ & \cup \{(i, m+1) : 1 \leq i \leq N\} \cup \{(m+1, j) : 1 \leq j \leq N\} \cup \{(m+1, m+1)\}. \end{aligned}$$

Note that this state space is of dimension $2(n+N) \leq 4m$, while S is of dimension m^2 .

$(Z_n)_{n=0}^{\infty}$ is an irreducible, aperiodic, positive recurrent Markov chain on E . Let $p(\dots)$ denote the one-step transition probabilities of $(Z_n)_{n=0}^{\infty}$ and $\pi(\cdot)$ its stationary distribution.

For $s \in E$ we define

$T(s) :=$ the time until next replacement, given that the present replacement is carried out in state s

$$\tau(s) := E T(s)$$

$c(s)$:= the expected costs incurred until next replacement, given that the present replacement is carried out in state s .

Then we have by the theory of regenerative processes (see e.g. Tijms et al. [8])

$$(4.1) \quad g(n, N) = \frac{\sum_{s \in E} c(s) \pi(s)}{\sum_{s \in E} \tau(s) \pi(s)}$$

Before we derive explicit expressions for $c(s)$ and $\tau(s)$ we note that because of symmetry it suffices to compute both functions in the states (i, N) for $i \leq N$ and $(i, m+1)$ of E .

For the function $c(s)$ we have

$$c(i, N) = r_1, \quad c(i, m+1) = b + r_1 \quad \text{for } i < n$$

$$c(i, N) = r_{12}, \quad c(i, m+1) = b + r_{12} \quad \text{for } i \geq n$$

For $i < n$ we have

$$(4.2) \quad \tau(i, N) = \tau(i, m+1) = \sum_{k=1}^{\infty} P(T(i, m+1) \geq k) = 1 + \sum_{k=2}^{N-i} r(k-1, i) r(k-1, 0)$$

where

$$r(k, i) := \prod_{\ell=0}^{k-1} p_{i+\ell}$$

Note that $r(k, i)$ equals $P(L \geq i+k | L \geq i)$, where L denotes the lifetime of a single component.

Similarly we find for $i \geq n$

$$(4.3) \quad \tau(i, N) = \tau(i, m+1) = \tau(m+1, m+1) = 1 + \sum_{k=2}^N r^2(k-1, 0).$$

Finally the stationary distribution can be obtained by solution of the set of stationary equations $\pi = \pi P$ together with the normalizing equation

$$\sum_{s \in E} \pi(s) = 1$$

This set of equations can be solved by standard procedures (see Tijms [7]) once the one step transition probabilities $p(s, t)$ are known.

In figure 3 below the possible transitions from (i, N) are shown.

Figure 3 The possible one-step transitions from (i, N)

For $i < n$ the transition probabilities are given by

$$p((i, N), (N, N-i)) = r(N-i, i) r(N-i, 0)$$

$$p((i, N), (i+k, m+1)) = r(k, i) r(k-1, 0) (1-p_{k-1}) \quad (1 \leq k \leq N-i)$$

$$p((i, N), (m+1, k)) = r(k-1, i) (1-p_{i+k-1}) r(k, 0) \quad (1 \leq k \leq N-i)$$

$$p((i, N), (m+1, m+1)) = 1 - \sum_{s \neq (m+1, m+1)} p((i, N), s)$$

For $p((i, N), s)$ with $i \geq n$ we get the same expressions with $i=0$ substituted in the right hand sides. Finally we note that

$$(4.4) \quad p((i, m+1), s) = p((i, N), s) \text{ for all } i.$$

Instead of the solution of the stationary equations $\pi = \pi P$ by a standard numerical procedure, like Gauss-Seidel, we may follow another approach to find the value of $g(n, N)$, namely the solution of the optimality equation corresponding with the semi-Markov decision process in which decisions can be taken at epochs of entrance into the states N and $m+1$ and where the decisions are prescribed by the given (n, N) -rule.

This yields an "optimality" equation

$$(4.5) \quad \begin{cases} v(s) = c(s) - g \tau(s) + \sum_{t \in E} p(s,t) v(t) & , s \in E. \\ v(m+1, m+1) = 0. \end{cases}$$

This system can be solved by the value-iteration method for semi-Markov decision processes (see Tijms [7]). Again we emphasize that the solution of (20) and (21) is much easier than solution of (3) because of the reduction of the state space. In section 5 we will show that (4.5) can even be reduced to a system of only n equations.

The choice between both methods is rather arbitrary and may depend on the preference of the user. The solution of the steady state equations yields the stationary distribution, while solution of (4.5) by the value iteration method yields bounds on the value of $g(n,N)$ during the computational procedure. In our calculations we used the value iteration method.

5. A Heuristic for the optimal (n,N) -policy

In this section we describe a heuristic to compute the optimal replacement policy within the class of (n,N) -policies. In this algorithm the method to compute $g(n,N)$ for fixed (n,N) -policy as described in the previous section is used as a building block. The main idea of the algorithm is contained in the following three steps.

Step 1. Choose good starting values for n and N . Go to 2.

Step 2. Compute $g(n,N)$ and the relative values $v_{(n,N)}(s)$, $s \in S$ and go to 3.

Step 3. Use a special form of the policy improvement procedure to establish whether improved values of n and N exist. If so go to 2 else stop.

Below we will elaborate on these steps in more detail.

Step 1. In step 1 we use as an initial choice of n and N the optimal control limit T^* for the one component replacement model with breakdown costs

b and replacement costs r_1 . By standard arguments from renewal theory it can be shown that T^* is the minimizing argument of

$$g(T) := \frac{r_1 + b \sum_{n=0}^T a_n}{\sum_{k=0}^T \sum_{\ell=k}^m a_\ell}$$

where

$$a_k := P(L_1=k), \quad k=0,1,2,\dots; \quad i=1,2.$$

Using assumption 2 it is easily verified that T^* is the smallest natural number k satisfying

$$(5.1) \quad a_k \sum_{i=0}^{k-1} \sum_{\ell=i}^m a_\ell - \sum_{i=0}^{k-1} a_i \geq r_1/b$$

A natural number k satisfying (5.1) exists if

$$(5.2) \quad \mu - 1 \geq r_1/b$$

If (5.2) is not satisfied we have $T^* = m+1$.

Step 2. In step 2 the $g(n,N)$ as well as the relative values $v_{(n,N)}(s)$, $s \in S$ are computed for a given (n,N) -policy. First we note that apart from the set of equations (2.2) the $g(n,N)$ and $v_{(n,N)}(s)$, $s \in E$ are also the unique solution to the embedded set (4.5). For a proof of this result we refer to Tijms [7] (pp. 229-230).

So we first solve (4.5). At this point it is worthwhile to note that (4.5) becomes a rather small set of only n equations with n unknowns due the special structure of our problem. To prove this we note that

$$v(i,m+1) = b + r_2 - g \tau(i,m+1) + \sum_{t \in E} p((i,m+1),t) v(t), \quad 1 \leq i < n$$

and

$$v(i,N) = r_2 - g \tau(i,N) + \sum_{t \in E} p((i,N),t) v(t), \quad 1 \leq i < n$$

while

$$(5.3) \quad v(i,m+1) = b + r_{12} - g \tau(i,m+1) + \sum_{t \in E} p((i,m+1),t) v(t), \quad n \leq i \leq N$$

and

$$v(i,N) = r_{12} - g \tau(i,N) + \sum_{t \in E} p((i,N),t) v(t), \quad n \leq i \leq N.$$

From (4.2), (4.3) and (4.4) we conclude that

$$v(i,m+1) = b + v(i,N) \quad \text{for all } 1 \leq i \leq N$$

Moreover we see from (4.3), (4.4) and the fact that $p((i,N), s)$ for $i \geq n$ is independent of i that the right hand side of (5.3) is also independent of i .

This, together with the symmetry of $v(.,.)$ (see proof of theorem 1) reduces (4.5) to a set of n equations.

Next the values of $v_{(n,N)}(s)$ for $s \in E^C$ can be computed by single pass calculations from $g(n,N)$ and $\{v_{(n,N)}(s) : s \in E\}$ as follows. Deleting in the notation the dependency on the (n,N) -policy we have

$$v(i,m+1) = b + r_1 - g + w(i,0) = b + v(i,j), \quad 1 \leq i < n; N \leq j \leq m$$

and

$$v(i,m+1) = b + r_{12} - g + w(0,0) = b + v(i,j), \quad n \leq i \leq N; N \leq j \leq m$$

Hence we conclude that

$$v(i,j) = v(i,m+1) - b \quad \text{for all } 1 \leq i \leq N; N \leq j \leq m.$$

$\{(n-1, j), N \leq j \leq m+1\}$ from 2 into 12 and in the states $\{(i, n-1), N \leq i \leq m+1\}$ from 1 into 12. In all other states no change of action occurs. Hence we compute the test value $T_{(n,N)}(s, 12)$ (see (2.3)) for

$$s \in \{(n-1, j), N \leq j \leq m+1\} \cup \{(i, n-1), N \leq i \leq m+1\}$$

which yields:

$$T_{(n,N)}((n-1, j), 12) = r_{12} - g + w(0, 0)$$

while

$$v_{(n,N)}(n-1, j) = r_2 - g + w(n-1, 0)$$

($w(\cdot, \cdot)$ is defined by (3.10))

Since neither $T_{(n,N)}((n-1, j), 12)$ nor $v_{(n,N)}(n-1, j)$ depends on j we conclude that either

$$(5.4) \quad T_{(n,N)}((n-1, j), 12) < v_{(n,N)}(n-1, j) \quad \text{for all } N \leq j \leq m+1$$

or

$$(5.5) \quad T_{(n,N)}((n-1, j), 12) \geq v_{(n,N)}(n-1, j) \quad \text{for all } N \leq j \leq m+1.$$

If (5.4) holds $(n-1, N)$ yields an improvement over the policy (n, N) . If, on the contrary (5.5) is satisfied we conclude that the policy (n, N) is not improved by $(n-1, N)$. So we can compare the policies (n, N) and $(n-1, N)$ without computation of $g(n-1, N)$. If $(n-1, N)$ yields an improvement over (n, N) we continue by comparing the policy (n, N) with $(n-2, N)$ and continue in this way until no improvement is obtained any more. On the other hand if (5.5) holds we compare (n, N) with $(n+1, N)$ in a similar way.

Next we consider possible improvements of the value of N , with n fixed. This turns out to be more delicate than changing n . A modification of (n, N) into $(n, N+1)$ yields a change of action on the set $\{(i, N) : 1 \leq i \leq N\} \cup \{(N, j) : 1 \leq j \leq N\}$.

For $1 \leq i \leq N$ we have

$$(5.6) \quad T_{(n,N)}((i,N),0) = -g + w(i,N)$$

while

$$(5.7) \quad v_{(n,N)}(i,N) = \begin{cases} r_1 - g + w(i,0) & \text{for } i < n \\ r_{12} - g + w(0,0) & \text{for } i \geq n \end{cases}$$

From (5.6) and (5.7) we conclude that in general no comparison between $T_{(n,N)}((i,N),0)$ and $v_{(n,N)}(i,N)$ is possible uniformly in $1 \leq i \leq N$. Hence no unambiguous comparison between the policies (n,N) and $(n,N+1)$ can be made based on this policy improvement step. Indeed this ambiguity did occur in our numerical examples. Therefore in our algorithm we compare (n,N) and $(n,N+1)$ directly on base of $g(n,N)$ and $g(n,N+1)$.

The complete algorithm now reads as follows.

Algorithm

Step 0. Determine T^* and choose $n = N = T^*$. Go to step 1.

Step 1. Compute $g(n,N)$ and $v_{(n,N)}(s)$, $s \in E$ and go to step 2.

Step 2. Compute $v_{(n,N)}(s)$, $s \in S$ and go to step 3.

Step 3. Apply the improvement procedure to compare (n,N) and $(n-1,N)$. If improvement is established compare (n,N) with $(n-2,N)$ and continue in this way until no further improvement is obtained. Set n equal to the last value for which improvement was established and go to step 1. If $(n-1,N)$ is not better than (n,N) apply the policy improvement procedure to compare (n,N) and $(n+1,N)$ and proceed analogously.

If neither $(n-1,N)$ nor $(n+1,N)$ is better than (n,N) than go to step 4.

Step 4. Compute $g(n, N+1)$. If $g(n, N+1) < g(n, N)$ then put $N := N+1$ and go to step 1 else compute $g(n, N-1)$. If $g(n, N-1) < g(n, N)$ and $n < N$ then put $N := N-1$ and go to step 1. If $g(n, N-1) < g(n, N)$ and $n=N$ then put $N := N-1$ and $n := n-1$ and go to step 1 else go to step 5.

Step 5. Perform an overall policy improvement step using (2.2) and (2.3) and stop.

The algorithm works quite satisfactorily. In table 2 below the computation times for several numerical examples are compared with those for the value-iteration algorithm. The numerical data used in this table are (5,4,5) and (5,7,10) for (b, r_1, r_{12}) while the lifetime distributions are given by $p^{(i)}$, $i=1, \dots, 5$ as given in the appendix. We have made the comparison with the value-iteration algorithm because this algorithm usually works better on large state spaces than the policy iteration method.

	(5,4,5)	(5,7,10)
$p^{(1)}$	4:18 (18:34)	7:03 (17:41)
$p^{(2)}$	4:18 (47:45)	6:10 (36:45)
$p^{(3)}$	4:88 (39:28)	7:14 (41:25)
$p^{(4)}$	20:10 (14:01)	31:40 (24:01)
$p^{(5)}$	8:09 (3:07:62)	8:62 (1:22:56)

Table 2 Computation times of the algorithm in hundreds of seconds. In brackets are given the computation times according to the value iteration algorithm, yielding the overall optimal policy.

Although our algorithm theoretically can stick to a local minimum this did not occur in our numerical examples.

Note that our algorithm is a hybrid method containing elements from the value iteration method (steps 1 and 2), from the policy iteration method (step 3) as well as brute comparisons (step 4).

Step 1 is in general the most time consuming step, in particular when m is large. Therefore we emphasize that this step has to be carried out only once each passage through step 3.

Step 5 can be carried out as a final step in the algorithm in order to check the final (n,N) -policy upon overall optimality. In most of our numerical experiments we only needed just one step in the overall policy improvement procedure starting at the optimal (n,N) -policy to get the overall optimal policy.

REFERENCES

- [1] Bäckert, W., and Rippin, D.W.T., "The determination of maintenance strategies for plants subject to breakdown", *Computers and Chemical Engineering* 9 (1985), 113-126.
- [2] Epstein, Sheldon, and Wilamowsky, Yonah, "An optimal replacement policy for life limited parts", *Opsearch* 23 (1986), 151-163.
- [3] Haurie, A., and L'Ecuyer, P., "A stochastic control approach to group preventive replacement in a multicomponent system", *IEEE Transactions on Automatic Control* 27 (1982), 387-393.
- [4] Howard, R.A., *Dynamic Programming and Markov Processes*, Wiley, New York, 1960.
- [5] Özekici, Süleyman, "Optimal replacement of one-unit systems under periodic inspection", *Siam J. Control and Optimization* 23 (1985), 122-128.
- [6] Özekici, Süleyman, "Optimal periodic replacement of multicomponent reliability systems", *Operations research* 36 (1988), 542-552.
- [7] Tijms, Henk C., *Stochastic modelling and analysis*, Wiley, Chichester, 1986.
- [8] Tijms, H.C., and Van Der Duyn Schouten, F.A., "Inventory control with two switch-over levels for a class of M/G/1 queueing systems with variable arrival and service rate", *Stochastic Processes* 6 (1978), 213-222.
- [9] Vergin, Roger C., and Scriabin, Michael, "Maintenance scheduling for multicomponent equipment", *AIIE Transactions* 9 (1977), 297-305.

APPENDIX Numerical results

In the tables below we present numerical results for a number of costs parameters (b, r_1, r_{12}) and probability vectors $p^{(i)}$.

In the first column we see whether the optimal strategy has the (n, N) -form or not (y/n) or (n, N) on the recurrent class (y') . From the 4th and 5th column we can make a comparison between the optimal strategy and the strategy $(n, N)^*$, where the 5th column gives the percentage increases of $g(n, N)^*$ over g^* . In addition we mention in the last column the number of times the average cost of a policy is calculated by the heuristic.

$$p^{(1)} = (0.80, 0.80, 0.75, 0.66, 0.55, 0.25, 0.15, 0.10, 0.05, 0.01) \quad (m=10)$$

b	r_1	r_{12}	$(n, N)^*$	T^*	g^*	%	#
5	1	1.6	n (2,3)	3	2.613	0.19	(4)
5	2	3	n (2,4)	4	3.212	0.03	(4)
5	2	4	n (4,4)	4	3.436	0.00	(3)
5	4	5	y' (2,5)	5	4.078	0.00	(4)
5	4	7.5	n (4,5)	5	4.645	0.06	(4)
5	7	8	y (2,11)	5	5.347	0.00	(7)
5	7	10	y' (3,11)	5	5.887	0.00	(6)
5	7	13	y' (5,11)	5	6.399	0.00	(6)
5	12	18	y' (4,11)	5	8.703	0.00	(4)

$$p^{(2)} = (0.96, 0.92, 0.87, 0.77, 0.60, 0.40, 0.31, 0.15, 0.05, 0.05) \quad (m=10)$$

b	r_1	r_{12}		$(n,N)^*$	T^*	g^*	%	#
5	1	1.6	y	(2,3)	3	1.348	0.00	(5)
5	2	3	n	(2,3)	3	1.863	0.21	(5)
5	2	4	n	(3,4)	3	2.160	0.18	(3)
5	4	5	y'	(1,4)	4	2.506	0.00	(4)
5	4	7.5	n	(3,5)	4	3.189	0.00	(4)
5	7	8	y'	(1,5)	5	3.396	0.00	(5)
5	7	10	n	(2,5)	5	3.919	0.02	(5)
5	7	13	n	(4,7)	5	4.587	0.13	(7)
5	12	18	y'	(3,11)	7	5.981	0.00	(5)

$$p^{(3)} = (0.90, 0.90, 0.88, 0.85, 0.65, 0.45, 0.25, 0.12, 0.10, 0.10) \quad (m=10)$$

b	r_1	r_{12}		$(n,N)^*$	T^*	g^*	%	#
5	1	1.6	y	(2,4)	4	1.583	0.00	(4)
5	2	3	y	(2,4)	4	2.045	0.00	(5)
5	2	4	y	(4,4)	4	2.254	0.00	(3)
5	4	5	n	(2,4)	4	2.724	0.51	(5)
5	4	7.5	n	(4,5)	4	3.300	0.27	(3)
5	7	8	n	(2,5)	5	3.655	0.03	(5)
5	7	10	y'	(3,6)	5	4.140	0.00	(5)
5	7	13	n	(4,8)	5	4.713	0.13	(6)
5	12	18	y'	(3,11)	6	6.234	0.00	(6)

$$p^{(4)} = (0.806, 0.703, 0.648, 0.609, 0.578, 0.552, 0.530, 0.510, 0.493, 0.477, 0.463, 0.450, 0.438, 0.427) \quad (m=14)$$

b	r_1	r_{12}		$(n,N)^*$	T^*	g^*	%	#
5	1	1.6	y'	(1,3)	2	3.003	0.00	(4)
5	2	3	y'	(2,7)	5	3.631	0.00	(5)
5	2	4	n	(6,11)	5	3.837	0.00	(9)
5	4	5	y	(2,14)	12	4.573	0.00	(5)
5	4	7.5	n	(11,14)	12	5.029	0.00	(6)
5	7	8	y	(1,15)	14	5.957	0.00	(4)
5	7	10	y	(5,15)	14	6.521	0.00	(5)
5	7	13	n	(14,14)	14	6.833	0.00	(3)
5	12	18	y	(9,15)	15	9.441	0.00	(3)

$$p^{(5)} = (0.995, 0.968, 0.916, 0.843, 0.754, 0.656, 0.555, 0.457, 0.366, 0.285, 0.216, 0.159, 0.114, 0.079) \quad (m=14)$$

b	r_1	r_{12}		$(n,N)^*$	T^*	g^*	%	#
5	1	1.6	y	(2,3)	3	0.928	0.00	(5)
5	2	3	n	(2,3)	4	1.407	0.07	(6)
5	2	4	n	(3,4)	4	1.678	0.48	(4)
5	4	5	y'	(1,4)	5	1.957	0.00	(5)
5	4	7.5	n	(3,5)	5	2.555	0.12	(4)
5	7	8	y'	(1,5)	6	2.689	0.00	(5)
5	7	10	y'	(2,6)	6	3.136	0.00	(5)
5	7	13	n	(4,8)	6	3.728	0.03	(5)
5	12	18	n	(3,10)	8	4.823	0.00	(5)

$p^{(4)}$ and $p^{(5)}$ are obtained from a discretization of the Weibull $(\alpha,1)$ distribution for $\alpha=1.4$ and $\alpha=3$ respectively.

IN 1988 REEDS VERSCHENEN

- 297 Bert Bettonvil
Factor screening by sequential bifurcation
- 298 Robert P. Gilles
On perfect competition in an economy with a coalitional structure
- 299 Willem Selen, Ruud M. Heuts
Capacitated Lot-Size Production Planning in Process Industry
- 300 J. Kriens, J.Th. van Lieshout
Notes on the Markowitz portfolio selection method
- 301 Bert Bettonvil, Jack P.C. Kleijnen
Measurement scales and resolution IV designs: a note
- 302 Theo Nijman, Marno Verbeek
Estimation of time dependent parameters in lineair models
using cross sections, panels or both
- 303 Raymond H.J.M. Gradus
A differential game between government and firms: a non-cooperative
approach
- 304 Leo W.G. Strijbosch, Ronald J.M.M. Does
Comparison of bias-reducing methods for estimating the parameter in
dilution series
- 305 Drs. W.J. Reijnders, Drs. W.F. Verstappen
Strategische bespiegelingen betreffende het Nederlandse kwaliteits-
concept
- 306 J.P.C. Kleijnen, J. Kriens, H. Timmermans and H. Van den Wildenberg
Regression sampling in statistical auditing
- 307 Isolde Woittiez, Arie Kapteyn
A Model of Job Choice, Labour Supply and Wages
- 308 Jack P.C. Kleijnen
Simulation and optimization in production planning: A case study
- 309 Robert P. Gilles and Pieter H.M. Ruys
Relational constraints in coalition formation
- 310 Drs. H. Leo Theuns
Determinanten van de vraag naar vakantiereizen: een verkenning van
materiële en immateriële factoren
- 311 Peter M. Kort
Dynamic Firm Behaviour within an Uncertain Environment
- 312 J.P.C. Blanc
A numerical approach to cyclic-service queueing models

- 313 Drs. N.J. de Beer, Drs. A.M. van Nunen, Drs. M.O. Nijkamp
Does Morkmon Matter?
- 314 Th. van de Klundert
Wage differentials and employment in a two-sector model with a dual labour market
- 315 Aart de Zeeuw, Fons Groot, Cees Withagen
On Credible Optimal Tax Rate Policies
- 316 Christian B. Mulder
Wage moderating effects of corporatism
Decentralized versus centralized wage setting in a union, firm, government context
- 317 Jörg Glombowski, Michael Krüger
A short-period Goodwin growth cycle
- 318 Theo Nijman, Marno Verbeek, Arthur van Soest
The optimal design of rotating panels in a simple analysis of variance model
- 319 Drs. S.V. Hannema, Drs. P.A.M. Versteijne
De toepassing en toekomst van public private partnership's bij de grote en middelgrote Nederlandse gemeenten
- 320 Th. van de Klundert
Wage Rigidity, Capital Accumulation and Unemployment in a Small Open Economy
- 321 M.H.C. Paardekooper
An upper and a lower bound for the distance of a manifold to a nearby point
- 322 Th. ten Raa, F. van der Ploeg
A statistical approach to the problem of negatives in input-output analysis
- 323 P. Kooreman
Household Labor Force Participation as a Cooperative Game; an Empirical Model
- 324 A.B.T.M. van Schaik
Persistent Unemployment and Long Run Growth
- 325 Dr. F.W.M. Boekema, Drs. L.A.G. Oerlemans
De lokale produktiestructuur doorgelicht.
Bedrijfstakverkenningen ten behoeve van regionaal-economisch onderzoek
- 326 J.P.C. Kleijnen, J. Kriens, M.C.H.M. Lafleur, J.H.F. Pardoel
Sampling for quality inspection and correction: AOQL performance criteria

- 327 Theo E. Nijman, Mark F.J. Steel
Exclusion restrictions in instrumental variables equations
- 328 B.B. van der Genugten
Estimation in linear regression under the presence of heteroskedasticity of a completely unknown form
- 329 Raymond H.J.M. Gradus
The employment policy of government: to create jobs or to let them create?
- 330 Hans Kremers, Dolf Talman
Solving the nonlinear complementarity problem with lower and upper bounds
- 331 Antoon van den Elzen
Interpretation and generalization of the Lemke-Howson algorithm
- 332 Jack P.C. Kleijnen
Analyzing simulation experiments with common random numbers, part II: Rao's approach
- 333 Jacek Osiewalski
Posterior and Predictive Densities for Nonlinear Regression. A Partly Linear Model Case
- 334 A.H. van den Elzen, A.J.J. Talman
A procedure for finding Nash equilibria in bi-matrix games
- 335 Arthur van Soest
Minimum wage rates and unemployment in The Netherlands
- 336 Arthur van Soest, Peter Kooreman, Arie Kapteyn
Coherent specification of demand systems with corner solutions and endogenous regimes
- 337 Dr. F.W.M. Boekema, Drs. L.A.G. Oerlemans
De lokale produktiestructuur doorgelicht II. Bedrijfstakverkenningen ten behoeve van regionaal-economisch onderzoek. De zeescheepsnieuwbouwindustrie
- 338 Gerard J. van den Berg
Search behaviour, transitions to nonparticipation and the duration of unemployment
- 339 W.J.H. Groenendaal and J.W.A. Vingerhoets
The new cocoa-agreement analysed
- 340 Drs. F.G. van den Heuvel, Drs. M.P.H. de Vor
Kwantificering van ombuigen en bezuinigen op collectieve uitgaven 1977-1990
- 341 Pieter J.F.G. Meulendijks
An exercise in welfare economics (III)

- 342 W.J. Selen and R.M. Heuts
A modified priority index for Günther's lot-sizing heuristic under capacitated single stage production
- 343 Linda J. Mittermaier, Willem J. Selen, Jeri B. Waggoner, Wallace R. Wood
Accounting estimates as cost inputs to logistics models
- 344 Remy L. de Jong, Rashid I. Al Layla, Willem J. Selen
Alternative water management scenarios for Saudi Arabia
- 345 W.J. Selen and R.M. Heuts
Capacitated Single Stage Production Planning with Storage Constraints and Sequence-Dependent Setup Times
- 346 Peter Kort
The Flexible Accelerator Mechanism in a Financial Adjustment Cost Model
- 347 W.J. Reijnders en W.F. Verstappen
De toenemende importantie van het verticale marketing systeem
- 348 P.C. van Batenburg en J. Kriens
E.O.Q.L. - A revised and improved version of A.O.Q.L.
- 349 Drs. W.P.C. van den Nieuwenhof
Multinationalisatie en coördinatie
De internationale strategie van Nederlandse ondernemingen nader beschouwd
- 350 K.A. Bubshait, W.J. Selen
Estimation of the relationship between project attributes and the implementation of engineering management tools
- 351 M.P. Tummers, I. Woittiez
A simultaneous wage and labour supply model with hours restrictions
- 352 Marco Versteijne
Measuring the effectiveness of advertising in a positioning context with multi dimensional scaling techniques
- 353 Dr. F. Boekema, Drs. L. Oerlemans
Innovatie en stedelijke economische ontwikkeling
- 354 J.M. Schumacher
Discrete events: perspectives from system theory
- 355 F.C. Bussemaker, W.H. Haemers, R. Mathon and H.A. Wilbrink
A (49,16,3,6) strongly regular graph does not exist
- 356 Drs. J.C. Caanen
Tien jaar inflatieneutrale belastingheffing door middel van vermogensaftrek en voorraadaftrek: een kwantitatieve benadering

- 357 R.M. Heuts, M. Bronckers
A modified coordinated reorder procedure under aggregate investment
and service constraints using optimal policy surfaces
- 358 B.B. van der Genugten
Linear time-invariant filters of infinite order for non-stationary
processes
- 359 J.C. Engwerda
LQ-problem: the discrete-time time-varying case
- 360 Shan-Hwei Nienhuys-Cheng
Constraints in binary semantical networks
- 361 A.B.T.M. van Schaik
Interregional Propagation of Inflationary Shocks
- 362 F.C. Drost
How to define UVMU
- 363 Rommert J. Casimir
Infogame users manual
Rev 1.2 December 1988
- 364 M.H.C. Paardeckooper
A quadratically convergent parallel Jacobi-process for diagonal
dominant matrices with nondistinct eigenvalues
- 365 Robert P. Gilles, Pieter H.M. Ruys
Characterization of Economic Agents in Arbitrary Communication
Structures
- 366 Harry H. Tigelaar
Informative sampling in a multivariate linear system disturbed by
moving average noise
- 367 Jörg Glombowski
Cyclical interactions of politics and economics in an abstract
capitalist economy

IN 1989 REEDS VERSCHENEN

- 368 Ed Nijssen, Will Reijnders
"Macht als strategisch en tactisch marketinginstrument binnen de
distributieketen"
- 369 Raymond Gradus
Optimal dynamic taxation with respect to firms
- 370 Theo Nijman
The optimal choice of controls and pre-experimental observations
- 371 Robert P. Gilles, Pieter H.M. Ruys
Relational constraints in coalition formation

Bibliotheek K. U. Brabant

17 000 01086008 9