Tilburg University

Controlled communication networks

Feltkamp, V.; van den Nouweland, C.G.A.M.

Publication date:
1992

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Feltkamp, V., & van den Nouweland, C. G. A. M. (1992). *Controlled communication networks*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 538). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

ECBM
R

7626
1992
538

UNIVERSITY
LIKE
UNIVERSITEIT
BRABANT

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS

DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM

CONTROLLED COMMUNICATION NETWORKS

V. Feltkamp
A. van den Nouweland

FEW 538

R40
Game Theory

Refereed by Prof.dr. S.H. Tijs

K.U.B.
BIBLIOTHEEK
TILBURG

Controlled Communication Networks

By V. Feltkamp* and A. van den Nouweland*

Abstract

This paper considers a generalisation of communication situations, namely controlled communication networks. Three solution concepts for controlled communication networks are introduced, the Myerson value, the position value and the mixed value, which are inspired by corresponding solution concepts for communication situations, and axiomatic characterisations of these concepts are provided. Further, network games associated with controlled communication networks are considered and it is shown that every TU -game can be obtained as a network game.

*Department of Econometrics, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands

1 Introduction and Preliminaries

In a cooperative *TU* (Transferable Utility) game (N, v) as modelled by von Neumann & Morgenstern (1944), N is a finite set of players, and the characteristic function v assigns to each subgroup of players a real number which is to be interpreted as the maximal gains (minimal costs) this coalition can achieve by cooperating, regardless of the actions of the other players.

It is generally assumed that each subgroup of players can form and cooperate to obtain its value. However, this approach fails to take into account communication restrictions that may cause deficiencies in cooperation in some coalitions. Myerson (1977) introduced communication graphs to model non-transitive communication restrictions. In such a graph the vertices are the players and an edge between two players represents the fact that these players can communicate directly. The general procedure is the following: given a *TU*-game (N, v) and a communication graph (N, A) , one defines a reward function r on collections of vertices and edges which takes the communication restrictions into account, and then new games are extracted. This model was elaborated further by Owen (1986), Borm, Owen & Tijs (1992), Van den Nouweland, Borm, Owen and Tijs (1991) and Van den Nouweland & Borm (1991). A survey on this subject is given in Borm, Van den Nouweland & Tijs (1991).

In this paper, we generalise communication situations by allowing the vertex set of the graph to differ from the set of players N , and by starting out with a reward function r , instead of deriving it. This means that we can assign different rewards to different connected graphs with the same vertex set. We assume that the players control the vertices and edges of the graph through so-called control games. In section 2 we provide the formal definition of controlled communication networks and we introduce three solution concepts for these networks, the Myerson value, the position value and the mixed value. These solution concepts are characterised axiomatically in section 3. Finally, in section 4 we present an alternative way of constructing a *TU*-game corresponding to a controlled communication network, and we show that all *TU*-games can be obtained as such a network game.

Preliminaries

Let $N := \{1, \dots, n\}$, and $2^N := \{S \mid S \subseteq N\}$. We denote the class of all transferable utility games with player set N by TU^N .

For each $S \in 2^N \setminus \{\emptyset\}$, the unanimity game $(N, u_S) \in TU^N$ is defined by

$$u_S(T) = \begin{cases} 1 & \text{if } S \subseteq T \\ 0 & \text{otherwise} \end{cases}$$

for all $T \subseteq N$.

It is well known that $\{(N, u_S) \mid S \in 2^N \setminus \{\emptyset\}\}$ forms a basis of TU^N .

The Shapley value Φ (cf. Shapley, 1953) is a linear function $TU^N \rightarrow \mathbb{R}^N$ defined by

$$\Phi_i(N, u_S) = \begin{cases} \frac{1}{|S|} & \text{if } i \in S \\ 0 & \text{if } i \in N \setminus S \end{cases}$$

for all $S \in 2^N \setminus \{\emptyset\}$.

2 Controlled Communication Networks

Consider a finite undirected graph (V, E) without loops or parallel edges. We assume that for each vertex $v \in V$ a *simple control game* (N, c_v) with *veto players* is given and, similarly, for each edge $e \in E$ a simple control game (N, c_e) with veto players is given. Here, a TU -game (N, c) is called *simple* if $c(S) \in \{0, 1\}$ for all $S \subseteq N$ and $c(N) = 1$. Moreover, a player $i \in N$ is called a *veto player* of the simple game (N, c) , denoted by $i \in \text{veto}(c)$, if i can block a positive outcome, i.e. $c(S) = 0$ for all $S \in 2^N \setminus \{\emptyset\}$ with $i \notin S$. If c_v is the control game for vertex $v \in V$, then a coalition $S \subseteq N$ is allowed to use vertex v if and only if $c_v(S) = 1$. The control games (N, c_e) for edges have a similar interpretation.

Furthermore, we assume that there is a reward function r on subsets of vertices and edges $r : 2^V \times 2^E \rightarrow \mathbb{R}$, measuring the economic value of subnetworks. Keeping in mind that edges are to model communication channels, it seems reasonable to assume that an edge is useless without both its end points, i.e. for all $W \subseteq V$ and $F \subseteq E$ it holds that $r(W, F) = r(W, F \setminus \{\{v_1, v_2\}\})$ if $\{v_1, v_2\} \in F$ is such that $\{v_1, v_2\}$ is not a subset of W . So, the reward of a network (W, F) does not depend on the edges not in $F(W) := \{\{v, w\} \in F \mid v \in W, w \in W\}$. Moreover, in a network (W, F) , W is partitioned into *communication components* in the following way: $C \subseteq W$ is a component within (W, F) if and only if $(C, F(C))$ is a connected subgraph of $(W, F(W))$ and is maximal with respect to this property. The resulting partition of W is denoted W/F .

Correspondingly, we assume that the reward function is additive with respect to these components, i.e.

$$r(W, F) = \sum_{C \in W/F} r(C, F(C))$$

for all $W \subseteq V$ and all $F \subseteq E$.

For simplicity, we assume that r is *zero-normalised*, i.e. $r(\{v\}, \emptyset) = 0$ for all $v \in V$.

A *controlled communication network* is a 6-tuple $\langle N, V, E, \{c_v | v \in V\}, \{c_e | e \in E\}, r \rangle$ as described above. The set of all controlled communication networks with player set N will be denoted CCN^N .

The purpose of this paper is to present and characterize some solutions to the problem of how to distribute the reward $r(V, E)$ among the players in N . Formally, a *solution concept* on CCN^N is a function $\gamma : CCN^N \rightarrow \mathbf{R}^N$ assigning $\gamma_i(C)$ to player i in the controlled communication network $C \in CCN^N$. One way to obtain solution concepts on CCN^N is to construct for each controlled communication network a *TU-game* corresponding to this network in which the players are the edges and/or vertices of the graph. To this game one can apply a solution concept from cooperative game theory, for example the Shapley value. This yields a value for edges and/or vertices and this value can be distributed among the players according to veto control. Concentrating on vertices this procedure yields the Myerson value (cf. Myerson, 1977) and concentrating on edges it yields the position value (cf. Borm, Owen, and Tijs, 1992). If no such distinction between vertices and edges is made, one obtains the mixed value.

Let $C = \langle N, V, E, \{c_v | v \in V\}, \{c_e | e \in E\}, r \rangle$ be a controlled communication network. Then the *Myerson value* $\mu(C) \in \mathbf{R}^N$ is defined by

$$\mu_i(C) := \sum_{v \in V : i \in \text{veto}(c_v)} \frac{\Phi_v(V, r_E)}{|\text{veto}(c_v)|}$$

for all $i \in N$, where the *vertex game* (V, r_E) is a game in which the vertices are the players, defined by $r_E(W) = r(W, E)$ for all $W \subseteq V$.

Further, the *position value* $\pi(C) \in \mathbf{R}^N$ is defined by

$$\pi_i(C) := \sum_{e \in E : i \in \text{veto}(c_e)} \frac{\Phi_e(E, r_V)}{|\text{veto}(c_e)|}$$

for all $i \in N$, where the edge game (E, r_V) is a game in which the edges are the players, defined by $r_V(F) = r(V, F)$ for all $F \subseteq E$.

Finally, the mixed value $\rho(\mathcal{C}) \in \mathbf{R}^N$ is defined by

$$\rho_i(\mathcal{C}) := \sum_{j \in V \cup E: i \in \text{veto}(c_j)} \frac{\Phi_j(V \cup E, \tilde{r})}{|\text{veto}(c_j)|}$$

for all $i \in N$, where the game $(V \cup E, \tilde{r})$ is defined by $\tilde{r}(J) = r(J \cap V, J \cap E)$ for all $J \subseteq V \cup E$.

3 Axiomatic characterisations

In this section we provide axiomatic characterisations of the three solution concepts introduced in section 2. It will turn out that all three concepts can be characterised by four axioms, three of them being the same for all three solution concepts.

For concision, if $\mathcal{C}$ is a controlled communication network and $F \subseteq E$ a set of edges we will use $\mathcal{C}^{-F}$ to denote the network where the edges in F have been omitted, and where the reward function has been restricted accordingly. We now introduce some properties. A solution concept γ on CCN^N is called *efficient* if for each controlled communication network $\mathcal{C}$, γ distributes exactly $r(V, E)$ among the players. In formula:

$$\sum_{i \in N} \gamma_i(\mathcal{C}) = r(V, E).$$

A solution concept γ on CCN^N is called *additive* if it is additive with respect to the reward function (*ceteris paribus*).

A solution concept γ on CCN^N is said to have the *superfluous edge property* if for all $\mathcal{C} \in CCN^N$ and all edges $e \in E$ that are superfluous for $\mathcal{C}$ it holds that

$$\gamma(\mathcal{C}) = \gamma(\mathcal{C}^{-\{e\}}).$$

Here, an edge $e \in E$ is called *superfluous* for $\mathcal{C}$ if for all $F \subseteq E$,

$$r(V, F) = r(V, F \setminus \{e\}).$$

Note that for an edge e to be superfluous, we only consider the whole set of vertices V ; we do not demand $r(W, F) = r(W, F \setminus \{e\})$ for all $W \subseteq V$ and $F \subseteq E$. However, this turns out to be an equivalent demand.

Lemma 1 Let $\mathcal{C} \in CCN^N$. Then an edge $e \in E$ is superfluous for $\mathcal{C}$ if and only if

$$r(W, F) = r(W, F \setminus \{e\})$$

for all $W \subseteq V$ and all $F \subseteq E$.

Proof. The "if" part is straightforward. For the "only if" part, note that

$$\begin{aligned} r(W, F) &= \sum_{C \in W/F} r(C, F(C)) \\ &= \sum_{C \in W/F(W)} r(C, F(C)) \\ &\stackrel{(*)}{=} \sum_{C \in V/F(W)} r(C, F(C)) = r(V, F(W)) \end{aligned}$$

for all $F \subseteq E$ and $W \subseteq V$.

Here, equality (*) follows from the fact that r is zero-normalised.

Hence, for a superfluous edge e

$$\begin{aligned} r(W, F \setminus \{e\}) &= r(V, (F \setminus \{e\})(W)) \\ &= r(V, F(W) \setminus \{e\}) \\ &= r(V, F(W)) \\ &= r(W, F) \end{aligned}$$

for all $F \subseteq E$ and $W \subseteq V$. This completes the proof. $\square$

In a graph (V, E) , we denote by $D(V, E)$ the set of vertices that have at least one neighbour in the graph, and we will shorten this notation to D whenever this does not lead to confusion.

The fourth property of solution concepts on CCN^N we introduce is anonymity. A controlled communication network is said to be *anonymous* if the reward function only depends on the number of non-isolated vertices and edges, i.e. there exists a function $f : \{0, \dots, |D \cup E|\} \rightarrow \mathbb{R}$ such that

$$r(W, F) = f(|(W \cap D) \cup F|)$$

for all $W \subseteq V$ and $F \subseteq E$.

A solution concept γ on CCN^N satisfies *anonymity* if for all anonymous $\mathcal{C} \in CCN^N$, the solution is proportional to the veto power of the players over the non-isolated parts of the graph or, in formula: there exists an $\alpha \in \mathbb{R}$ such that for all $i \in N$

$$\gamma_i(\mathcal{C}) = \alpha \cdot \sum_{j \in D \cup E : i \in \text{veto}(c_j)} \frac{1}{|\text{veto}(c_j)|}$$

The mixed value ρ satisfies the four properties mentioned. This is shown in

Lemma 2 The mixed value ρ satisfies efficiency, additivity, anonymity and the superfluous edge property.

Proof. Let $\mathcal{C}$ be a controlled communication network. Then

$$\begin{aligned} \sum_{i \in N} \rho_i(\mathcal{C}) &= \sum_{i \in N} \sum_{j \in V \cup E : i \in \text{veto}(c_j)} \frac{\Phi_j(V \cup E, \tilde{r})}{|\text{veto}(c_j)|} \\ &= \sum_{j \in V \cup E} \Phi_j(V \cup E, \tilde{r}) \cdot \sum_{i \in \text{veto}(c_j)} \frac{1}{|\text{veto}(c_j)|} \\ &= \sum_{j \in V \cup E} \Phi_j(V \cup E, \tilde{r}) \stackrel{(*)}{=} r(V, E), \end{aligned}$$

where equality (*) follows from efficiency of the Shapley value Φ . Hence, ρ is efficient.

Additivity of ρ follows straightforward by using additivity for Φ .

In order to prove the superfluous edge property, take $\mathcal{C} \in CCN^N$ and $e \in E$ that is superfluous for $\mathcal{C}$. It clearly suffices to prove that $\Phi_e(V \cup E, \tilde{r}) = 0$ and $\Phi_j(V \cup E, \tilde{r}) = \Phi_j(V \cup E \setminus \{e\}, \tilde{r})$ for all $j \in V \cup E \setminus \{e\}$. Using Lemma 1 we easily obtain $\tilde{r}(J) = \tilde{r}(J \setminus \{e\})$ for any $J \subseteq V \cup E$. Hence, e is a zero player in the game $(V \cup E, \tilde{r})$, and consequently $\Phi_e(V \cup E, \tilde{r}) = 0$ and $\Phi_j(V \cup E, \tilde{r}) = \Phi_j(V \cup E \setminus \{e\}, \tilde{r})$ for all $j \in V \cup E \setminus \{e\}$. We

conclude that ρ satisfies the superfluous edge property.

Now let $\mathcal{C} \in CCN^N$ be anonymous, and let $f : \{0, \dots, |D \cup E|\} \rightarrow \mathbf{R}$ be such that $r(W, F) = f(|(D \cap W) \cup F|)$ for all $F \subseteq E$ and $W \subseteq V$. Then all vertices $v \in V \setminus D$ are zero players in the game $(V \cup E, \tilde{r})$, and all $j \in D \cup E$ are symmetric in this game. By symmetry, efficiency and the dummy property of Φ this implies

$$\Phi_j(V \cup E, \tilde{r}) = \begin{cases} \frac{f(|D \cup E|)}{|D \cup E|} & \text{if } j \in D \cup E \\ 0 & \text{if } j \in V \setminus D. \end{cases}$$

Hence, $\rho_j(\mathcal{C}) = \alpha \cdot \sum_{j \in D \cup E : i \in \text{veto}(c_j)} \frac{1}{|\text{veto}(c_j)|}$, where $\alpha := \frac{f(|D \cup E|)}{|D \cup E|}$. □

Before we prove that ρ is characterised by the four properties, we introduce two more definitions.

Let (V, E) be a graph. Then we denote by $\mathcal{R}(V, E)$ the set of (V, E) -admissible reward functions, i.e.

$$\mathcal{R}(V, E) := \{r : 2^V \times 2^E \rightarrow \mathbf{R} \mid r \text{ is additive w.r.t. components and zero-normalised}\}.$$

Moreover,

$$\begin{aligned} \mathcal{B}(V, E) := \{ & u_{W,F} : 2^V \times 2^E \rightarrow \mathbf{R} \mid (W, F) \text{ is a connected subgraph of } (V, E), \\ & \text{and } |W| \geq 2\}, \end{aligned}$$

where $u_{W,F}$ is defined by

$$u_{W,F}(W', F') := \begin{cases} 1 & \text{if } W \subseteq W' \text{ and } F \subseteq F' \\ 0 & \text{otherwise.} \end{cases}$$

Then we have the following

Lemma 3 Let (V, E) be a fixed graph. Then $\mathcal{B}(V, E)$ forms a basis of the vector space $\mathcal{R}(V, E)$.

This lemma is an easy corollary of the fact that the set of unanimity games $\{(N, u_S) \mid S \in 2^N \setminus \{\emptyset\}\}$ forms a basis of TU^N , and that the reward functions are zero-normalized. However, the proof is rather technical, and therefore we omit it.

Theorem 4 The mixed value ρ is the unique solution concept on CCN^N satisfying efficiency, additivity, anonymity and the superfluous edge property.

Proof. According to Lemma 2 ρ satisfies the four properties. Hence, we only have to show that there is at most one solution concept satisfying these properties. Suppose γ is a solution concept on CCN^N that satisfies the four properties. Using lemma 3 and additivity of γ and ρ , we see that it suffices to prove $\gamma = \rho$ for situations in which $r = \beta u_{W,F}$ for some $\beta \in \mathbb{R}$ and some connected subgraph (W, F) with $|W| \geq 2$. Hence, let $\mathcal{C} \subseteq CCN^N$ be a controlled communication network with $r = \beta u_{W,F}$ for some $\beta \in \mathbb{R}$ and some connected subgraph (W, F) with $|W| \geq 2$. Since every edge e in $E \setminus F$ is superfluous for $\mathcal{C}$, the superfluous edge property implies that $\gamma(\mathcal{C}) = \gamma(\mathcal{C}^{-E \setminus F})$. Furthermore,

$$r(W', F') = \beta u_{W,F}(W', F') = \begin{cases} \beta & \text{if } W \subseteq W' \text{ and } F' = F \\ 0 & \text{otherwise} \end{cases}$$

for all $W' \subseteq V$ and $F' \subseteq F$.

Since (W, F) is a connected graph, it holds that $D(V, F) = W$. So, defining $f : \{0, \dots, |D \cup E|\} \rightarrow \mathbb{R}$ by

$$f(k) = \begin{cases} \beta & \text{if } k = |W \cup F| \\ 0 & \text{otherwise} \end{cases}$$

we see that the controlled communication network $\mathcal{C}^{-E \setminus F}$ is anonymous. Now, by anonymity of γ we know that there exists an $\alpha \in \mathbb{R}$ such that

$$\gamma_i(\mathcal{C}^{-E \setminus F}) = \alpha \cdot \sum_{j \in W \cup F : i \in \text{veto}(c_j)} \frac{1}{|\text{veto}(c_j)|}$$

for all $i \in N$.

Using efficiency, we obtain

$$\beta = r(V, F) = \sum_{i \in N} \gamma_i(\mathcal{C}^{-E \setminus F}) = \sum_{i \in N} \alpha \cdot \sum_{j \in W \cup F : i \in \text{veto}(c_j)} \frac{1}{|\text{veto}(c_j)|} = \alpha \cdot |W \cup F|.$$

Hence, $\alpha = \beta \cdot |W \cup F|^{-1}$ and recalling $\gamma(\mathcal{C}) = \gamma(\mathcal{C}^{-E \setminus F})$, we see

$$\gamma_i(\mathcal{C}) = \frac{\beta}{|W \cup F|} \cdot \sum_{j \in W \cup F: i \in \text{veto}(c_j)} \frac{1}{|\text{veto}(c_j)|} = \rho_i(\mathcal{C}).$$

□

We proceed by providing axiomatic characterisations of both the Myerson value and the position value. Both values can be characterised by efficiency, additivity, the superfluous edge property and an anonymity axiom. The anonymity axioms we need are *vertex anonymity* and *edge anonymity*.

A solution concept γ on CCN^N is said to be *vertex anonymous* if for every controlled communication network $\mathcal{C} \in CCN^N$ such that there exists a function $f: \{0, \dots, |D(V, E)|\} \rightarrow \mathbb{R}$ with $r(W, E) = f(|D \cap W|)$ for all $W \subseteq V$, there is an $\alpha \in \mathbb{R}$ such that for all $i \in N$

$$\gamma_i(\mathcal{C}) = \alpha \cdot \sum_{v \in D: i \in \text{veto}(c_v)} |\text{veto}(c_v)|^{-1}.$$

A solution concept γ on CCN^N is called *edge anonymous* if for every $\mathcal{C} \in CCN^N$ such that there exists a function $f: \{0, \dots, |E|\} \rightarrow \mathbb{R}$ with $r(V, F) = f(|F|)$ for all $F \subseteq E$, there is an $\alpha \in \mathbb{R}$ such that for all i in N

$$\gamma_i(\mathcal{C}) = \alpha \cdot \sum_{e \in E: i \in \text{veto}(c_e)} |\text{veto}(c_e)|^{-1}.$$

Theorem 5

- i) The Myerson value μ is the unique solution concept on CCN^N that satisfies efficiency, additivity, the superfluous edge property and vertex anonymity.
- ii) The position value π is the unique solution concept on CCN^N that satisfies efficiency, additivity, the superfluous edge property and edge anonymity.

The proof of theorem 5 runs along the same lines as the proof of lemma 2 and theorem 4 and therefore it is left to the reader.

Remark. The axiomatic characterisations of the Myerson value and the position value provided in theorem 5 are similar to axiomatic characterisations provided in Borm, Owen

and Tijs (1992) for *TU*-communication situations. However, the reader should note that our characterisations hold for all controlled communication networks whereas Borm, Owen and Tijs (1992) had to restrict to cycle-free communication graphs.

4 Network games

In the previous sections we approached the problem how to divide the reward $r(V, E)$ of a controlled communication network amongst the players in an indirect way, by first determining the value of vertices and edges and then distributing these values among the veto players in the corresponding control games. In this section we will describe a direct way of dealing with the problem.

Let $\mathcal{C}$ be a controlled communication network. Now we define an associated game with player set N in the following way:

Let $S \subseteq N$. Then $V(S) := \{v \in V \mid c_v(S) = 1\}$ is the set of all vertices that coalition S can control, and $E(S) := \{e \in E \mid c_e(S) = 1\}$ is the set of edges that coalition S can control. Correspondingly, coalition S can obtain

$$v_{\mathcal{C}}(S) := r(V(S), E(S)).$$

Hence, we associate with $\mathcal{C} \in CCN^N$ the network game $(N, v_{\mathcal{C}})$ as defined above. Consequently, some solution concept for *TU*-games could be applied to the game $(N, v_{\mathcal{C}})$.

This approach seems interesting, because a number of games associated with economic situations can be seen to be network games in a more or less natural way. Some examples are sequencing games (Curiel, Pederzoli & Tijs, 1989), permutation games (Tijs *et al.*, 1984) and assignment games (Shapley & Shubik, 1972).

However, some scepticism is in place here, because every *TU*-game is a network game in a trivial way:

Let (N, v) be an arbitrary *TU*-game. We proceed by defining a controlled communication network corresponding to (N, v) . Let $V := N$ and $E := \{\{i, j\} \mid i, j \in N, i \neq j\}$. Hence, (V, E) is the complete graph with vertex set N . The control game for each vertex $i \in N$ is $(N, u_{\{i\}})$ and the control game for each edge $\{i, j\}$ is $(N, u_{\{i, j\}})$. Hence, every player is a dictator for his own vertex and an edge between two vertices is controlled by

the two players it connects. Finally, the reward function r assigns $v(S)$ to the subgraph $(S, E(S))$ for all $S \subseteq N$, and is extended in some feasible way to all subsets of vertices and edges. It is easily seen that the network game associated with the controlled communication network described above is the game (N, v) .

Note that in the above discussion we did not restrict to zero-normalised games. However, the restrictions to zero-normalised reward functions was only made for simplicity and is not essential. Hence, we do not have to worry about it here.

References

- Borm, P., Owen, G., and Tijs, S.H. (1992). "On the position value for communication situations", to appear in *SIAM J. Discr. Math.*
- Borm, P., Nouweland, A. van den, and Tijs, S.H. (1991). "Cooperation and communication restrictions: a survey", in: 'Economic Behaviour in an Imperfect Environment', Gilles, R.P. and Ruys, P.H.M. (Eds.), North-Holland, Amsterdam. (To appear)
- Curiel, I., Pederzoli, G., and Tijs, S.H. (1989). "Sequencing games", *Eur. J. Oper. Res.* 40, 344-351
- Myerson, R.B. (1977). "Graphs and cooperation in games", *Math. Oper. Res.* 2, 225-229.
- Neumann, J. von and Morgenstern, O. (1944). "Theory of Games and Economic Behavior" (3rd edition, 1953), Princeton University Press, Princeton.
- Nouweland, A. van den and Borm, P. (1991). "On the convexity of communication games", *Int. J. Game Th.* 19, 421-430.
- Nouweland, A. van den, Borm, P., Owen, G., and Tijs, S.H. (1991). "Cost allocation and communication", *Res. Mem. FEW 469*, Dept. of Econometrics, Tilburg University, The Netherlands.
- Owen, G. (1986). "Values of graph-restricted games", *SIAM J. Alg. Discr. Meth.* 7, 210-220.
- Shapley, L.S. (1953). "A value for n-person games", in: *Contributions to the Theory of Games, II* (Eds. Tucker, A.W. and Kuhn, H.), *Ann. Math. Studies* 28, Princeton University Press, 307-317.
- Shapley, L.S. and Shubik, M. (1972). "The assignment game I: the core", *Int. J. Game Th.* 1, 111-130.
- Tijs, S.H., Parthasarathy, T., Potters, J., and Rajendra Prasad, V. (1984). "Permutation games: another class of totally balanced games", *OR Spectrum* 6, 119-123.

IN 1991 REEDS VERSCHENEN

- 466 Prof.Dr. Th.C.M.J. van de Klundert - Prof.Dr. A.B.T.M. van Schaik
Economische groei in Nederland in een internationaal perspectief
- 467 Dr. Sylvester C.W. Eijffinger
The convergence of monetary policy - Germany and France as an example
- 468 E. Nijssen
Strategisch gedrag, planning en prestatie. Een inductieve studie binnen de computerbranche
- 469 Anne van den Nouweland, Peter Borm, Guillermo Owen and Stef Tijs
Cost allocation and communication
- 470 Drs. J. Grazell en Drs. C.H. Veld
Motieven voor de uitgifte van converteerbare obligatieleningen en warrant-obligatieleningen: een agency-theoretische benadering
- 471 P.C. van Batenburg, J. Kriens, W.M. Lammerts van Bueren and R.H. Veenstra
Audit Assurance Model and Bayesian Discovery Sampling
- 472 Marcel Kerkhofs
Identification and Estimation of Household Production Models
- 473 Robert P. Gilles, Guillermo Owen, René van den Brink
Games with Permission Structures: The Conjunctive Approach
- 474 Jack P.C. Kleijnen
Sensitivity Analysis of Simulation Experiments: Tutorial on Regression Analysis and Statistical Design
- 475 C.P.M. van Hoesel
An $O(n \log n)$ algorithm for the two-machine flow shop problem with controllable machine speeds
- 476 Stephan G. Vanneste
A Markov Model for Opportunity Maintenance
- 477 F.A. van der Duyn Schouten, M.J.G. van Eijs, R.M.J. Heuts
Coordinated replenishment systems with discount opportunities
- 478 A. van den Nouweland, J. Potters, S. Tijs and J. Zarzuelo
Cores and related solution concepts for multi-choice games
- 479 Drs. C.H. Veld
Warrant pricing: a review of theoretical and empirical research
- 480 E. Nijssen
De Miles and Snow-typologie: Een exploratieve studie in de meubel-
branche
- 481 Harry G. Barkema
Are managers indeed motivated by their bonuses?

- 482 Jacob C. Engwerda, André C.M. Ran, Arie L. Rijkeboer
Necessary and sufficient conditions for the existence of a positive definite solution of the matrix equation $X + A^T X^{-1} A = I$
- 483 Peter M. Kort
A dynamic model of the firm with uncertain earnings and adjustment costs
- 484 Raymond H.J.M. Gradus, Peter M. Kort
Optimal taxation on profit and pollution within a macroeconomic framework
- 485 René van den Brink, Robert P. Gilles
Axiomatizations of the Conjunctive Permission Value for Games with Permission Structures
- 486 A.E. Brouwer & W.H. Haemers
The Gewirtz graph - an exercise in the theory of graph spectra
- 487 Pim Adang, Bertrand Melenberg
Intratemporal uncertainty in the multi-good life cycle consumption model: motivation and application
- 488 J.H.J. Roemen
The long term elasticity of the milk supply with respect to the milk price in the Netherlands in the period 1969-1984
- 489 Herbert Hamers
The Shapley-Entrance Game
- 490 Rezaul Kabir and Theo Vermaelen
Insider trading restrictions and the stock market
- 491 Piet A. Verheyen
The economic explanation of the jump of the co-state variable
- 492 Drs. F.L.J.W. Manders en Dr. J.A.C. de Haan
De organisatorische aspecten bij systeemontwikkeling een beschouwing op besturing en verandering
- 493 Paul C. van Batenburg and J. Kriens
Applications of statistical methods and techniques to auditing and accounting
- 494 Ruud T. Frambach
The diffusion of innovations: the influence of supply-side factors
- 495 J.H.J. Roemen
A decision rule for the (des)investments in the dairy cow stock
- 496 Hans Kremers and Dolf Talman
An SLSPP-algorithm to compute an equilibrium in an economy with linear production technologies

- 497 L.W.G. Strijbosch and R.M.J. Heuts
Investigating several alternatives for estimating the compound lead
time demand in an (s,Q) inventory model
- 498 Bert Bettonvil and Jack P.C. Kleijnen
Identifying the important factors in simulation models with many
factors
- 499 Drs. H.C.A. Roest, Drs. F.L. Tijssen
Beheersing van het kwaliteitsperceptieproces bij diensten door middel
van keurmerken
- 500 B.B. van der Genugten
Density of the F-statistic in the linear model with arbitrarily
normal distributed errors
- 501 Harry Barkema and Sytse Douma
The direction, mode and location of corporate expansions
- 502 Gert Nieuwenhuis
Bridging the gap between a stationary point process and its Palm
distribution
- 503 Chris Veld
Motives for the use of equity-warrants by Dutch companies
- 504 Pieter K. Jagersma
Een etiologie van horizontale internationale ondernemingsexpansie
- 505 B. Kaper
On M-functions and their application to input-output models
- 506 A.B.T.M. van Schaik
Produktiviteit en Arbeidsparticipatie
- 507 Peter Borm, Anne van den Nouweland and Stef Tijs
Cooperation and communication restrictions: a survey
- 508 Willy Spanjers, Robert P. Gilles, Pieter H.M. Ruys
Hierarchical trade and downstream information
- 509 Martijn P. Tummers
The Effect of Systematic Misperception of Income on the Subjective
Poverty Line
- 510 A.G. de Kok
Basics of Inventory Management: Part 1
Renewal theoretic background
- 511 J.P.C. Blanc, F.A. van der Duyn Schouten, B. Pourbabai
Optimizing flow rates in a queueing network with side constraints
- 512 R. Peeters
On Coloring j-Unit Sphere Graphs

- 513 Drs. J. Dagevos, Drs. L. Oerlemans, Dr. F. Boekema
Regional economic policy, economic technological innovation and networks
- 514 Erwin van der Krabben
Het functioneren van stedelijke onroerend-goed-markten in Nederland - een theoretisch kader
- 515 Drs. E. Schaling
European central bank independence and inflation persistence
- 516 Peter M. Kort
Optimal abatement policies within a stochastic dynamic model of the firm
- 517 Pim Adang
Expenditure versus consumption in the multi-good life cycle consumption model
- 518 Pim Adang
Large, infrequent consumption in the multi-good life cycle consumption model
- 519 Raymond Gradus, Sjak Smulders
Pollution and Endogenous Growth
- 520 Raymond Gradus en Hugo Keuzenkamp
Arbeidsongeschiktheid, subjectief ziektegevoel en collectief belang
- 521 A.G. de Kok
Basics of inventory management: Part 2
The (R,S)-model
- 522 A.G. de Kok
Basics of inventory management: Part 3
The (b,Q)-model
- 523 A.G. de Kok
Basics of inventory management: Part 4
The (s,S)-model
- 524 A.G. de Kok
Basics of inventory management: Part 5
The (R,b,Q)-model
- 525 A.G. de Kok
Basics of inventory management: Part 6
The (R,s,S)-model
- 526 Rob de Groof and Martin van Tuijl
Financial integration and fiscal policy in interdependent two-sector economies with real and nominal wage rigidity

- 527 A.G.M. van Eijs, M.J.G. van Eijs, R.M.J. Heuts
Gecoördineerde bestelsystemen
een management-georiënteerde benadering
- 528 M.J.G. van Eijs
Multi-item inventory systems with joint ordering and transportation
decisions
- 529 Stephan G. Vanneste
Maintenance optimization of a production system with buffercapacity
- 530 Michel R.R. van Bremen, Jeroen C.G. Zijlstra
Het stochastische variantie optiewaarderingsmodel
- 531 Willy Spanjers
Arbitrage and Walrasian Equilibrium in Economies with Limited Information

IN 1992 REEDS VERSCHENEN

- 532 F.G. van den Heuvel en M.R.M. Turlings
Privatisering van arbeidsongeschiktheidsregelingen
Refereed by Prof.Dr. H. Verbon
- 533 J.C. Engwerda, L.G. van Willigenburg
LQ-control of sampled continuous-time systems
Refereed by Prof.dr. J.M. Schumacher
- 534 J.C. Engwerda, A.C.M. Ran & A.L. Rijkeboer
Necessary and sufficient conditions for the existence of a positive
definite solution of the matrix equation $X + A^*X^{-1}A = Q$.
Refereed by Prof.dr. J.M. Schumacher
- 535 Jacob C. Engwerda
The indefinite LQ-problem: the finite planning horizon case
Refereed by Prof.dr. J.M. Schumacher
- 536 Gert-Jan Otten, Peter Borm, Ton Storcken, Stef Tijs
Effectivity functions and associated claim game correspondences
Refereed by Prof.dr. P.H.M. Ruys
- 537 Jack P.C. Kleijnen, Gustav A. Alink
Validation of simulation models: mine-hunting case-study
Refereed by Prof.dr.ir. C.A.T. Takkenberg

Bibliotheek K. U. Brabant

17 000 01087530 1