Tilburg University

Sampling for quality inspection and correction

Kleijnen, J.P.C.; Kriens, J.; Lafleur, M.

Publication date:
1986

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Kleijnen, J. P. C., Kriens, J., & Lafleur, M. (1986). *Sampling for quality inspection and correction: AOQL performance criteria*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 246). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

8
CBM
R

1982-246
7626

1986

246

UNIVERSITEIT
BRABANT

POSTBOX 90153
5000 LE TILBURG
THE NETHERLANDS

* C I N O O 3 4 2 *

DEPARTMENT OF ECONOMICS
RESEARCH MEMORANDUM

**SAMPLING FOR QUALITY INSPECTION AND
CORRECTION: AOQL PERFORMANCE CRITERIA**

J.P.C. Kleijnen
J. Kriens
M.C.H.M. Lafleur
J.H.F. Pardoel

510.8

FEW 246

School of Business and Economics
Tilburg University
Post Box 90153
5000 LE Tilburg
Netherlands

Version 1: December 1986

SAMPLING FOR QUALITY INSPECTION AND CORRECTION:
AOQL PERFORMANCE CRITERIA

J.P.C. KLEIJNEN
J. KRIENS
M.C.H.M. LAFLEUR
J.H.F. PARDOEL

School of Business and Economics
Tilburg University
Post Box 90153
5000 LE Tilburg
Netherlands

Keywords: Percentage defectives, quality control, Monte Carlo simulation, auditing.

ABSTRACT

The Average Outgoing Quality Limit (AOQL) denotes a sampling plan leading to inspection of the whole population, if the sample shows a number of defective items k exceeding an acceptance number k_0 . According to the literature, this constant k_0 is chosen such that the expected value of $\tilde{p}$, the fraction of defectives after inspection and possible correction, does not exceed a prespecified constant $\tilde{p}_m$; moreover several other performance criteria are estimated, using an extensive Monte Carlo simulation. The main conclusion is that the AOQL scheme is useful in practice, including applications in auditing, but the chance that the quality constraint is violated, $P\left[\tilde{p} > \tilde{p}_m\right]$, may be sizable.

1. INTRODUCTION: AOQL

AOQL was introduced by Dodge and Romig around 1930; its application to auditing was studied by Kriens and Veenstra (1985). We can summarize this sampling scheme as follows. The goal of AOQL as introduced by Dodge and Romig, is to guarantee a minimum quality of the outgoing populations expressed as a maximum for the average fraction of defectives in the population. Kriens and Veenstra (1985) split up the yearly population into a number of subpopulations. The quality of a yearly population - before sampling and correction - is quantified by p , the percentage of "defective items" in the yearly population. The Number of items (defect plus correct) per Year is (say) NY , for example, a company produces NY cars; in auditing, accounts are sampled and NY is measured in dollars per years; see Kriens and Veenstra (1985, p. 387). Consequently, after inspection and correction the minimum quality corresponds to a maximum value for $\tilde{p}$.

The sampling scheme has the following steps (also see Table 1 later on).

- (i) The expected yearly population is divided into a number of subperiods S , for example, $S = 52$ corresponds to production per week. These subpopulations may have different sizes, in expectation and certainly in realization. We denote the realized size of the subpopulation by N .
- (ii) From each (realized) subpopulation a sample of size n is taken (n depends on several parameters, as we shall see).
- (iii) Per sample (of size n) the number of defective items $\underline{k}$ is determined by inspection; obviously $\underline{k}$ is random (denoted by an underscore). And its integer values k satisfy: $0 \leq k \leq n$.
- (iv) If and only if k exceeds a critical constant k_0 (which varies with n ; see step ii) then the whole subpopulation is

inspected and, by assumption, all defective items in the subpopulation are corrected. If $k \leq k_0$ then the defective items in the sample are corrected. So after sampling the quality of the subpopulation is improved, unless no defectives at all were found ($k = 0$) which probably indicates that the subpopulation had perfect quality already.

After step (iv) the fraction of defectives per subpopulation $\tilde{p}$ should satisfy the minimum-quality requirement $\tilde{p}_m$. So given a correct selection of the sampling plan's parameters n and k_0 (see next paragraph), $\tilde{p}$ should satisfy the condition $E[\tilde{p}] \leq \tilde{p}_m$. Obviously, if the original fraction of defectives (before sampling) was very good already (say, $p = 0$), then $E[\tilde{p}] < p$. If this quality was very bad ($p \gg \tilde{p}_m$), then the sampling plan implies that sampling is (nearly) always followed by inspection and correction of the whole subpopulation so that $(0^*) \tilde{p} \ll \tilde{p}_m$. See Figure 1 where p^* is the "least favorable" value of p .

Obviously k follows the hypergeometric distribution with parameters n , p and N . The critical constant k_0 and n can be computed such that, not only holds $E[\tilde{p}] \leq \tilde{p}_m$, but also the expected costs are minimized. The original tables in Dodge and Romig (1959), however, contain some inaccuracies. Therefore we use the recent tables computed by Kriens and Winters (1987); see also Veenstra and Buysse (1985) and Van Batenburg, Kriens and Veenstra (1987). Table 1 illustrates some typical results.

In practice p , the before-sampling or prior probability, is unknown and often only the left-most columns in tables like Table 1 are used (low p). Even if the prior probability is estimated wrongly, the quality constraint $E[\tilde{p}] \leq \tilde{p}_m$ is met; the expected costs, however, may increase. A conjecture is that, not only does the expected value meet the quality constraint, but also the chance of too bad a yearly quality is negligible.

Figure 1: Expected fraction of defectives after sampling $E(\bar{p})$ versus fraction before sampling p .

Table 1: Sample size n and acceptance number k_o depending on prior probability p , subpopulation size N , and quality limit $\tilde{p}_m$ ($\tilde{p}_m = 1\%$).

Subpopulation size N	Before sampling p							
	0-0.02		...	0.21-0.40		...	0.81-1.00	
N	n	k _o		n	k _o		n	k _o
1-25	11	0		11	0		11	0
26-50	22	0		22	0		22	0
...								
801-1000	35	0		80	1		120	2
...								
1001-2000	36	0		80	1		180	3
20001-50000	85	1		255	4		990	15
50001-100000	85	1		255	4		1520	22

i.e., if $\tilde{p}$ denotes the average yearly outgoing quality then

$$P\left[\tilde{p} > \tilde{p}_m\right] \approx 0 \quad (1.1)$$

This conjecture is investigated in our simulation. Moreover, it hardly takes more computer time to estimate how bad $\tilde{p}$ is if the constraint $\tilde{p} < \tilde{p}_m$ is violated, i.e., we estimate the following conditional expectations:

$$E\left[\tilde{p} - \tilde{p}_m \mid \tilde{p} > \tilde{p}_m\right] \quad (1.2)$$

2. DESIGN OF MONTE CARLO EXPERIMENT

As Table 1 showed, the sample size n and acceptance number k_0 are completely determined by the subpopulation size N , the prior probability p , and the quality limit $\tilde{p}_m$. In turn the subpopulation size N depends on the yearly population size NY and the number of subperiods S . In the simulation we study three values for S , namely 4, 13, and 52 which correspond to quarters, "months", and weeks. Our selection of the yearly population size NY is based in the experience of one of the authors with auditing applications: 10,000; 100,000 and 1,000,000. The expected subpopulation size $E(N)$ equals NY/S . We assume that the actual sizes N follow a uniform distribution with expected value NY/S ; its range is such that the coefficient of variation is always (roughly) 6%, an arbitrarily selected value.

We selected the following six values for the quality limit $\tilde{p}_m$: 0.1%, 0.5%, 1%, 2%, 5%, 10%. Selection of the prior probability p in the simulation must be related to the quality limit $\tilde{p}_m$. There are no tables available for $p > 2 \tilde{p}_m$. This is no problem if only the left columns of the tables are used (see section 1); obviously if p is very high, then the scheme breaks down, that is, sampling is (nearly) always followed by inspection of the whole subpopulation; therefore we restrict our simulation of the "practitioner" to, $p \leq 6 \tilde{p}_m$. Obviously not all subpopulations must have the same p , even if all subpopulations have the same expected value $E(p)$. Therefore we sample p in the simulation. As figure 1 demonstrates, the performance $E(\tilde{p})$ improves as p deviates from the least favourable value p^* . In preliminary simulation experiments we sampled p from a distribution with a high variance, and indeed $E(\tilde{p})$ decreased (not further reported in this paper). Therefore we concentrate our simulation on worst case situations, that is, p has a small

range. We further assume that p is uniformly distributed with a range of only $0.2 \tilde{p}_m$. We do change the expected value $E[p]$; as we explained above, we vary p between 0 and $6 \tilde{p}_m$. So we sample p from the uniform distribution between 0 and $0.2 \tilde{p}_m$, between $0.2 \tilde{p}_m$ and $0.4 \tilde{p}_m$, ... , between $5.8 \tilde{p}_m$ and $6 \tilde{p}_m$.

Summarizing, we simulate 1620 factor combinations using only the left-most columns of the tables ("practitioner's approach") and 540 combinations with the optimal $[n, k_o]$ combinations ("theoretical approach").

There is a technical issue in the simulation: how often should each factor combination be simulated in order to obtain reliable estimates of performance criteria such as $P\left[\tilde{p} > p_m\right]$? By definition, one replicate yields a binomial variable (say) $\underline{x}$ with $q = P(\underline{x} = 0) = P\left[\tilde{p} > \tilde{p}_m\right]$. Using the normal approximation to the binomial distribution, it is straightforward to derive the number of replications needed to estimate q with either a relative precision of 10% or an absolute precision of 0.001; also see Kleijnen (1987, pp. 46-51). This approach shows that at most 16,221 replications are needed (when $\hat{q} = .01$) to satisfy either the relative precision or the absolute precision requirement, with 90% probability. Actually we do not know q . So we substitute the "current" estimate for q (after at least 100 replications) that is, the estimate available after (say) r replications where $r = 101, 102, \dots$. We found that the average number of replications is roughly 1000. We emphasize that the simulation not only estimates the performance criterion $q = P\left[\tilde{p} > \tilde{p}_m\right]$ but several more criteria. The main criterion, however, is q so that we concentrate on q to select the number of replications.

It turns out that it takes 40 hours of computer time to simulate $1620 + 540$ factor combinations, each combination replicated roughly 1000 times. To keep computer time below this (sizable) value, we have to introduce the following technical refinement. The number of defectives $\underline{k}$ has a hypergeometric distribution. The binomial distribution provides a good approximation of $n \ll N$ which is often the case (but not always: if N is small then $n > N$ may occur); see Table 1. The Poisson distribution is a good approximation to the binomial distribution, if p is small. We simulate the Poisson distribution using the subroutine in Naylor et al. (1966, p. 114), so that our program runs 20 times faster on our computer (a VAX 780 running under VMS using the NAG subroutine for the multiplicative congruential random number generator).

3. MONTE CARLO RESULTS

The Monte Carlo experiment yields a mass of data. We analyze these data through regression analysis (using SAS). Preliminary plots look like a gamma function. Therefore we fit such a type of non-linear regression model for $\tilde{p}$; its R^2 adjusted for the number of explanatory variables, is higher than 0.95 and yields Figure 2, which looks like the theoretical Figure 1: $\tilde{p} \leq \tilde{p}_m$.

If the prior probability satisfies $p \leq \tilde{p}_m$ then obviously $q = P\left[\tilde{p} > \tilde{p}_m\right] = 0$. If $p > \tilde{p}_m$ then we again fit a function like the gamma function, with $R^2 = 0.99$ for the theoretical case and 0.74 for the practitioner's approach; see Figure 3. So there is a sizable chance (up to 40% in Figure 3) of violating the quality constraint $\tilde{p} \leq \tilde{p}_m$. However, we repeat that our simulation is a worst case, since p of the subpopulation is sampled from a uniform distribution with a small range.

Figure 2: Average yearly outgoing quality $\bar{p}$ versus fraction of defectives before sampling p (yearly population $NY = 1,000,000$; subperiods $S = 52$).

Figure 3: Estimated probability of quality violation, $P(\tilde{p} > \tilde{p}_m)$ ($NY = 100,000$; $S = 52$).

Figure 4: Estimated size of quality violation, $E(\tilde{p} - \tilde{p}_m \mid \tilde{p} > \tilde{p}_m)$ in theoretical approach ($NY = 1,000,000$).

Figure 5: Estimated fraction of yearly population, actually inspected
 (NY = 1,000,000 ; S = 52).

If $\bar{\tilde{p}} > \tilde{p}_m$ then we wonder how bad the quality violation is: $E[\bar{\tilde{p}} - \tilde{p}_m | \bar{\tilde{p}} > \tilde{p}_m]$. Figure 4 shows that smaller subperiods (higher S) give extra protection.

Next we consider the costs of the AOQL scheme. This scheme implies that all N units (of a subperiod) are inspected if $k > k_0$. Figure 5 shows the fraction of the yearly population actually inspected. That fraction increases drastically if $p > \tilde{p}_m$. Obviously the practitioner's approach is more expensive. We add that the curves are hardly effected by S, the number of subperiods (not displayed).

Our simulation shows that is it important to have a good idea about p, the before sampling fraction of defectives. Therefore we suggest to obtain an estimate of p, using $\hat{p} = k/n$ if $k \leq k_0$ and $\hat{p} = K/N$ if $k > k_0$ where K denotes the number of defectives in the subpopulation (of size N). As time t goes on, the estimators $\hat{p}_t$ can be combined; for example, we may weigh the $\hat{p}_t$ with the sample sizes n_t or the subpopulation sizes N_t (if $k \leq k_0$ or $k > k_0$ respectively). If $\hat{p}_t$ shows serial correlation or non-stationary behavior, then time series techniques may be applied.

4. CONCLUSIONS

The AOQL sampling plan is indeed used in practice; see Kriens and Veenstra (1985). Then it is assumed that if the expected yearly fraction of defectives after inspection and correction $E(\tilde{p})$ meets the quality constraint $\tilde{p}_m$ then the probability of exceeding the constraint, $P[\bar{\tilde{p}} > \tilde{p}_m]$ is negligible. Figure 3 (based on our simulation data analyzed by regression) shows that this probability is sizable, if the "before inspection" or prior probability p is higher than the limit $\tilde{p}_m$ but

not extremely high (if $p \gg \tilde{p}_m$ then most times sampling is followed by inspection of the whole subpopulation). If in practice p varies much over subperiods, then $P\left[\tilde{p} > \tilde{p}_m\right]$ decreases (we simulated worst case situations: small range of p). One can get an estimate of p from the sampling procedure: if $k \leq k_0$ then $\hat{p} = k/n$; else $\hat{p} = K/N$. Increasing the number of periods S decreases the size of the expected quality violation; see Figure 4. Underestimating p is not wise: it does not give extra quality protection (Figure 3) and yet more inspection work is done (Figure 5). So in practice one should build up knowledge about p .

REFERENCES

- Dodge, H.F. and H.G. Romig, *Sampling inspection tables*. John Wiley, New York, second edition, 1959.
- Kleijnen, J.P.C., *Statistical tools for simulation practitioners*. Marcel Dekker, Inc., New York, 1987.
- Kriens, J. and R.H. Veenstra, Statistical sampling in internal control by using the AOQL-system. *The Statistician*, 34, 1985, pp. 383-390.
- Kriens, J. and A.A.G.M. Winters (1987), The A.O.Q.L. tables by Dodge and Romig: some corrections. (to appear.)
- Naylor, T.H., J.L. Balintfy, D.S. Burdick and K. Chu, *Computer simulation techniques*. John Wiley & Sons, Inc., New York, 1966.
- Van Batenburg, P.C., J. Kriens and R.H. Veenstra (1987), Average Outgoing Quality Limit - a revised and improved version. (to appear in Dutch.)
- Veenstra, R.H. and J.C. Buysse (1985), Optimizing sampling applications in the administration, *De Accountant*, 10, no. 198, pp. 561-563. (in Dutch.)

IN 1986 REEDS VERSCHENEN

- 202 J.H.F. Schilderink
Interregional Structure of the European Community. Part III
- 203 Antoon van den Elzen and Dolf Talman
A new strategy-adjustment process for computing a Nash equilibrium in a noncooperative more-person game
- 204 Jan Vingerhoets
Fabrication of copper and copper semis in developing countries. A review of evidence and opportunities
- 205 R. Heuts, J. van Lieshout, K. Baken
An inventory model: what is the influence of the shape of the lead time demand distribution?
- 206 A. van Soest, P. Kooreman
A Microeconomic Analysis of Vacation Behavior
- 207 F. Boekema, A. Nagelkerke
Labour Relations, Networks, Job-creation and Regional Development. A view to the consequences of technological change
- 208 R. Alessie, A. Kapteyn
Habit Formation and Interdependent Preferences in the Almost Ideal Demand System
- 209 T. Wansbeek, A. Kapteyn
Estimation of the error components model with incomplete panels
- 210 A.L. Hempenius
The relation between dividends and profits
- 211 J. Kriens, J.Th. van Lieshout
A generalisation and some properties of Markowitz' portfolio selection method
- 212 Jack P.C. Kleijnen and Charles R. Standridge
Experimental design and regression analysis in simulation: an FMS case study
- 213 T.M. Doup, A.H. van den Elzen and A.J.J. Talman
Simplicial algorithms for solving the non-linear complementarity problem on the simplotope
- 214 A.J.W. van de Gevel
The theory of wage differentials: a correction
- 215 J.P.C. Kleijnen, W. van Groenendaal
Regression analysis of factorial designs with sequential replication
- 216 T.E. Nijman and F.C. Palm
Consistent estimation of rational expectations models

- 217 P.M. Kort
The firm's investment policy under a concave adjustment cost function
- 218 J.P.C. Kleijnen
Decision Support Systems (DSS), en de kleren van de keizer ...
- 219 T.M. Doup and A.J.J. Talman
A continuous deformation algorithm on the product space of unit simplices
- 220 T.M. Doup and A.J.J. Talman
The 2-ray algorithm for solving equilibrium problems on the unit simplex
- 221 Th. van de Klundert, P. Peters
Price Inertia in a Macroeconomic Model of Monopolistic Competition
- 222 Christian Mulder
Testing Korteweg's rational expectations model for a small open economy
- 223 A.C. Meijdam, J.E.J. Plasmans
Maximum Likelihood Estimation of Econometric Models with Rational Expectations of Current Endogenous Variables
- 224 Arie Kapteyn, Peter Kooreman, Arthur van Soest
Non-convex budget sets, institutional constraints and imposition of concavity in a flexible household labor supply model
- 225 R.J. de Groof
Internationale coördinatie van economische politiek in een twee-regio-twee-sectoren model
- 226 Arthur van Soest, Peter Kooreman
Comment on 'Microeconomic Demand Systems with Binding Non-Negativity Constraints: The Dual Approach'
- 227 A.J.J. Talman and Y. Yamamoto
A globally convergent simplicial algorithm for stationary point problems on polytopes
- 228 Jack P.C. Kleijnen, Peter C.A. Karremans, Wim K. Oortwijn, Willem J.H. van Groenendaal
Jackknifing estimated weighted least squares
- 229 A.H. van den Elzen and G. van der Laan
A price adjustment for an economy with a block-diagonal pattern
- 230 M.H.C. Paardekooper
Jacobi-type algorithms for eigenvalues on vector- and parallel computer
- 231 J.P.C. Kleijnen
Analyzing simulation experiments with common random numbers

- 232 A.B.T.M. van Schaik, R.J. Mulder
On Superimposed Recurrent Cycles
- 233 M.H.C. Paardekooper
Sameh's parallel eigenvalue algorithm revisited
- 234 Pieter H.M. Ruys and Ton J.A. Storcken
Preferences revealed by the choice of friends
- 235 C.J.J. Huys en E.N. Kertzman
Effectieve belastingtarieven en kapitaalkosten
- 236 A.M.H. Gerards
An extension of König's theorem to graphs with no odd- K_4
- 237 A.M.H. Gerards and A. Schrijver
Signed Graphs - Regular Matroids - Grafts
- 238 Rob J.M. Alessie and Arie Kapteyn
Consumption, Savings and Demography
- 239 A.J. van Reeken
Begrippen rondom "kwaliteit"
- 240 Th.E. Nijman and F.C. Palmer
Efficiency gains due to using missing data. Procedures in regression models
- 241 Dr. S.C.W. Eijffinger
The determinants of the currencies within the European Monetary System

IN 1987 REEDS VERSCHENEN

- 242 Gerard van den Berg
Nonstationarity in job search theory
- 243 Annie Cuyt, Brigitte Verdonk
Block-tridiagonal linear systems and branched continued fractions
- 244 J.C. de Vos, W. Vervaat
Local Times of Bernoulli Walk
- 245 Arie Kapteyn, Peter Kooreman, Rob Willemsse
Some methodological issues in the implementation
of subjective poverty definitions

Bibliotheek K. U. Brabant

17 000 01059975 2