Tilburg University

Eigenvalues and the diameter of graphs

van Dam, E.R.; Haemers, W.H.

Publication date:
1993

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
van Dam, E. R., & Haemers, W. H. (1993). *Eigenvalues and the diameter of graphs*. (CentER Discussion Paper; Vol. 1993-43). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

entER
for
Economic Research

Discussion paper

ECO
CBM
RR
8414
991993
43

Faint handwritten text, possibly "Discussion paper" and "Sijpe"

Center
for
Economic Research

No. 9343

**Eigenvalues and the
Diameter of Graphs**

by Edwin R. van Dam
and Willem H. Haemers

June 1993

ISSN 0924-7815

K.U.B.
BIBLIOTHEEK
TILBURG

Eigenvalues and the Diameter of Graphs

EDWIN R. VAN DAM WILLEM H. HAEMERS

Abstract. Using eigenvalue interlacing and Chebyshev polynomials we find upper bounds for the diameter of regular and bipartite biregular graphs in terms of their eigenvalues. This improves results of Chung and Delorme and Solé. The same method gives upper bounds for the number of vertices at a given minimum distance from a given vertex set. These results have some applications to the covering radius of error-correcting codes.

1. Introduction

It is an old result that the diameter of a connected graph is smaller than the number of different eigenvalues of its adjacency matrix (cf. [3]). More recently, Chung [2] and Delorme and Solé [4] derived bounds for the diameter of regular graphs in terms of the actual value of (some of) the eigenvalues. In this paper we shall derive a tool which we can use to derive upper bounds for the diameter of regular (multi)graphs. The tool leaves us some freedom to choose different polynomials. The bounds then depend on the values of the polynomial evaluated at the eigenvalues. By suitable choices of the polynomial we find all bounds mentioned above. But we can do better by using Chebyshev polynomials. Our method also gives bounds for the number of vertices at a given minimum distance from a given vertex set, and we also improve the bounds of Delorme and Solé for the diameter of bipartite biregular graphs. We should mention that Mohar [8] found diameter bounds in terms of the Laplacian eigenvalues of the graph. Though our method also applies to the Laplacian matrix of a (possibly nonregular) graph (see Section 2.5), Mohar's results don't seem to fit in our framework. However, our bound seems to be better (see Section 4.2 for examples). Finally, we apply our bounds to Ramanujan graphs, and to the coset graphs of linear codes to obtain an upper bound for the covering radius of a linear code.

Throughout the paper, G will be an undirected graph with n vertices, adjacency matrix A and diameter $d(G)$. We allow G to be a multigraph, that is, G may have multiple edges and loops (a loop counts for one edge in the degree). The eigenvalues (of the adjacency matrix) of G are denoted by $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$. Similarly we denote the eigenvalues of an $m \times m$ matrix M by $\lambda_1(M) \geq \lambda_2(M) \geq \dots \geq \lambda_m(M)$, if the spectrum of M is real. (All the matrices we use have real spectrum.) The polynomials we use have real coefficients.

2. Regular Graphs

2.1. The Tool

THEOREM 2.1. *Let G be connected and regular of degree k . Let m be a nonnegative integer and let X and Y be sets of sizes x and y , respectively, such that the distance between any vertex of X and any vertex of Y is at least $m + 1$. If p is a polynomial of degree m such that $p(k) = 1$, then*

$$\frac{xy}{(n-x)(n-y)} \leq \max_{i \neq 1} p^2(\lambda_i) .$$

Proof. If we take $\{1, 2, \dots, n\}$ for our vertex set, and (without loss of generality) $X = \{1, \dots, x\}$, $Y = \{n - y + 1, \dots, n\}$, then $p(A)_{ij} = 0$ for all $i = 1, \dots, x$, and $j = n - y + 1, \dots, n$. Now consider the matrix

$$M = \begin{bmatrix} O & : & p(A) \\ \dots & & \dots \\ p(A) & : & O \end{bmatrix} .$$

Note that M is symmetric, has row and column sums equal to 1, and its spectrum is $\{\pm p(\lambda_i) \mid i = 1, 2, \dots, n\}$. Let M be partitioned symmetrically in the following way.

$$M = \begin{bmatrix} O & : & O & : & : & O \\ \dots & & \dots & & \dots & \dots \\ O & : & O & : & : & \dots \\ \dots & & \dots & & \dots & \dots \\ : & & : & & O & : & O \\ \dots & & \dots & & \dots & & \dots \\ O & : & : & & O & : & O \end{bmatrix} \begin{matrix} \} x \\ \\ \} n-x \\ \\ \} n-y \\ \\ \} y \end{matrix} .$$

Let B be the matrix of average row sums in the blocks of this partition, then

$$B = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 1 - \frac{y}{n-x} & \frac{y}{n-x} \\ \frac{x}{n-y} & 1 - \frac{x}{n-y} & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} ,$$

and the eigenvalues of B are $\lambda_1(B) = -\lambda_4(B) = 1$, $\lambda_2(B) = -\lambda_3(B) = \sqrt{\frac{xy}{(n-x)(n-y)}}$. Since the eigenvalues of B interlace those of M (cf. [5]), we have that

$$\lambda_2(B) \leq \lambda_2(M) \leq \max_{i \neq 1} |p(\lambda_i)|, \text{ so } \frac{xy}{(n-x)(n-y)} \leq \max_{i \neq 1} p^2(\lambda_i) .$$

□

2.2. The Diameter

The first application of Theorem 2.1 is to prove that the diameter of G is smaller than the number of different eigenvalues of G . Let G have r different eigenvalues $k = \mu_1 > \mu_2 > \dots > \mu_r$. Now let p be the polynomial of degree $r - 1$, given by

$$p(z) = \prod_{i \neq 1} \frac{z - \mu_i}{k - \mu_i} ,$$

then $p(k) = 1$ and $p(\mu_i) = 0$ for $i \neq 1$. So if X is a set of x vertices and Y is a set of y vertices, such that the distance between X and Y is at least r , then

$$\frac{xy}{(n-x)(n-y)} \leq 0,$$

so $x = 0$ or $y = 0$. This implies that any two vertices have distance smaller than r , so $d(G) < r$.

If we take $m = d(G) - 1$, then there are (at least) two vertices which have distance $m + 1$.

Let X and Y be singletons consisting of two such vertices, and let p be given by $p(z) = \left(\frac{z}{k}\right)^m$.

Now apply Theorem 2.1, then we find that

$$\frac{1}{n-1} \leq \left[\frac{\lambda}{k}\right]^m, \text{ so } d(G) \leq \frac{\log(n-1)}{\log\left(\frac{k}{\lambda}\right)} + 1, \text{ where } \lambda = \max_{i \neq 1} |\lambda_i|.$$

This bound (which is only applicable to nonbipartite graphs) was found by Chung [2]. Similarly

we find the bound of Delorme and Solé [4] by taking $p(z) = \frac{z+t}{k+t} \left(\frac{z}{k}\right)^{m-1}$ for arbitrary (real) t .

But we can do better. Theorem 2.1 allows us to use any polynomial p of degree m such that $p(k) = 1$. To get the sharpest bound possible in this way, however, we must choose p such that the right hand side of the inequality in Theorem 2.1 is minimized. An obvious approach is to minimize over all polynomials p of the form

$$p(z) = \left(\frac{z-\alpha}{k-\alpha}\right)^m, \alpha \in \mathbb{R}.$$

This gives the following bound.

THEOREM 2.2. *Let G be connected and regular of degree k (not complete), then*

$$d(G) \leq \frac{\log(n-1)}{\log\left[\frac{2k-\lambda_2-\lambda_n}{\lambda_2-\lambda_n}\right]} + 1.$$

Proof. Like in the previous example, take $m = d(G) - 1$ and let X and Y be singletons at distance $d(G)$. For p we take

$$p(z) = \left[\frac{z - \frac{1}{2}(\lambda_2 + \lambda_n)}{k - \frac{1}{2}(\lambda_2 + \lambda_n)}\right]^m, \text{ then we find that } \frac{1}{n-1} \leq \left[\frac{\frac{1}{2}(\lambda_2 - \lambda_n)}{k - \frac{1}{2}(\lambda_2 + \lambda_n)}\right]^m.$$

Since G is not complete, $\lambda_2 \neq \lambda_n$, and the bound follows. $\square$

For $m = 1$ this is the best possible, but in general we can still do better by looking at a "relaxation" of our problem. This is the problem of minimizing $\max \{p^2(x) \mid \lambda_n \leq x \leq \lambda_2\}$ over all polynomials p of degree m such that $p(k) = 1$. The solution of this problem can be given in terms of Chebyshev polynomials (cf. [9, Th. 2.1, Ex. 2.5.12]). We have to take

$$\check{C}(z) = \frac{T_m \left[\frac{2z - \lambda_2 - \lambda_n}{\lambda_2 - \lambda_n} \right]}{T_m \left[\frac{2k - \lambda_2 - \lambda_n}{\lambda_2 - \lambda_n} \right]}, \text{ where } T_m(x) = \frac{1}{2}(x + \sqrt{x^2 - 1})^m + \frac{1}{2}(x - \sqrt{x^2 - 1})^m$$

is the m -th Chebyshev polynomial. We shall use the following properties of this polynomial:

- (i) $\max_{x \in [-1, 1]} |T_m(x)| = 1$,
- (ii) $T_m \left[\frac{2k - \beta - \alpha}{\beta - \alpha} \right] = \frac{1}{2} \left[\frac{\sqrt{k - \alpha} + \sqrt{k - \beta}}{\sqrt{k - \alpha} - \sqrt{k - \beta}} \right]^m + \frac{1}{2} \left[\frac{\sqrt{k - \alpha} - \sqrt{k - \beta}}{\sqrt{k - \alpha} + \sqrt{k - \beta}} \right]^m$,
- (iii) $T_m(x)$ is an even, respectively odd polynomial, if m is even, respectively odd.

THEOREM 2.3. *Let G be connected and regular of degree k (not complete), then*

$$d(G) < \frac{\log 2(n-1)}{\log \left[\frac{\sqrt{k - \lambda_n} + \sqrt{k - \lambda_2}}{\sqrt{k - \lambda_n} - \sqrt{k - \lambda_2}} \right]} + 1.$$

Proof. Take the suggested polynomial $\check{C}(z)$, then we find that

$$n - 1 \geq T_m \left[\frac{2k - \lambda_2 - \lambda_n}{\lambda_2 - \lambda_n} \right] > \frac{1}{2} \left[\frac{\sqrt{k - \lambda_n} + \sqrt{k - \lambda_2}}{\sqrt{k - \lambda_n} - \sqrt{k - \lambda_2}} \right]^m,$$

from which the bound follows. (Again, $m = d(G) - 1$ and $x = y = 1$.) □

2.3. The Number of Distant Vertices

Now let X be a set of vertices of size x in the graph G . We shall derive a bound on the number y of vertices, which have distance at least d from X .

THEOREM 2.4. *Let G be connected and regular of degree k (not complete). Let X be an arbitrary set of vertices of size x , and y the number of vertices at distance at least d from X , where d is a positive integer, then*

$$y < \frac{n}{1 + \frac{x}{4(n-x)} \left[\frac{\sqrt{k - \lambda_n} + \sqrt{k - \lambda_2}}{\sqrt{k - \lambda_n} - \sqrt{k - \lambda_2}} \right]^{2(d-1)}}.$$

Proof. Again, we take the polynomial $\check{C}(z)$, with $m = d - 1$. Furthermore we let Y be the set of vertices at distance at least d from X . Now Theorem 2.1 implies that

$$\frac{(n-x)(n-y)}{xy} \geq T_m \left(\frac{2k - \lambda_2 - \lambda_n}{\lambda_2 - \lambda_n} \right)^2 > \frac{1}{4} \left(\frac{\sqrt{k - \lambda_n} + \sqrt{k - \lambda_2}}{\sqrt{k - \lambda_n} - \sqrt{k - \lambda_2}} \right)^{2m},$$

and the bound follows. $\square$

2.4. An Extreme Case

Let G be connected and regular with r different eigenvalues $k = \mu_1 > \mu_2 > \dots > \mu_r$. We have already proven that $d(G) \leq r - 1$. In case of equality we are able to determine the polynomial which minimizes our upper bound. Then we take p such that $p(\mu_j) = (-1)^j c$, $j = 2, \dots, r$, where c is chosen such that $p(k) = 1$. That this p is really the best polynomial of degree $r - 2$ we can choose, can be seen in the following way. Suppose there is a polynomial g of degree $r - 2$ such that $g(k) = 1$, and $|g(\mu_i)| < c$, for all $i = 2, 3, \dots, r$. Let h be given by $h = p - g$, then $h(k) = 0$, and $h(\mu_i) = c - g(\mu_i) > 0$ for even i , and $h(\mu_i) = -c - g(\mu_i) < 0$ for odd $i \neq 1$. This implies that h has at least $r - 1$ roots, which is a contradiction.

THEOREM 2.5. *Let G be a connected regular graph of degree k , with r different eigenvalues $k = \mu_1 > \mu_2 > \dots > \mu_r$. If $d(G) = r - 1$, then*

$$n \geq 1 + \sum_{j=2}^r \prod_{i \neq j, 1} \frac{k - \mu_i}{|\mu_j - \mu_i|}.$$

Proof. As suggested, we have to take

$$p(z) = \sum_{j=2}^r (-1)^j c \mathcal{P}_j(z), \text{ where } \mathcal{P}_j(z) = \prod_{i \neq j, 1} \frac{z - \mu_i}{\mu_j - \mu_i}, j = 2, \dots, r,$$

and where c is such that $p(k) = 1$. Since the degree of p is $r - 2$, it follows from Theorem 2.1 that $\frac{1}{n-1} \leq c$. Since $p(k) = 1$ we have that

$$\frac{1}{c} = \sum_{j=2}^r (-1)^j \prod_{i \neq j, 1} \frac{k - \mu_i}{\mu_j - \mu_i} = \sum_{j=2}^r \prod_{i \neq j, 1} \frac{k - \mu_i}{|\mu_j - \mu_i|},$$

from which the result follows. $\square$

2.5. Laplacian Eigenvalues

If Γ is not a regular graph, it can be transformed into a regular graph G by adding a suitable number of loops to every vertex. If k is the maximum degree in Γ , we add $k - \text{degree}(v)$ loops to every vertex v , so that G is regular of degree k . Moreover, there is a relation between the eigenvalues of the Laplacian matrix Q of Γ and the eigenvalues of the adjacency matrix A of G . The Laplacian matrix Q of Γ is defined by $Q = \text{diag}(\text{degree}(v)) - A(\Gamma)$, where $A(\Gamma)$ is the adjacency matrix of Γ . Now it follows that $Q = kI - A$, so if $\theta_1 \leq \theta_2 \leq \dots \leq \theta_n$ are the Laplacian eigenvalues of Γ , then $\theta_i = k - \lambda_i$, $i = 1, 2, \dots, n$. Since the distance between two vertices is the same in Γ and G , we can get bounds in terms of the Laplacian eigenvalues of Γ . For example, Theorem 2.3 now says that

$$d(\Gamma) < \frac{\log 2(n-1)}{\log \left[\frac{\sqrt{\theta_n} + \sqrt{\theta_2}}{\sqrt{\theta_n} - \sqrt{\theta_2}} \right]} + 1.$$

3. Bipartite Biregular Graphs

A bipartite graph G is a graph that allows a partition of the vertex set into two parts V_1 and V_2 such that all edges are between V_1 and V_2 . (Note that this implies that there are no loops.) It is well known that if λ is an eigenvalue of G , then $-\lambda$ is also an eigenvalue of G with the same multiplicity. A bipartite graph G is biregular if every vertex in V_1 is adjacent to k_1 vertices and every vertex in V_2 is adjacent to k_2 vertices, for some k_1 and k_2 . Note that

$$\lambda_1 = \sqrt{k_1 k_2}, \text{ and } \lambda_i = -\lambda_{n-i+1} \text{ for all } i = 1, 2, \dots, n.$$

Delorme and Solé [4] found a bound for the diameter of bipartite biregular graphs. Just like them we shall distinguish between the distance of vertices in the same part, and the distance of vertices in different parts. However, the maximum distance between two vertices in the same part, and the maximum distance between two vertices in different parts differ only one.

3.1. The Tool

From the next two theorems, which will be proven in a similar way as Theorem 2.1 (actually, one will be deduced from it), it is easy to derive the bounds of Delorme and Solé. However, we can improve them.

THEOREM 3.1. *Let G be connected and bipartite biregular, with n_1 vertices (V_1) of degree k_1 , and n_2 vertices (V_2) of degree k_2 . Let m be odd, and X_1 a subset of V_1 of size x_1 , and Y_2 a subset of V_2 of size y_2 , such that the distance between any vertex in X_1 and any vertex in Y_2 is at least $m + 2$. Let p be an odd polynomial of degree m , such that $p(\lambda_i) = 1$, then*

$$\frac{x_1 y_2}{(n_1 - x_1)(n_2 - y_2)} \leq \max_{i \neq 1, n} p^2(\lambda_i).$$

Proof. Let $p(A)$ be partitioned symmetrically as follows.

$$p(A) = \begin{bmatrix} O & O & & & O \\ \dots & \dots & & & \dots \\ O & O & & & O \\ \dots & \dots & & & \dots \\ & & & O & O \\ \dots & \dots & & \dots & \dots \\ O & & & O & O \end{bmatrix} \begin{matrix} x_1 \\ n_1 - x_1 \\ n_2 - y_2 \\ y_2 \end{matrix}$$

Let B be the matrix of average row sums in the blocks of this partition, then

$$B = \begin{bmatrix} 0 & 0 & \kappa & 0 \\ 0 & 0 & \kappa - \frac{y_2}{n_1 - x_1} \frac{1}{\kappa} & \frac{y_2}{n_1 - x_1} \frac{1}{\kappa} \\ \frac{x_1}{n_2 - y_2} \kappa & \frac{1}{\kappa} - \frac{x_1}{n_2 - y_2} \kappa & 0 & 0 \\ 0 & \frac{1}{\kappa} & 0 & 0 \end{bmatrix}, \text{ where } \kappa = \sqrt{\frac{k_1}{k_2}}.$$

The eigenvalues of B are $\pm 1, \pm \sqrt{\frac{x_1 y_2}{(n_1 - x_1)(n_2 - y_2)}}$, and since they interlace those of $p(A)$, we have

$$\lambda_2(B) \leq \lambda_2(p(A)) \leq \max_{i \neq 1, n} |p(\lambda_i)|, \text{ and}$$

$$\lambda_1(B) \leq \lambda_1(p(A)) = \max_i |p(\lambda_i)| = \max\{1, \max_{i \neq 1, n} |p(\lambda_i)|\},$$

and the result follows. $\square$

THEOREM 3.2. *Let G be connected and bipartite biregular, with n_1 vertices (V_1) of degree k_1 , and n_2 vertices (V_2) of degree k_2 . Let m be even, $j = 1$ or 2 , and X_j a subset of V_j of size x_j , and Y_j a subset of V_j of size y_j , such that the distance between any vertex in X_j and any vertex in Y_j is at least $m + 2$. Let p be an even polynomial of degree m , such that $p(\lambda_1) = 1$, then*

$$\frac{x_j y_j}{(n_j - x_j)(n_j - y_j)} \leq \max_{i \neq 1, n} p^2(\lambda_i).$$

Proof. Since p is an even polynomial of degree m , there is a polynomial q of degree $m/2$ such that $p(z) = q(z^2)$. Now we consider the matrix A^2 , which has row and column sums equal to $k_1 k_2$, and eigenvalues λ_i^2 , $i = 1, 2, \dots, n$. Furthermore, we can write A^2 as

$$A^2 = \begin{bmatrix} M_1 & \vdots & O \\ \vdots & \ddots & \vdots \\ O & \vdots & M_2 \end{bmatrix} \begin{matrix} n_1 \\ \\ n_2 \end{matrix}.$$

Note that the spectrum of A^2 is the union of the spectra of M_1 and M_2 , and both M_1 and M_2 have one eigenvalue $k_1 k_2$. By applying Theorem 2.1 to the graph on V_j with adjacency matrix M_j , and the polynomial q , we prove the theorem, since the distance in this new graph (which is one of the so-called halved (multi)graphs) is half of the original distance (in G). $\square$

3.2. The Diameter

In the following, we denote by $d_{ij}(G)$ the maximum distance between any vertex in V_i and any vertex in V_j , for $i, j = 1$ or 2 .

THEOREM 3.3. *Let G be connected and bipartite biregular, with n_1 vertices (V_1) of degree k_1 , and n_2 vertices (V_2) of degree k_2 , then*

$$d_{12}(G) < \frac{\frac{1}{2} \log 4(n_1 - 1)(n_2 - 1)}{\log \left(\frac{\lambda_1}{\lambda_2} + \sqrt{\frac{\lambda_1^2}{\lambda_2^2} - 1} \right)} + 2 .$$

Proof. Let $m = d_{12}(G) - 2$, so m is odd, and let X_1, Y_2 be singletons consisting of vertices, which have maximum distance. Let p be given by

$$p(z) = \frac{T_m\left(\frac{z}{\lambda_1}\right)}{T_m\left(\frac{\lambda_1}{\lambda_2}\right)} ,$$

so p is also odd. Now we apply Theorem 3.1 and find that

$$(n_1 - 1)(n_2 - 1) \geq T_m^2\left(\frac{\lambda_1}{\lambda_2}\right) > \frac{1}{4} \left(\frac{\lambda_1}{\lambda_2} + \sqrt{\frac{\lambda_1^2}{\lambda_2^2} - 1} \right)^{2m} ,$$

from which the bound follows. □

In a similar way the next theorem can be derived from Theorem 3.2.

THEOREM 3.4. *Let G be connected and bipartite biregular, with n_1 vertices (V_1) of degree k_1 , and n_2 vertices (V_2) of degree k_2 . Let $j = 1$ or 2 , then*

$$d_{jj}(G) < \frac{\log 2(n_j - 1)}{\log \left(\frac{\lambda_1}{\lambda_2} + \sqrt{\frac{\lambda_1^2}{\lambda_2^2} - 1} \right)} + 2 .$$

□

If G is bipartite and regular (so $k = k_1 = k_2$), we can combine Theorems 3.3 and 3.4 to obtain the following.

COROLLARY 3.5. *Let G be connected, bipartite and regular of degree k , then*

$$d(G) < \frac{\log(n - 2)}{\log \left(\frac{k}{\lambda_2} + \sqrt{\frac{k^2}{\lambda_2^2} - 1} \right)} + 2 .$$

□

The proof of Theorem 3.2 shows a way for a subtle improvement of Theorem 3.4 if the graph G has no eigenvalue equal to zero.

THEOREM 3.6. *Let G be connected and bipartite biregular, with n_1 vertices (V_1) of degree k_1 , and n_2 vertices (V_2) of degree k_2 . Let $j = 1$ or 2 , then*

$$d_{jj}(G) < \frac{2 \log 2(n_j - 1)}{\log \left[\frac{\sqrt{\lambda_1^2 - \lambda_i^2} + \sqrt{\lambda_1^2 - \lambda_2^2}}{\sqrt{\lambda_1^2 - \lambda_i^2} - \sqrt{\lambda_1^2 - \lambda_2^2}} \right]} + 2, \text{ where } t = \left\lceil \frac{n}{2} \right\rceil.$$

Proof. Our polynomial p is now given by

$$p(z) = \frac{T_{\frac{m}{2}} \left[\frac{2z^2 - \lambda_2^2 - \lambda_i^2}{\lambda_2^2 - \lambda_i^2} \right]}{T_{\frac{m}{2}} \left[\frac{2\lambda_1^2 - \lambda_2^2 - \lambda_i^2}{\lambda_2^2 - \lambda_i^2} \right]},$$

where $m = d_{jj}(G) - 2$. Now Theorem 3.2 implies that

$$n_j - 1 \geq T_{\frac{m}{2}} \left[\frac{2\lambda_1^2 - \lambda_2^2 - \lambda_i^2}{\lambda_2^2 - \lambda_i^2} \right] > \frac{1}{2} \left[\frac{\sqrt{\lambda_1^2 - \lambda_i^2} + \sqrt{\lambda_1^2 - \lambda_2^2}}{\sqrt{\lambda_1^2 - \lambda_i^2} - \sqrt{\lambda_1^2 - \lambda_2^2}} \right]^{\frac{m}{2}},$$

from which the bound follows. □

4. Ramanujan Graphs and Other Examples

4.1 Ramanujan Graphs

Let G be regular of degree k , and let λ be the maximum of the absolute values of the eigenvalues of G , unequal to $\pm k$. The graphs G for which

$$\lambda \leq 2\sqrt{k-1}$$

are called Ramanujan graphs. If we apply our bounds for the diameter to these graphs, we get

$$d(G) < \frac{2 \log 2(n-1)}{\log(k-1)} + 1$$

for nonbipartite Ramanujan graphs and

$$d(G) < \frac{2 \log(n-2)}{\log(k-1)} + 2$$

for bipartite Ramanujan graphs, which is (a minor improvement of) the bound of Lubotzky, Phillips and Sarnak [6]. This means that these graphs have a small diameter, since the upper bound is approximately twice the lower bound

$$d(G) \geq \frac{\log((n-1)^{\frac{k-2}{k}} + 1)}{\log(k-1)},$$

which holds for every regular graph of degree k . (We get this lower bound by observing that the number of points at a given distance d from a given point is at most $k(k-1)^{d-1}$.)

4.2 Other Examples

• In the following table we give some examples of spectra of graphs and compute the various upper bounds for the diameter, to get some idea of how good they can be.

# points and spectrum	# different eigenvalues - 1	Th. 2.3	Th. 2.2	Delorme and Solé [4]	Chung [2]	Mohar [8]	Th. 2.5
$n = 10$, eigenvalues $3, 1, -2$. (Petersen graph, $d(G) = 2$.)	2	2	3	3	6	4	-
$n = 28$, eigenvalues $3, 2, \sqrt{2} - 1, -1,$ $-1 - \sqrt{2}$. (Coxeter graph, $d(G) = 4$.)	4	5	9	8	16	8	-
$n = 125$, eigenvalues $64, 4, -1, -16$. (K_5^3 , $d(G) = 2$.)	3	3	3	3	4	6	2
$n = 5^{10}$, eigenvalues $4^{10}, 4^8, \dots, 1, -4,$ $-4^3, \dots, -4^9$. (K_5^{10} , $d(G) = 2$.)	10	6	8	7	12	18	9

Note that K_5^n is the product of n copies of the complete graph on five points, so two vertices are adjacent if they differ in all coordinates. In the two examples of such graphs, Theorem 2.5 gives a contradiction if we assume that the diameter is one less than the number of different eigenvalues.

5. The Covering Radius of Error-Correcting Codes

In this section we give some applications of the diameter bounds to error-correcting codes. The reader, who is not familiar with the basic concepts of coding theory is referred to MacWilliams and Sloane [7].

5.1. The Hamming Graph

Let C be a code of length N over an alphabet Q of q elements. We can apply Theorem 2.1 to the Hamming graph $H(N, q)$ to derive an upper bound for the covering radius of C in terms of N , q and the size of C , since the code C is nothing but a subset of vertices in the Hamming graph, and the covering radius of C is the maximum distance of any of the vertices to C . However, we only derive a useless bound. In the next section we give another approach for linear codes over $GF(q)$.

5.2. The Coset Graph of a Linear Code

Given a linear code we can make a graph that has diameter equal to the covering radius of the code. The eigenvalues of the graph can be expressed in terms of the weights of the dual code. In this way we can derive bounds for the covering radius of a code, using the bounds for the diameter of the graph (cf. [4]).

Let C be a linear code over $GF(q)$ of length N and dimension K . The coset graph of C is a graph G with vertex set $GF(q)^N/C$ (the cosets of C) of size $n = q^{N-K}$. The number of edges between two cosets $x + C$ and $y + C$ is the number of words of weight one in $x - y + C$ (which is equal to the number of words of weight one in $y - x + C$). In this way we get a regular graph of degree $k = N(q - 1)$, possibly with loops (which occurs whenever there is a codeword of weight one) or multiple edges (which occurs whenever there is a coset containing more than one word of weight one, which is the case if and only if there is a codeword of weight two or there are at least two codewords of weight one (in the latter case we have multiple loops)). It is easily seen that the covering radius of C is equal to the diameter of G . Furthermore it is known [1] that the spectrum of G is (the multiset) $\{N(q - 1) - qw(x) \mid x \in C^\perp\}$.

The first observation we can make now is that the covering radius is not bigger than the external distance, since the latter equals the number of nonzero weights in the dual code. (Note that in case of equality, we can apply Theorem 2.5.) Application of our bounds for the diameter of G give the following results.

THEOREM 5.1. *Let C be a linear code over $GF(q)$ of length N and dimension K , and covering radius ρ . Let $\Delta^\perp$ and $\delta^\perp$ be the maximum, respectively minimum nonzero weight (distance) of the dual code $C^\perp$. Then*

$$\rho < \frac{\log 2(q^{N-K} - 1)}{\log \left[\frac{\sqrt{\Delta^\perp} + \sqrt{\delta^\perp}}{\sqrt{\Delta^\perp} - \sqrt{\delta^\perp}} \right]} + 1.$$

Proof. This follows immediately from Theorem 2.3, using that $\lambda_2 = N(q - 1) - q\delta^\perp$ and $\lambda_n = N(q - 1) - q\Delta^\perp$. $\square$

LEMMA 5.2. *The coset graph G of a linear code C over $GF(q)$ is bipartite if and only if $q = 2$ and $C^\perp$ contains the all-one word.*

Proof. Let N be the length of the code, and λ_n be the smallest eigenvalue of G . Suppose $q = 2$ and $C^\perp$ contains the all-one word, so $\Delta^\perp = N$, then $\lambda_n = -N$, from which we may conclude that G is bipartite. Conversely, if G is bipartite, then $\lambda_n = -N(q - 1)$, so there is an $x \in C^\perp$ such that $N(q - 1) - qw(x) = -N(q - 1)$, which can only be the case if $q = 2$ and $w(x) = N$, so x is the all-one word. $\square$

THEOREM 5.3. *Let C be a binary linear code of length N and dimension K , and covering radius ρ , such that $C^\perp$ contains the all-one word. Let $\delta^\perp$ be the minimum nonzero weight (distance) of the dual code $C^\perp$. Then*

$$\rho < \frac{\log (2^{N-K} - 2)}{\log \left[\frac{\sqrt{N - \delta^\perp} + \sqrt{\delta^\perp}}{\sqrt{N - \delta^\perp} - \sqrt{\delta^\perp}} \right]} + 2.$$

Proof. This follows from Corollary 3.5, using that $\lambda_2 = N - 2\delta^\perp$. $\square$

References

1. A.R. Calderbank and J.-M. Goethals, On a Pair of Dual Subschemes of the Hamming Scheme $H_n(q)$, *Europ. J. Combinatorics* **6** (1985), 133-147.
2. F.R.K. Chung, Diameters and Eigenvalues, *J. Am. Math. Soc.* **2** (1989), 187-196.
3. D.M. Cvetković, M. Doob and H. Sachs, *Spectra of Graphs*, V.E.B. Deutscher Verlag der Wissenschaften, Berlin, 1979.
4. C. Delorme and P. Solé, Diameter, Covering Index, Covering Radius and Eigenvalues, *Europ. J. Combinatorics* **12** (1991), 95-108.
5. W.H. Haemers, *Eigenvalue Techniques in Design and Graph Theory* (thesis Eindhoven University of Technology, 1979), Math. Centre Tract 121, Mathematical Centre, Amsterdam, 1980.
6. A. Lubotzky, R. Phillips and P. Sarnak, Ramanujan Graphs, *Combinatorica* **8** (1988), 261-277.
7. F.J. MacWilliams and N.J.A. Sloane, *The Theory of Error-Correcting Codes*, North-Holland, Amsterdam, 1977.
8. B. Mohar, Eigenvalues, Diameter, and Mean Distance in Graphs, *Graphs Combin.* **7** (1991), 53-64.
9. T.J. Rivlin, *Chebyshev Polynomials (2nd ed.)*, John Wiley and Sons, New York, 1990.

Discussion Paper Series, CentER, Tilburg University, The Netherlands:

(For previous papers please consult previous discussion papers.)

No.	Author(s)	Title
9206	W. Härdle	Applied Nonparametric Models
9207	M. Verbeek and Th. Nijman	Incomplete Panels and Selection Bias: A Survey
9208	W. Härdle and A.B. Tsybakov	How Sensitive Are Average Derivatives?
9209	S. Albæk and P.B. Overgaard	Upstream Pricing and Advertising Signal Downstream Demand
9210	M. Cripps and J. Thomas	Reputation and Commitment in Two-Person Repeated Games
9211	S. Albæk	Endogenous Timing in a Game with Incomplete Information
9212	T.J.A. Storcken and P.H.M. Ruys	Extensions of Choice Behaviour
9213	R.M.W.J. Beetsma and F. van der Ploeg	Exchange Rate Bands and Optimal Monetary Accommodation under a Dirty Float
9214	A. van Soest	Discrete Choice Models of Family Labour Supply
9215	W. Güth and K. Ritzberger	On Durable Goods Monopolies and the (Anti-) Coase-Conjecture
9216	A. Simonovits	Indexation of Pensions in Hungary: A Simple Cohort Model
9217	J.-L. Ferreira, I. Gilboa and M. Maschler	Credible Equilibria in Games with Utilities Changing During the Play
9218	P. Borm, H. Keiding, R. Mclean, S. Oortwijn and S. Tijs	The Compromise Value for NTU-Games
9219	J.L. Horowitz and W. Härdle	Testing a Parametric Model against a Semiparametric Alternative
9220	A.L. Bovenberg	Investment-Promoting Policies in Open Economies: The Importance of Intergenerational and International Distributional Effects
9221	S. Smulders and Th. van de Klundert	Monopolistic Competition, Product Variety and Growth: Chamberlin vs. Schumpeter
9222	H. Bester and E. Petrakis	Price Competition and Advertising in Oligopoly

No.	Author(s)	Title
9223	A. van den Nouweland, M. Maschler and S. Tijs	Monotonic Games are Spanning Network Games
9224	H. Suehiro	A "Mistaken Theories" Refinement
9225	H. Suehiro	Robust Selection of Equilibria
9226	D. Friedman	Economically Applicable Evolutionary Games
9227	E. Bomhoff	Four Econometric Fashions and the Kalman Filter Alternative - A Simulation Study
9228	P. Borm, G.-J. Otten and H. Peters	Core Implementation in Modified Strong and Coalition Proof Nash Equilibria
9229	H.G. Bloemen and A. Kapteyn	The Joint Estimation of a Non-Linear Labour Supply Function and a Wage Equation Using Simulated Response Probabilities
9230	R. Beetsma and F. van der Ploeg	Does Inequality Cause Inflation? - The Political Economy of Inflation, Taxation and Government Debt
9231	G. Almekinders and S. Eijffinger	Daily Bundesbank and Federal Reserve Interventions - Do they Affect the Level and Unexpected Volatility of the DM/\$-Rate?
9232	F. Vella and M. Verbeek	Estimating the Impact of Endogenous Union Choice on Wages Using Panel Data
9233	P. de Bijl and S. Goyal	Technological Change in Markets with Network Externalities
9234	J. Angrist and G. Imbens	Average Causal Response with Variable Treatment Intensity
9235	L. Meijdam, M. van de Ven and H. Verbon	Strategic Decision Making and the Dynamics of Government Debt
9236	H. Houba and A. de Zeeuw	Strategic Bargaining for the Control of a Dynamic System in State-Space Form
9237	A. Cameron and P. Trivedi	Tests of Independence in Parametric Models: With Applications and Illustrations
9238	J.-S. Pischke	Individual Income, Incomplete Information, and Aggregate Consumption
9239	H. Bloemen	A Model of Labour Supply with Job Offer Restrictions
9240	F. Drost and Th. Nijman	Temporal Aggregation of GARCH Processes

No.	Author(s)	Title
9241	R. Gilles, P. Ruys and J. Shou	Coalition Formation in Large Network Economies
9242	P. Kort	The Effects of Marketable Pollution Permits on the Firm's Optimal Investment Policies
9243	A.L. Bovenberg and F. van der Ploeg	Environmental Policy, Public Finance and the Labour Market in a Second-Best World
9244	W.G. Gale and J.K. Scholz	IRAs and Household Saving
9245	A. Bera and P. Ng	Robust Tests for Heteroskedasticity and Autocorrelation Using Score Function
9246	R.T. Baillie, C.F. Chung and M.A. Tieslau	The Long Memory and Variability of Inflation: A Reappraisal of the Friedman Hypothesis
9247	M.A. Tieslau, P. Schmidt and R.T. Baillie	A Generalized Method of Moments Estimator for Long- Memory Processes
9248	K. Wärneryd	Partisanship as Information
9249	H. Huizinga	The Welfare Effects of Individual Retirement Accounts
9250	H.G. Bloemen	Job Search Theory, Labour Supply and Unemployment Duration
9251	S. Eijffinger and E. Schaling	Central Bank Independence: Searching for the Philosophers' Stone
9252	A.L. Bovenberg and R.A. de Mooij	Environmental Taxation and Labor-Market Distortions
9253	A. Lusardi	Permanent Income, Current Income and Consumption: Evidence from Panel Data
9254	R. Beetsma	Imperfect Credibility of the Band and Risk Premia in the European Monetary System
9301	N. Kahana and S. Nitzan	Credibility and Duration of Political Contests and the Extent of Rent Dissipation
9302	W. Güth and S. Nitzan	Are Moral Objections to Free Riding Evolutionarily Stable?
9303	D. Karotkin and S. Nitzan	Some Peculiarities of Group Decision Making in Teams
9304	A. Lusardi	Euler Equations in Micro Data: Merging Data from Two Samples

No.	Author(s)	Title
9305	W. Güth	A Simple Justification of Quantity Competition and the Cournot-Oligopoly Solution
9306	B. Peleg and S. Tijs	The Consistency Principle For Games in Strategic Form
9307	G. Imbens and A. Lancaster	Case Control Studies with Contaminated Controls
9308	T. Ellingsen and K. Wärneryd	Foreign Direct Investment and the Political Economy of Protection
9309	H. Bester	Price Commitment in Search Markets
9310	T. Callan and A. van Soest	Female Labour Supply in Farm Households: Farm and Off-Farm Participation
9311	M. Pradhan and A. van Soest	Formal and Informal Sector Employment in Urban Areas of Bolivia
9312	Th. Nijman and E. Sentana	Marginalization and Contemporaneous Aggregation in Multivariate GARCH Processes
9313	K. Wärneryd	Communication, Complexity, and Evolutionary Stability
9314	O.P. Attanasio and M. Browning	Consumption over the Life Cycle and over the Business Cycle
9315	F. C. Drost and B. J. M. Werker	A Note on Robinson's Test of Independence
9316	H. Hamers, P. Borm and S. Tijs	On Games Corresponding to Sequencing Situations with Ready Times
9317	W. Güth	On Ultimatum Bargaining Experiments - A Personal Review
9318	M.J.G. van Eijs	On the Determination of the Control Parameters of the Optimal Can-order Policy
9319	S. Hurkens	Multi-sided Pre-play Communication by Burning Money
9320	J.J.G. Lemmen and S.C.W. Eijffinger	The Quantity Approach to Financial Integration: The Feldstein-Horioka Criterion Revisited
9321	A.L. Bovenberg and S. Smulders	Environmental Quality and Pollution-saving Technological Change in a Two-sector Endogenous Growth Model
9322	K.-E. Wärneryd	The Will to Save Money: an Essay on Economic Psychology
9323	D. Talman, Y. Yamamoto and Z. Yang	The $(2^{n+m+1} - 2)$ -Ray Algorithm: A New Variable Dimension Simplicial Algorithm For Computing Economic Equilibria on $S^n \times R_+^m$

No.	Author(s)	Title
9324	H. Huizinga	The Financing and Taxation of U.S. Direct Investment Abroad
9325	S.C.W. Eijffinger and E. Schaling	Central Bank Independence: Theory and Evidence
9326	T.C. To	Infant Industry Protection with Learning-by-Doing
9327	J.P.J.F. Scheepens	Bankruptcy Litigation and Optimal Debt Contracts
9328	T.C. To	Tariffs, Rent Extraction and Manipulation of Competition
9329	F. de Jong, T. Nijman and A. Röell	A Comparison of the Cost of Trading French Shares on the Paris Bourse and on SEAQ International
9330	H. Huizinga	The Welfare Effects of Individual Retirement Accounts
9331	H. Huizinga	Time Preference and International Tax Competition
9332	V. Feltkamp, A. Koster, A. van den Nouweland, P. Borm and S. Tijs	Linear Production with Transport of Products, Resources and Technology
9333	B. Lauterbach and U. Ben-Zion	Panic Behavior and the Performance of Circuit Breakers: Empirical Evidence
9334	B. Melenberg and A. van Soest	Semi-parametric Estimation of the Sample Selection Model
9335	A.L. Bovenberg and F. van der Ploeg	Green Policies and Public Finance in a Small Open Economy
9336	E. Schaling	On the Economic Independence of the Central Bank and the Persistence of Inflation
9337	G.-J. Otten	Characterizations of a Game Theoretical Cost Allocation Method
9338	M. Gradstein	Provision of Public Goods With Incomplete Information: Decentralization vs. Central Planning
9339	W. Güth and H. Kliemt	Competition or Co-operation
9340	T.C. To	Export Subsidies and Oligopoly with Switching Costs
9341	A. Demirgüç-Kunt and H. Huizinga	Barriers to Portfolio Investments in Emerging Stock Markets
9342	G.J. Almekinders	Theories on the Scope for Foreign Exchange Market Intervention
9343	E.R. van Dam and W.H. Haemers	Eigenvalues and the Diameter of Graphs

Bibliotheek K. U. Brabant

17 000 01163437 6