Tilburg University

A dynamical characterization of evolutionarily stable states

Bomze, I.M.; van Damme, E.E.C.

Publication date:
1990

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Bomze, I. M., & van Damme, E. E. C. (1990). *A dynamical characterization of evolutionarily stable states*. (CentER Discussion Paper; Vol. 1990-45). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM
CBM
R
8414
1990
45

entER
for
omic Research

Discussion paper

No. 9045

**A DYNAMICAL CHARACTERIZATION
OF EVOLUTIONARILY STABLE STATES** *R20*

by Immanuel M. Bomze *518.55*
and Eric E.C. van Damme

August 1990

ISSN 0924-7815

A dynamical characterization of evolutionarily stable states

June 1990

*Immanuel M. Bomze**

Eric E. C. van Damme†

Abstract. Evolutionary stability, the central solution concept in evolutionary game theory, is closely related to local asymptotic stability in a certain nonlinear dynamical system operating on the state space, the so-called "replicator dynamics". However, a purely dynamical characterization of evolutionary stability is not available in an elementary manner. This characterization can be achieved by investigating so-called "derived games" which consist of mixed strategies corresponding to successful states in the original game. Within this context, several characterization results are obtained, which also may shed some light on the extremality properties of evolutionary stability.

Acknowledgement. The authors thank *Franz Weissing* for stimulating discussions and for drawing attention to the fact that he had done some closely related work in his master thesis [15].

Key words and phrases: asymptotic stability; evolutionary games; mixed strategies; polymorphism; replicator dynamics; stable sets.

Running head: Evolutionary game dynamics.

* Department for Statistics and Computer Science, University of Vienna.

Address: Universitätsstraße 5, A-1010 Wien, Austria

† CentER, Tilburg University.

Address: P.O.Box 90153, NL-5000 LE Tilburg, The Netherlands

1. Introduction

Evolutionary stability, the central solution concept in evolutionary game theory, is closely related to local asymptotic stability in a certain nonlinear dynamical system operating on the state space, the so-called “replicator dynamics”. However, a purely dynamical characterization of evolutionary stability is not available in an elementary manner. This characterization can be achieved by investigating so-called “derived games” which consist of mixed strategies corresponding to successful states in the original game. Within this context, several characterization results are obtained in this paper, which also may shed some light on the extremality properties of evolutionary stability.

The prototypical examples in the seminal papers of *Maynard Smith* and *Price* [5,6] dealt with stability of population states with the aim to explain polymorphism of behaviour (a state where different individuals may behave differently; cf. also the forerunning article [9]). Also, the concept of evolutionary stability seems – at least from the point of view of frequency-dependent selection – to be more stringent in the context of polymorphic states than in models under a monomorphistic interpretation, featuring evolutionarily stable strategies (played by everyone in the population) instead of evolutionarily stable population states. So we start in the present article with the basic, polymorphic model, but the results obtained below may to some extent support the monomorphistic approach, which plays a prominent role in recent literature. To be more precise, the paper is organized as follows: section 1 deals with (possibly) polymorphic populations consisting of individuals that display pure strategies only, while the subsequent sections treat derived games where the original states now play the role of a single (mixed) strategy, and where the dynamical outcomes (i.e. the stable equilibria) are monomorphisms centered at that strategies. In section 2, the concurring states are dimorphisms throughout (only two types of behaviour present), while section 3 deals with a k -morphism setup for the derived games. Section 4 is devoted to weak (*Lya-punov*) stability, and section 5 characterizes evolutionary stability in terms of stable sets, focusing on the convergence of the average strategy played within the k -morphic population in the derived game.

To begin with, let us shortly recapitulate the simplest formal setting of an evolutionary game Γ : assume that, in a certain contest-like situation, individuals are capable to display n different types of behaviour which we for short call strategies in the sequel. With respect to this contest, the state of the population in question is then fully

described by the state space

$$S^n = \{[x_1, \dots, x_n]' \in \mathbb{R}^n : x_i \geq 0, 1 \leq i \leq n, \sum_{i=1}^n x_i = 1\},$$

where x_i represents the relative frequency of individuals displaying strategy i (for short called i -individuals in the sequel) within the population (the symbol $'$ denotes transposition).

There are monomorphic populations (e.g. in state $e = [1, 0, \dots, 0]' \in S^n$) where all individuals display one strategy (e.g. strategy 1), but more interesting are of course polymorphic states (e.g. $[\frac{1}{n}, \dots, \frac{1}{n}]' \in S^n$), where more than one type of behaviour is present (e.g., all with equal frequency). It should be noted that in this setting, every individual can only display one strategy throughout its lifetime, irrespective whether the population is monomorphic or polymorphic.

In this article, we assume that the mean payoff (incremental fitness) to an i -individual contesting with individuals belonging to a population in state $x \in S^n$, depends linearly upon the state x . As shown in [3, pp.61–64], this assumption essentially means that the conflict is pairwise, i.e. there are always exactly two individuals participating in a contest. Therefore, if we denote by a_{ij} the payoff of an i -individual in a confrontation with a j -individual, then

$$A = [a_{ij}]_{1 \leq i \leq n, 1 \leq j \leq n}$$

forms the $n \times n$ payoff matrix which fully characterizes the game Γ . Indeed, the mean payoff to an i -individual contesting with individuals belonging to a population in state $x = [x_1, \dots, x_n]' \in S^n$, amounts to

$$(Ax)_i = \sum_{j=1}^n a_{ij} x_j,$$

and hence the average mean payoff within a (sub-) population in state $y = [y_1, \dots, y_n]' \in S^n$ against a population in state x is given by

$$y'Ax = \sum_{i=1}^n y_i (Ax)_i = \sum_{i=1}^n \sum_{j=1}^n y_i a_{ij} x_j.$$

Now we are in a position to formulate the notion of evolutionary stability introduced in [5] and [6]:

Definition 1:

A state $p \in S^n$ is said to be "evolutionarily stable" in a game Γ , if for all different states $q \in S^n$, $q \neq p$,

$$\text{either } q'Ap < p'Ap \text{ or } q'Ap = p'Ap \text{ and } p'Aq > q'Aq$$

holds.

In other words, a state p is evolutionarily stable if

- the average mean payoff for a population in a different state q against p does not exceed the average mean payoff within the population in state p (equilibrium condition), and
- if, in case of equality in the equilibrium condition, the state q has a lower average mean payoff within itself than p has against q (stability condition).

Following *Taylor and Jonker* [10], we introduce the replicator dynamics $\mathcal{D}(\Gamma)$ corresponding to the game Γ , which operates on the state space S^n (a dot ' denotes differentiation with respect to time t):

$$\dot{x}_i = x_i[(Ax)_i - x'Ax], \quad 1 \leq i \leq n.$$

This dynamical system is one of the simplest models describing the evolution of the population over time: indeed, the amount $(Ax)_i - x'Ax$ by which the mean payoff for i -individuals exceeds – or is exceeded by – the average mean payoff, is considered to be equal to the relative growth – or shrinking – rate $\dot{x}_i/x_i$ of i -individuals. By means of the replicator dynamics we now can introduce a second stability notion as follows:

Definition 2:

A state $p \in S^n$ is said to be "dynamically stable" in a game Γ if and only if p constitutes a locally asymptotically stable fixed point for the replicator dynamics $\mathcal{D}(\Gamma)$.

In other words, a state p is dynamically stable in Γ if every trajectory $x(t)$ that describes the evolution of the population according to $\mathcal{D}(\Gamma)$ tends to p (i.e., $x(t) \rightarrow p$ as $t \rightarrow +\infty$), whenever it started in a state sufficiently close to p (i.e., if the distance from $x(0)$ to p is small enough).

Taylor and Jonker showed for general $n \times n$ -games that every evolutionarily stable state p is dynamically stable [10]. The converse is true for 2×2 -games: here evolutionary and dynamical stability coincide (see, e.g. [2, Thm. 30], or [14, p.226]). On the other hand, some 3×3 -games have dynamically stable equilibria that are not evolutionarily stable, as the following example due to *Zeeman* [17] shows:

Example: For the game Γ given by

$$A = \begin{bmatrix} 0 & 1 & 1 \\ -1 & 0 & 3 \\ 1 & 1 & 0 \end{bmatrix}, \quad p = \begin{bmatrix} 1/3 \\ 1/3 \\ 1/3 \end{bmatrix}$$

is dynamically stable, but not evolutionarily stable, since for $q = [\frac{6}{9}, \frac{1}{9}, \frac{2}{9}]'$ we have $p'Aq = \frac{10}{27} < \frac{32}{81} = q'Aq$.

Remark: A notion stronger than dynamical stability is that of global (dynamical) stability: a state $p \in S^n$ is called "globally stable", if every trajectory $x(t)$ starting in a non-fixed point $x(0) \in S^n$ tends to p as $t \rightarrow +\infty$. One readily sees that global and evolutionary stability are incompatible with each other: indeed, since there are games Γ with more than one evolutionarily stable state, these cannot be globally stable for $\mathcal{D}(\Gamma)$. On the other hand, the state p in the above example is even globally stable though not evolutionarily stable.

2. Mixed strategies; derived 2×2 -games

As the remarks at the end of the previous section show, a purely dynamical characterization of evolutionary stability is not available in a straightforward manner. Nevertheless such a characterization seems to be desirable, and this is the main objective of the present paper. To achieve this, we shall pass from pure strategists' models to mixed strategists' models (see, e.g. [4], [11, 12, 13], [8], and [3, pp.7–15]). Now it is important to distinguish between

- polymorphic (population) states describing the distribution of behaviour within a population, and
- mixed (individual) strategies, which include probabilities π_i for an individual to display behaviour pattern i .

Hence, (polymorphic or monomorphic) states describe a statistical aspect while (mixed or pure) strategies refer to an individualistic aspect of behaviour. The term "probability π_i " may have different interpretations, e.g., π_i may correspond to relative frequencies in sequential contests, or to a change in behaviour with relative duration π_i over lifetime.

A first step towards our goal is done by the following theorem, in which we use the notion of a derived 2×2 -game which is a special case of an evolutionary mixed strategy

game investigated by Sigmund [8] (following his lines, we shall treat the general case of derived $k \times k$ -games below). The rationale for this concept is the idea that individuals belonging to a population in an evolutionarily stable state p , might adopt a mixed strategy in displaying behaviour pattern i with a probability π_i that equals the frequency p_i of i -individuals in state p .

Suppose there is a second state $q \neq p$ that is adopted as a mixed strategy by some other individuals (putting $\pi_i = q_i$) in the same way, perhaps because q is also evolutionarily stable. Recall that there are games that have several evolutionarily stable states (e.g. n in the game with payoff matrix $A = I$, the $n \times n$ -identity matrix); cf. [3, pp.76–81]. Furthermore assume that in this new situation, every individual adopts either p or q as a mixed strategy. Hence all population states considered here are dimorphic: there are only two types of individuals, confrontations between them would best be described in a game $\Gamma_{p,q}$ which we call "derived game", and which has the 2×2 -payoff matrix

$$A_{p,q} = \begin{bmatrix} p'Ap & p'Aq \\ q'Ap & q'Aq \end{bmatrix} = \begin{bmatrix} p' \\ q' \end{bmatrix} A[p, q];$$

indeed, the first row of $A_{p,q}$ is constituted by the payoffs for p , while the second represents the payoffs for q . With respect to $\Gamma_{p,q}$, a state of the population is of course fully described by a vector $\begin{bmatrix} x \\ 1-x \end{bmatrix} \in S^2$ (i.e. $0 \leq x \leq 1$), where x is the relative frequency of p -individuals and $1-x$ is the relative frequency of q -individuals. If $0 < x < 1$, then the state is genuinely dimorphic, while a monomorphic population of p -individuals is given by the state $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$. The next theorem now shows that a population state p (which may be a polymorphism w.r.t. the original game) is evolutionarily stable, if and only if the corresponding monomorphism, based on p played as a mixed strategy, beats any dimorphism based on both p and another mixed strategy arising from a state $q \neq p$ in the original game, provided enough individuals play p . This means that the p -monomorphism $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ is dynamically stable w.r.t. the replicator dynamics $\Gamma_{p,q}$ (see also the remark after Theorem 4 below).

Theorem 1:

A state $p \in S^n$ is evolutionarily stable for Γ if and only if p corresponds to a dynamically stable equilibrium state $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$ in any derived 2×2 -game $\Gamma_{p,q}$ of dimorphic populations playing p and q as mixed strategies, where $q \neq p$ is a different state $q \in S^n$.

Proof: Assume that Γ is represented by the payoff matrix A . Then the payoff matrix

$A_{p,q}$ of the 2×2 -game $\Gamma_{p,q}$ equals

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} p'Ap & p'Aq \\ q'Ap & q'Aq \end{bmatrix}.$$

The corresponding replicator dynamics $\mathcal{D}(\Gamma_{p,q})$ on $S^2 = \{[\frac{x}{1-x}] : 0 \leq x \leq 1\}$ is then given by

$$\dot{x} = x(1-x)[(a-c)x + (b-d)(1-x)],$$

Since $q \neq p$, evolutionary stability of p yields

$$a - c = p'Ap - q'Ap \geq 0$$

and

$$b - d = p'Aq - q'Aq > 0 \quad \text{whenever} \quad a - c = 0.$$

Thus, for x sufficiently close to 1, $\dot{x}$ has to be strictly positive. This implies that the state $[\frac{x}{1-x}]$ evolves towards $[\frac{1}{0}]$ as time goes on, if the initial state is sufficiently close to $[\frac{1}{0}]$. Therefore, the monomorphic state where every individual displays p will be fixed. Assume, conversely, that p is dynamically stable in any game $\Gamma_{p,q}$ with $q \neq p$. Since there are only two possible behavioural patterns, p and q , (local) fixation of p in turn yields $\dot{x} > 0$ if x is close to 1, which entails, by reversion of the above arguments, evolutionary stability of the state p in the original game Γ . $\square$

Example: Let us investigate the example from section 1 in light of the above results: the derived game $\Gamma_{p,q}$ has payoff matrix

$$A_{p,q} = \begin{bmatrix} 2/3 & 10/27 \\ 2/3 & 32/81 \end{bmatrix}.$$

The corresponding replicator dynamics reads

$$\dot{x} = -\frac{2}{81}x(1-x)^2 < 0, \quad 0 < x < 1,$$

so that $x(t) \searrow 0$, i.e. $[\frac{x(t)}{1-x(t)}] \rightarrow [\frac{0}{1}]$ as $t \rightarrow \infty$. Hence the state $[\frac{1}{0}]$ corresponding to p is not dynamically stable in the derived game, although p is dynamically stable in the original game.

3. Characterization in terms of derived $k \times k$ -games

Since the original game can have more than two evolutionarily stable states, it is tempting to consider derived games of k -morphic populations. Here individuals can adopt $k \geq 2$ mixed strategies corresponding, say, to $p_1, \dots, p_k$, where $p_i \in S^n$ describe (possibly) polymorphic states of the "old" population. The state of the "new", k -morphic population comprising these p_i -individuals is then fully described by a vector $x = [x_1, \dots, x_k]' \in S^k$, where x_i is the relative frequency of individuals adopting a mixed strategy adapted to the state p_i . For instance, a monomorphic population consisting exclusively of p_1 -individuals is described by the state $e = [1, 0, \dots, 0]' \in S^k$. The derived game $\Gamma_{p_1, \dots, p_k}$ of k -morphisms has the $k \times k$ -payoff matrix

$$A_{p_1, \dots, p_k} = \begin{bmatrix} p_1' \\ \vdots \\ p_k' \end{bmatrix} A[p_1, \dots, p_k] = C'AC,$$

where C is an $n \times k$ -matrix consisting of the columns $p_1, \dots, p_k$.

Remark: To emphasize that x describes the population state with respect to $\Gamma_{p_1, \dots, p_k}$, one could write $\langle x, p_1, \dots, p_k \rangle$ instead of x . Formally, x corresponds to a statistical distribution (i.e. a probability measure) on S^n of the form

$$\sum_{j=1}^k x_j \delta_{p_j},$$

where δ_p denotes *Dirac measure* (unit point mass) located at p . Models involving states that correspond to more general distributions on S^n are treated in [3]. To avoid lengthy notation, we however shall denote a state simply by x , if it is clear from the context that this is the description of a population with respect to the game $\Gamma_{p_1, \dots, p_k}$.

It is useful to introduce the notion of the "population strategy" m_x which denotes the average strategy adopted within a population in state x :

$$m_x = \sum_{j=1}^k x_j p_j = Cx.$$

In terms of the population strategy, the replicator dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$ on the new state space S^k takes the simple form

$$\dot{x}_i = x_i[p_i' A m_x - m_x' A m_x], \quad 1 \leq i \leq k.$$

Indeed, we have $(C'ACx)_i = p_i'ACx = p_i'Am_x$ and $x'C'ACx = m_x'Am_x$. Of course the dynamics $\mathcal{D}(\Gamma)$ and $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$ are strongly interrelated. For instance, in [8] it is shown that if $\mathcal{D}(\Gamma)$ is a certain type of gradient system, then so is $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$.

The results presented in this and the subsequent section deal with dynamical stability properties of an evolutionarily stable state p in the original game, with respect to the derived k -morphism game $\Gamma_{p_1, \dots, p_k}$, where $p_1 = p$ while p_j , $2 \leq j \leq k$, are different states of the original population. At first let us generalize Theorem 1 (note that the converse of Theorem 2 below follows by Theorem 1):

Theorem 2:

Consider k different states $p_1, \dots, p_k$ in S^n . If p_1 is evolutionarily stable for Γ , and if p_1 is no convex combination of $p_2, \dots, p_k$, then p_1 corresponds to a dynamically stable state e in $\Gamma_{p_1, \dots, p_k}$.

Proof: (1) Let $e = [1, 0, \dots, 0]' \in S^k$ denote the monomorphic state of the mixed strategists' population, where every individual displays strategy p_1 . We at first show that, under the assumption of the theorem, the population strategy m_e determines the state e uniquely. Indeed, suppose that for some $x \in S^k$, $x \neq e$,

$$\sum_{j=1}^k x_j p_j = m_x = m_e = p_1$$

holds. Putting $\lambda_j = x_j / (1 - x_1) \geq 0$, $2 \leq j \leq k$, this implies

$$p_1 = \sum_{j=2}^k \lambda_j p_j,$$

which by assumption is absurd. Hence $m_x \neq m_e = p_1$ if $x \neq e$.

(2) Now we prove $(p_1 - m_x)'Am_x > 0$ whenever $x \neq e$ is close enough to e . Indeed, since the map $x \mapsto m_x$ is continuous, m_x is close to $m_e = p_1$ provided that the distance between x and e is sufficiently small. Evolutionary stability of p_1 in the original game Γ now yields

$$(p_1 - q)'Aq > 0 \quad \text{whenever } q \neq p_1 \text{ is close enough to } p_1 \quad (*)$$

([7]; this result essentially depends upon the fact that an evolutionarily stable state in a pairwise conflict with finitely many pure strategies is uninvadable, cf. [3, p.95]). Thus the claimed assertion follows, putting $q = m_x$.

(3) Finally consider the first equation of the replicator dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$:

$$\dot{x}_1 = x_1(p_1 - m_x)'Am_x > 0$$

whenever $x \neq e$ is close enough to e . Hence $x_1(t)$ increases towards unity if $t \rightarrow +\infty$, implying $x(t) \rightarrow e$ as $t \rightarrow +\infty$. Thus e constitutes a dynamically stable state for the dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$. $\square$

Remark: In (1) and (2) above, we in fact proved that

$$(e - x)' C' A C x = (p_1 - m_x)' A m_x > 0 \quad \text{whenever } x \neq e \text{ is close to } e,$$

which amounts to evolutionary stability of the state e in the derived game $\Gamma_{p_1, \dots, p_k}$. Thus, invoking the result of [10] implying dynamical stability, we obtain an alternative proof of Thm. 2. However, the above proof has the advantage that it admits an immediate generalization adapted to games with nonlinear payoff; see [3, pp. 104 and 109]. Remark further that the condition on p_1 in Theorem 2 above to be no convex combination of the remaining states $p_2, \dots, p_k$, is equivalent to extremality of p_1 in the convex hull of all the states $p_1, \dots, p_k$.

4. Weak dynamical stability

This section deals with the case where the extremality assumption of Theorem 2 is violated. Putting $p_2 = p_1$ one immediately sees that one cannot hope for dynamical stability of p_1 in the derived game under these circumstances; moreover we have the following general negative result:

Theorem 3:

Consider k states $p_1, \dots, p_k$ in S^n , where p_1 is a convex combination of $p_2, \dots, p_k$. Then the state e in $\Gamma_{p_1, \dots, p_k}$ corresponding to p_1 cannot be dynamically stable, even if p_1 were evolutionarily stable for the original game Γ .

Proof: Assume that $p_1 = \sum_{j=2}^k \lambda_j p_j$ for some $\lambda_j \geq 0$, $2 \leq j \leq k$, with $\sum_{j=2}^k \lambda_j = 1$. Put $S = \{x \in S^k : x_1 > 0\}$ and consider the function

$$x \mapsto V(x) = \frac{1}{x_1} \prod_{j=2}^k x_j^{\lambda_j}, \quad x \in S,$$

which is continuous on S and vanishes at $e = [1, 0, \dots, 0]' \in S$. If e were dynamically stable, every trajectory $x(t)$ starting in a neighbourhood of e would fulfill $x(t) \rightarrow e$ and

hence, by continuity, $V(x(t)) \rightarrow V(e) = 0$ as $t \rightarrow +\infty$. If $x(0) = [x_1, \dots, x_k]' \in S^k$ with $x_i > 0$, all i , $1 \leq i \leq k$, then $V(x(0)) > 0$. Therefore we would obtain the relation

$$V(x(t)) < V(x(0)), \quad t \text{ large enough,}$$

which is absurd since V is a constant of motion for the dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$: indeed, taking logarithms and differentiating with respect to time t , we get

$$\begin{aligned} [\log V(x(t))]' &= -\frac{\dot{x}_1(t)}{x_1(t)} + \sum_{j=2}^k \lambda_j \frac{\dot{x}_j(t)}{x_j(t)} \\ &= (m_{x(t)} - p_1)' Am_{x(t)} + \sum_{j=2}^k \lambda_j (p_j - m_{x(t)})' Am_{x(t)} \\ &= (m_{x(t)} - p_1)' Am_{x(t)} + \left(\sum_{j=2}^k \lambda_j p_j \right)' Am_{x(t)} - \sum_{j=2}^k \lambda_j m_{x(t)}' Am_{x(t)} \\ &= (m_{x(t)} - p_1)' Am_{x(t)} + p_1' Am_{x(t)} - m_{x(t)}' Am_{x(t)} = 0. \end{aligned}$$

Hence e cannot be dynamically stable. □

Remark: One could relate the above result to the proof of Theorem 2 in that the population strategy m_x now does not determine the state x uniquely: more precisely, in any neighbourhood of e there are states $x \neq e$ with $m_x = m_e = p_1$ and therefore $\dot{x}_i = x_i(p_i - p_1)' Ap_1 = 0$. Hence e cannot be dynamically stable. The proof above is a variant of an argument of Sigmund [8] who showed that there is a one-dimensional foliation of S^k under the dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$, provided $\{p_1, \dots, p_k\}$ are linearly dependent. However, as Theorem 2 shows, this foliation needs not be incompatible with dynamical stability of e . On the other hand, the existence of such a foliation might suggest that e is weakly dynamically stable in Lyapunov's sense:

Definition 3:

A state $p \in S^n$ is said to be "weakly dynamically stable" in a game Γ if and only if p constitutes a Lyapunov-stable fixed point for the replicator dynamics $\mathcal{D}(\Gamma)$. This means that, given any (small) neighbourhood U of p in S^n , every trajectory starting in S^n , and closely enough to p , will not leave U .

In other words, a state p is weakly dynamically stable in Γ , if for any prescribed (small) barrier $\varepsilon > 0$, every state $x(t)$ will remain within a distance from p smaller than ε , provided that $t \geq 0$ and that the distance from $x(0) \in S^n$ to p is small enough. Again, the trajectory $x(t)$, $t \geq 0$, describes the evolution of the population according to $\mathcal{D}(\Gamma)$.

Theorem 4:

Let $p_1, \dots, p_k$ be k states in S^n . If p_1 is evolutionarily stable for Γ , then p_1 corresponds to a weakly dynamically stable equilibrium in $\Gamma_{p_1, \dots, p_k}$.

Proof: Similarly to the proof of Thm. 2, we derive from evolutionary stability of p_1 that $\dot{x}_1(t) \geq 0$ holds in the dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$, whenever $x(t)$ is close to e , e.g. if $x(t) \in U_\epsilon$, where we put

$$U_\epsilon = \{x \in S^k : x_1 > 1 - \epsilon\},$$

$\epsilon > 0$ being sufficiently small. Since this means that $x_1(t)$ cannot decrease as time $t \geq 0$ increases, provided that $x(0) \in U_\epsilon$, we conclude $x(t) \in U_\epsilon$ for all $t \geq 0$. Because U_ϵ constitutes a neighbourhood of e in S^k , which shrinks to e as $\epsilon \searrow 0$, weak dynamical stability of e is thus established. $\square$

Example: Consider the meanwhile classical hawk-dove game introduced in [5]. If we rescale payoffs to reduce calculational effort, the payoff matrix for this 2×2 -game reads

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

The unique evolutionarily stable state in Γ then is $p = \begin{bmatrix} 1/2 \\ 1/2 \end{bmatrix}$. Indeed, we have

$$x'Ap = 1/2 = p'Ap \quad \text{and} \quad p'Ax = 1/2 > 2x_1(1 - x_1) = x'Ax \quad \text{for all } x \in S^2, x \neq p.$$

Taking $p_1 = p$, $p_2 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $p_3 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, we get

$$C_1 = \begin{bmatrix} 1/2 & 1 & 0 \\ 1/2 & 0 & 1 \end{bmatrix}$$

and hence

$$C_1'AC_1 = \begin{bmatrix} 1/2 & 1/2 & 1/2 \\ 1/2 & 0 & 1 \\ 1/2 & 1 & 0 \end{bmatrix}$$

yielding a dynamics $\mathcal{D}(\Gamma_{p_1, p_2, p_3})$ on S^3 where $e = [1, 0, 0]'$ is *Lyapunov-stable*, but not locally asymptotically stable. By contrast, if we form C_2 stacking p , $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$, and $\begin{bmatrix} 3/4 \\ 1/4 \end{bmatrix}$, together, then the resulting payoff matrix

$$C_2'AC_2 = \begin{bmatrix} 1/2 & 1/2 & 1/2 \\ 1/2 & 0 & 1/4 \\ 1/2 & 1/4 & 3/8 \end{bmatrix}$$

induces a dynamics where e is even globally asymptotically stable (see Figure 1).

Figure 1: Dynamics of two games derived from $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$.

5. Characterization in terms of stable sets

This last section deals with a more general characterization result along lines similar to that followed by *Thomas* in [12, 13]. If one is interested only in the time evolution of observable characteristics like the population strategy $m_{x(t)}$ rather than in the evolution of the states themselves, then the following question becomes important: suppose that $\bar{p} \in S^n$ is an evolutionarily stable state with respect to a game Γ , and consider the derived game $\Gamma_{p_1, \dots, p_k}$, where $\bar{p}$ is a convex combination of the states $p_1, \dots, p_k$; if $x(t)$ denotes a trajectory subject to the dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$, will the population strategy $m_{x(t)}$ converge to $\bar{p}$ or not? In other words, we ask whether or not the states $x(t)$ will approach the set

$$R_{\bar{p}} = \{x \in S^k : m_x = \bar{p}\}$$

as time goes on (observe that the map $x \mapsto m_x = \sum_{j=1}^k x_j p_j$ is uniformly continuous on S^k so that both formulations of the question are indeed equivalent). Using a notion from the theory of dynamical systems, this question amounts to asking whether or not the set $R_{\bar{p}}$ is stable in the following sense:

Definition 4:

Consider a dynamics $\mathcal{D}$ operating on S^k ; a set $R \subseteq S^k$ is said to be "(dynamically) stable", if there is a set $U \subseteq S^k$ which is open in S^k and which contains R , such that every trajectory starting in U approaches R .

In other words, $R_{\bar{p}}$ is dynamically stable if every trajectory $x(t)$ subject to the dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$ with $x(0) \in U$ fulfills $d(x(t), R_{\bar{p}}) \rightarrow 0$ as $t \rightarrow +\infty$, where $U \supseteq R_{\bar{p}}$ is a suitably chosen open set in S^k and

$$d(x, R_{\bar{p}}) = \inf\{\|x - r\| : r \in R_{\bar{p}}\}$$

is the minimum Euclidean distance from x to $R_{\bar{p}}$ ($\|z\| = \sqrt{z'z}$ denotes the usual Euclidean norm of a vector $z \in \mathbb{R}^n$).

The following result shows that the question formulated above can be answered positively, and moreover provides a general dynamical characterization of evolutionary stability that covers all cases discussed in sections 2 and 3.

Theorem 5:

Let $\bar{p} \in S^n$ be a state in the game Γ . Then the following assertions are equivalent:

- (1) $\bar{p}$ is evolutionary stable (with respect to Γ);
- (2) for all derived games $\Gamma_{p_1, \dots, p_k}$, such that $\bar{p}$ is a convex combination of the states $p_1, \dots, p_k$, the set $R_{\bar{p}}$ is dynamically stable with respect to the dynamics $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$.

Proof: (1) $\Rightarrow$ (2): consider an arbitrary state $\bar{x} \in R_{\bar{p}}$, i.e. fulfilling $m_{\bar{x}} = \bar{p}$; we first construct an open neighbourhood $U_{\bar{x}}$ of $\bar{x}$ in S^k that is positively invariant under $\mathcal{D}(\Gamma_{p_1, \dots, p_k})$, i.e. fulfills

$$x(t) \in U_{\bar{x}} \text{ for all } t \geq 0 \text{ whenever } x(0) \in U_{\bar{x}}$$

(similarly to the proof of Theorem 4, where $\bar{x} = e$ and $U_{\bar{x}} = U_e$). To this end we employ evolutionary stability of $\bar{p}$ which guarantees the existence of an $\varepsilon > 0$ such that

$$(\bar{p} - q)' A q > 0 \text{ whenever } 0 < \|q - \bar{p}\| < \varepsilon,$$

see (*) in the proof of Theorem 2. By continuity of the map $x \mapsto m_x$, there is a $\rho > 0$ fulfilling

$$\|m_x - \bar{p}\| = \|m_x - m_{\bar{x}}\| < \varepsilon \text{ whenever } \|x - \bar{x}\| < \rho.$$

Putting these two conditions together, we obtain

$$\text{either } x \in R_{\bar{p}} \text{ or } (\bar{p} - m_x)' A m_x > 0 \text{ if } \|x - \bar{x}\| < \rho.$$

Now consider the function

$$V_{\bar{x}}(x) = \prod_{j=1}^k x_j^{x_j}, \quad x \in S^k;$$

this function is continuous and strictly positive in a neighbourhood of $\bar{x}$ in S^k . Since $x = \bar{x}$ is the only maximizer of $V_{\bar{x}}$ on S^k , there is an $\alpha > 0$ such that

$$\|x - \bar{x}\| < \rho \text{ whenever } V_{\bar{x}}(x) > V_{\bar{x}}(\bar{x}) - \alpha.$$

We claim that

$$U_{\bar{x}} = \{x \in S^k : V_{\bar{x}}(x) > V_{\bar{x}}(\bar{x}) - \frac{\alpha}{2}\}$$

is positively invariant (for simplicity of proof, we choose α smaller than $V_{\bar{x}}(\bar{x})$ so that $V_{\bar{x}}(x) \geq \frac{1}{2} V_{\bar{x}}(\bar{x}) > 0$ for all $x \in U_{\bar{x}}$): indeed, the considerations above ensure the relation

$$(\bar{p} - m_x)' A m_x \geq 0 \text{ for all } x \in U_{\bar{x}};$$

furthermore, the function $V_{\bar{x}}(x(t))$ increases with time t along any trajectory $x(t)$ starting in $U_{\bar{x}}$, because we obtain – similarly to the proof of Theorem 3 –

$$\begin{aligned} [\log V_{\bar{x}}(x(t))] &= \sum_{j=1}^k \bar{x}_j \frac{\dot{x}_j(t)}{x_j(t)} \\ &= \sum_{j=1}^k \bar{x}_j (p_j - m_{x(t)})' Am_{x(t)} \\ &= (m_{\bar{x}} - m_{x(t)})' Am_{x(t)} \\ &= (\bar{p} - m_{x(t)})' Am_{x(t)} \geq 0 \quad \text{if } x(t) \in U_{\bar{x}}, \end{aligned}$$

and hence $x(0) \in U_{\bar{x}}$ entails $x(t) \in U_{\bar{x}}$ for all $t \geq 0$. Moreover, by continuity any accumulation point $\tilde{x} = \lim_{\nu \rightarrow \infty} x(t_{\nu})$ of the trajectory $x(t)$ (where $t_{\nu} \nearrow +\infty$ as $\nu \rightarrow \infty$) fulfills

$$V_{\bar{x}}(\tilde{x}) = \lim_{\nu \rightarrow \infty} V_{\bar{x}}(x(t_{\nu})) \geq \frac{1}{2} V_{\bar{x}}(\tilde{x}) > 0$$

as well as

$$(\bar{p} - m_{\tilde{x}})' Am_{\tilde{x}} = \lim_{\nu \rightarrow \infty} (\bar{p} - m_{x(t_{\nu})})' Am_{x(t_{\nu})} = \lim_{\nu \rightarrow \infty} [\log V_{\bar{x}}(x(t_{\nu}))]' = 0.$$

By construction of $U_{\bar{x}}$, this equality can prevail only if $m_{\tilde{x}} = \bar{p}$, i.e. if $\tilde{x} \in R_{\bar{p}}$. Hence we have shown $d(x(t), R_{\bar{p}}) \rightarrow 0$ as $t \rightarrow +\infty$, whenever $x(0) \in U_{\bar{x}}$. Now observe that the set

$$U = \bigcup_{\bar{x} \in R_{\bar{p}}} U_{\bar{x}} \subseteq S^k$$

is open in S^k and contains $R_{\bar{p}}$. The reasoning above proves $d(x(t), R_{\bar{p}}) \rightarrow 0$ as $t \rightarrow +\infty$ whenever $x(0) \in U$.

(2) $\Rightarrow$ (1) follows from Theorem 1, taking $p_1 = \bar{p}$, $p_2 = q$ (an arbitrary state different from $\bar{p}$), and observing that in this case $R_{\bar{p}} = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$ holds, so that stability of $R_{\bar{p}}$ is the same as asymptotic stability of $\begin{bmatrix} 1 \\ 0 \end{bmatrix}$. $\square$

The proof of (2) $\Rightarrow$ (1) above indicates already why Theorem 5 is a generalization of Theorems 1 and 2: just take $\bar{p} = p_1$, and observe that the extremality condition on p_1 is exactly the relation $R_{\bar{p}} = \{e\}$. Theorem 4 is no direct consequence of Theorem 5, since the latter cannot exclude the case that a trajectory approaches $R_{\bar{p}}$ in an "almost cycling manner", coming eventually arbitrarily close to different states $\tilde{x} \in R_{\bar{p}}$. However, the proof of Theorem 5 also proves Theorem 4, since a trajectory starting in $U_{\bar{x}}$ never leaves it, and because the neighbourhoods $U_{\bar{x}}$ shrink towards $\bar{x}$ as α decreases to zero.

Remark: An alternative approach to Theorem 5 would consist in proceeding similarly as Zeeman in [17], namely considering the dynamics for the population strategies:

$$[m_{\mathbf{x}(t)}]' = [C\mathbf{x}(t)]' = C\dot{\mathbf{x}}(t) = D_{\mathbf{x}(t)}A m_{\mathbf{x}(t)},$$

where $D_{\mathbf{x}} = \sum_{j=1}^k x_j p_j p_j' - m_{\mathbf{x}} m_{\mathbf{x}}'$ is the $n \times n$ -dispersion matrix of the state $\mathbf{x}$, i.e. the variance/covariance-matrix of the distribution $\sum_{j=1}^k x_j \delta_{p_j}$ (cf. the first remark in section 3). The proof in [17, Lemma 7] however seems to work only under additional regularity assumptions both on $\bar{p}$ in the original game and on the dispersion matrices $D_{\mathbf{x}(t)}$ along trajectories $\mathbf{x}(t)$ near $R_{\bar{p}}$. These assumptions are not used in the proof presented above. For instance, Zeeman's method does not apply without modifications to the setting of Theorem 3, if $k \leq n$; more generally, if the rank of C is less than n , then the rank of $D_{\mathbf{x}}$ is necessarily less than $n - 1$ (cf. [1]).

References

- [1] E. Akin, Exponential families and game dynamics. *Can. J. Math.* **34** (1982) 374.
- [2] I.M. Bomze, Non-cooperative two-person games in biology: a classification. *Int. J. Game Theory* **15** (1986) 31.
- [3] I.M. Bomze and B.M. Pötscher, *Game theoretical foundations of evolutionary stability* (Springer, Berlin, 1989).
- [4] W.G.S. Hines, Strategy stability in complex randomly mating diploid populations. *J. appl. Prob.* **19** (1982) 653.
- [5] J. Maynard Smith and G.R. Price, The logic of animal conflict. *Nature (London)* **246** (1973) 15.
- [6] J. Maynard Smith, The theory of games and the evolution of animal conflicts. *J. theor. Biol.* **47** (1974) 209.
- [7] P. Schuster, K. Sigmund, J. Hofbauer and R. Wolff, Selfregulation of behaviour in animal societies, I: symmetric contests. *Biol. Cybern.* **40** (1981) 1.
- [8] K. Sigmund, Game dynamics, mixed strategies, and gradient systems. *Theor. Pop. Biol.* **32** (1987) 114.

- [9] F.M. Stewart, Evolution of dimorphism in a predator-prey model, *Theor. Pop. Biol.* **2** (1971) 493.
- [10] P. Taylor and L. Jonker, Evolutionarily stable strategies and game dynamics. *Math. Biosci.* **40** (1978) 145.
- [11] B. Thomas, Evolutionarily stability: states and strategies. *Theor. Pop. Biol.* **26** (1984) 49.
- [12] B. Thomas, On evolutionarily stable sets. *J. Math. Biol.* **22** (1985) 105.
- [13] B. Thomas, Evolutionarily stable sets in mixed strategist models. *Theor. Pop. Biol.* **28** (1985) 332.
- [14] E.E.C. van Damme, *Stability and perfection of Nash equilibria* (Springer, Berlin, 1987).
- [15] F.J. Weissing, Populationsgenetische Grundlagen der evolutionären Spieltheorie. Thesis, Univ. Bielefeld, FRG (1983).
- [16] E.C. Zeeman, Population dynamics from game theory, in: *Global theory of dynamical systems*, ed. E. Nitecki and C. Robinson (Springer, Berlin, 1980).
- [17] E.C. Zeeman, Dynamics of the evolution of animal conflicts. *J. theor. Biol.* **89** (1981) 249.

Discussion Paper Series, CentER, Tilburg University, The Netherlands:

(For previous papers please consult previous discussion papers.)

No.	Author(s)	Title
8932	E. van Damme, R. Selten and E. Winter	Alternating Bid Bargaining with a Smallest Money Unit
8933	H. Carlsson and E. van Damme	Global Payoff Uncertainty and Risk Dominance
8934	H. Huizinga	National Tax Policies towards Product- Innovating Multinational Enterprises
8935	C. Dang and D. Talman	A New Triangulation of the Unit Simplex for Computing Economic Equilibria
8936	Th. Nijman and M. Verbeek	The Nonresponse Bias in the Analysis of the Determinants of Total Annual Expenditures of Households Based on Panel Data
8937	A.P. Barten	The Estimation of Mixed Demand Systems
8938	G. Marini	Monetary Shocks and the Nominal Interest Rate
8939	W. Güth and E. van Damme	Equilibrium Selection in the Spence Signaling Game
8940	G. Marini and P. Scaramozzino	Monopolistic Competition, Expected Inflation and Contract Length
8941	J.K. Dagsvik	The Generalized Extreme Value Random Utility Model for Continuous Choice
8942	M.F.J. Steel	Weak Exogeneity in Misspecified Sequential Models
8943	A. Roell	Dual Capacity Trading and the Quality of the Market
8944	C. Hsiao	Identification and Estimation of Dichotomous Latent Variables Models Using Panel Data
8945	R.P. Gilles	Equilibrium in a Pure Exchange Economy with an Arbitrary Communication Structure
8946	W.B. MacLeod and J.M. Malcomson	Efficient Specific Investments, Incomplete Contracts, and the Role of Market Alterna- tives
8947	A. van Soest and A. Kapteyn	The Impact of Minimum Wage Regulations on Employment and the Wage Rate Distribution
8948	P. Kooreman and B. Melenberg	Maximum Score Estimation in the Ordered Response Model

No.	Author(s)	Title
8949	C. Dang	The D_3 -Triangulation for Simplicial Deformation Algorithms for Computing Solutions of Nonlinear Equations
8950	M. Cripps	Dealer Behaviour and Price Volatility in Asset Markets
8951	T. Wansbeek and A. Kapteyn	Simple Estimators for Dynamic Panel Data Models with Errors in Variables
8952	Y. Dai, G. van der Laan, D. Talman and Y. Yamamoto	A Simplicial Algorithm for the Nonlinear Stationary Point Problem on an Unbounded Polyhedron
8953	F. van der Ploeg	Risk Aversion, Intertemporal Substitution and Consumption: The CARA-LQ Problem
8954	A. Kapteyn, S. van de Geer, H. van de Stadt and T. Wansbeek	Interdependent Preferences: An Econometric Analysis
8955	L. Zou	Ownership Structure and Efficiency: An Incentive Mechanism Approach
8956	P. Kooreman and A. Kapteyn	On the Empirical Implementation of Some Game Theoretic Models of Household Labor Supply
8957	E. van Damme	Signaling and Forward Induction in a Market Entry Context
9001	A. van Soest, P. Kooreman and A. Kapteyn	Coherency and Regularity of Demand Systems with Equality and Inequality Constraints
9002	J.R. Magnus and B. Pesaran	Forecasting, Misspecification and Unit Roots: The Case of AR(1) Versus ARMA(1,1)
9003	J. Driffill and C. Schultz	Wage Setting and Stabilization Policy in a Game with Renegotiation
9004	M. McAleer, M.H. Pesaran and A. Bera	Alternative Approaches to Testing Non-Nested Models with Autocorrelated Disturbances: An Application to Models of U.S. Unemployment
9005	Th. ten Raa and M.F.J. Steel	A Stochastic Analysis of an Input-Output Model: Comment
9006	M. McAleer and C.R. McKenzie	Keynesian and New Classical Models of Unemployment Revisited
9007	J. Osiewalski and M.F.J. Steel	Semi-Conjugate Prior Densities in Multivariate t Regression Models

No.	Author(s)	Title
9007	J. Osiewalski and M.F.J. Steel	Semi-Conjugate Prior Densities in Multi- variate t Regression Models
9008	G.W. Imbens	Duration Models with Time-Varying Coefficients
9009	G.W. Imbens	An Efficient Method of Moments Estimator for Discrete Choice Models with Choice-Based Sampling
9010	P. Deschamps	Expectations and Intertemporal Separability in an Empirical Model of Consumption and Investment under Uncertainty
9011	W. Güth and E. van Damme	Gorby Games - A Game Theoretic Analysis of Disarmament Campaigns and the Defense Efficiency-Hypothesis
9012	A. Horsley and A. Wrobel	The Existence of an Equilibrium Density for Marginal Cost Prices, and the Solution to the Shifting-Peak Problem
9013	A. Horsley and A. Wrobel	The Closedness of the Free-Disposal Hull of a Production Set
9014	A. Horsley and A. Wrobel	The Continuity of the Equilibrium Price Density: The Case of Symmetric Joint Costs, and a Solution to the Shifting-Pattern Problem
9015	A. van den Elzen, G. van der Laan and D. Talman	An Adjustment Process for an Exchange Economy with Linear Production Technologies
9016	P. Deschamps	On Fractional Demand Systems and Budget Share Positivity
9017	B.J. Christensen and N.M. Kiefer	The Exact Likelihood Function for an Empirical Job Search Model
9018	M. Verbeek and Th. Nijman	Testing for Selectivity Bias in Panel Data Models
9019	J.R. Magnus and B. Pesaran	Evaluation of Moments of Ratios of Quadratic Forms in Normal Variables and Related Statistics
9020	A. Robson	Status, the Distribution of Wealth, Social and Private Attitudes to Risk
9021	J.R. Magnus and B. Pesaran	Evaluation of Moments of Quadratic Forms in Normal Variables

No.	Author(s)	Title
9022	K. Kamiya and D. Talman	Linear Stationary Point Problems
9023	W. Emons	Good Times, Bad Times, and Vertical Upstream Integration
9024	C. Dang	The D_2 -Triangulation for Simplicial Homotopy Algorithms for Computing Solutions of Nonlinear Equations
9025	K. Kamiya and D. Talman	Variable Dimension Simplicial Algorithm for Balanced Games
9026	P. Skott	Efficiency Wages, Mark-Up Pricing and Effective Demand
9027	C. Dang and D. Talman	The D_1 -Triangulation in Simplicial Variable Dimension Algorithms for Computing Solutions of Nonlinear Equations
9028	J. Bai, A.J. Jakeman and M. McAleer	Discrimination Between Nested Two- and Three- Parameter Distributions: An Application to Models of Air Pollution
9029	Th. van de Klundert	Crowding out and the Wealth of Nations
9030	Th. van de Klundert and R. Gradus	Optimal Government Debt under Distortionary Taxation
9031	A. Weber	The Credibility of Monetary Target Announce- ments: An Empirical Evaluation
9032	J. Osiewalski and M. Steel	Robust Bayesian Inference in Elliptical Regression Models
9033	C. R. Wichers	The Linear-Algebraic Structure of Least Squares
9034	C. de Vries	On the Relation between GARCH and Stable Processes
9035	M.R. Baye, D.W. Jansen and Q. Li	Aggregation and the "Random Objective" Justification for Disturbances in Complete Demand Systems
9036	J. Driffill	The Term Structure of Interest Rates: Structural Stability and Macroeconomic Policy Changes in the UK
9037	F. van der Ploeg	Budgetary Aspects of Economic and Monetary Integration in Europe

No.	Author(s)	Title
9038	A. Robson	Existence of Nash Equilibrium in Mixed Strategies for Games where Payoffs Need not Be Continuous in Pure Strategies
9039	A. Robson	An "Informationally Robust Equilibrium" for Two-Person Nonzero-Sum Games
9040	M.R. Baye, G. Tian and J. Zhou	The Existence of Pure-Strategy Nash Equilibrium in Games with Payoffs that are not Quasiconcave
9041	M. Burnovsky and I. Zang	"Costless" Indirect Regulation of Monopolies with Substantial Entry Cost
9042	P.J. Deschamps	Joint Tests for Regularity and Autocorrelation in Allocation Systems
9043	S. Chib, J. Osiewalski and M. Steel	Posterior Inference on the Degrees of Freedom Parameter in Multivariate-t Regression Models
9044	H.A. Keuzenkamp	The Probability Approach in Economic Methodology: On the Relation between Haavelmo's Legacy and the Methodology of Economics
9045	I.M. Bomze and E.E.C. van Damme	A Dynamical Characterization of Evolutionarily Stable States

P.O. BOX 90153. 5000 LE TILBURG. THE NETHERLANDS

Bibliotheek K. U. Brabant

17 000 01117581 8