Tilburg University

Sensitivity to prior independence via Farlie-Gumbel-Morgenstern model

de La Horra, J.; Fernández, C.

Publication date:
1994

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

de La Horra, J., & Fernández, C. (1994). *Sensitivity to prior independence via Farlie-Gumbel-Morgenstern model*. (CentER Discussion Paper; Vol. 1994-17). CentER.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

R
CBM
841
R
1994-17
8414
1994
NR.17

entER
for
omic Research

Discussion paper

Center
for
Economic Research

8414
1994
17
No. 9417
44

**SENSITIVITY TO PRIOR
INDEPENDENCE VIA FARLIE-GUMBEL-
MORGENSTERN MODEL**

by Julian de la Horra and
Carmen Fernandez

February 1994

ISSN 0924-7815

Sensitivity to prior independence via Farlie-Gumbel-Morgenstern model

By JULIAN DE LA HORRA and CARMEN FERNANDEZ

Departamento de Matemáticas
Universidad Autónoma de Madrid
28049 Madrid, Spain

Key Words and Phrases: Bayesian robustness; Farlie-Gumbel-Morgenstern system; Prior independence; Posterior bounds.

Abstract

When the parameter space is multidimensional, the prior marginals are sometimes simple to specify, whereas the joint distribution may be extremely hard to elicit; as a consequence of this, prior independence is often assumed. When this is the case, it seems convenient to perform a sensitivity analysis, i.e., to consider a class of prior distributions (containing the independence as a particular case) and to compute the range of variation of the posterior quantity of interest, when the prior moves in that class. In this paper, the well-known Farlie-Gumbel-Morgenstern system (or a generalization of it) is proposed as the class of priors. We show that the robustness analysis is easy to carry out and, although this family may sometimes seem too narrow, it could be used as a first step in a robustness study.

1. Introduction

Let x be an observation from a sampling model $l(x|\theta)$, where $\theta = (\theta_1, \dots, \theta_n) \in \Theta = \Theta_1 \times \dots \times \Theta_n$. In order to perform a Bayesian analysis we need to specify a prior density $\pi(\theta_1, \dots, \theta_n)$ on the parameters. The posterior density will then take the form

$$\pi(\theta_1, \dots, \theta_n|x) \propto l(x|\theta_1, \dots, \theta_n)\pi(\theta_1, \dots, \theta_n).$$

Suppose that it is relatively simple to elicit our prior marginal densities $g_1(\theta_1), \dots, g_n(\theta_n)$. If the parameters could be considered independent, the prior density would be $\pi(\theta_1, \dots, \theta_n) = \prod_{i=1}^n g_i(\theta_i)$. However, in some cases we may not have a strong belief in the latter assumption and we will want to carry out a study of robustness with respect to departures from prior independence. For this purpose it will be convenient to consider a class of prior densities, all having the specified marginals $g_1(\theta_1), \dots, g_n(\theta_n)$, while containing the density under independence as a particular case, and to compute the ranges of the posterior quantities of interest as the prior moves in that class.

The paper by Lavine, Wasserman and Wolpert (1991) follows these lines. They consider an ϵ -contamination class where the contaminations allowed are all probability distributions with the given marginals. Although their approach is very interesting, it presents some inconveniences: in some cases the class may be too large, containing distributions that are not reasonable in view of our prior beliefs, resulting into a misleading impression of lack of robustness. Furthermore, in order to solve the problem, they need to simplify it, considering a finite partition for each component of the parameter space and assigning the probability corresponding to each set of the partition induced on Θ , to a single point in that set. Moreno and Cano (1992) propose other classes of prior distributions that could also be used as approximations to this ϵ -contamination class. Other studies of Bayesian robustness in multidimensional parameter spaces can be found in Lavine (1991a), and Berger and Moreno (1992).

Families of multivariate distributions with fixed marginals have been investigated in the literature for a long time. See, for instance, Mardia (1970) and Whitt (1976). There are well-known, simple models that, keeping the marginals fixed, allow for different degrees of dependence between the components. It seems quite reasonable to use some of these models as classes of prior distributions to analyze robustness with respect to departures from the assumption of prior independence. In particular, the Farlie-Gumbel-Morgenstern (FGM) family is very simple and could be appropriate for our purposes. It will be described in the next section. As a matter of fact, the FGM system is a particular case of a model with a very clear and intuitive meaning. This more general model will be introduced in Section 3. Ranges of posterior expectations as the prior moves over this class are computed in Section 4. Finally, some examples are analyzed in Section 5.

2. The Farlie-Gumbel-Morgenstern model

The FGM system was discussed by Morgenstern (1956), Gumbel (1958) and Farlie (1960); see also Johnson and Kotz (1975, 1977). This model is described next. For notational convenience, we shall restrict ourselves to the bidimensional case; however, the FGM system and the results in this paper can easily be extended to higher dimensions.

Let $F(\theta_1)$ and $G(\theta_2)$ be fixed marginal distribution functions. The Farlie-Gumbel-Morgenstern model corresponds to the following distribution function:

$$\Pi(\theta_1, \theta_2) = F(\theta_1)G(\theta_2)[1 + \lambda\{1 - F(\theta_1)\}\{1 - G(\theta_2)\}], \quad \text{where } \lambda \in [-1, 1].$$

(Of course, $\Pi(\theta_1, \theta_2)$ depends on λ but, for the sake of simplicity, this dependence will not be explicitly expressed.)

If $F(\theta_1)$ and $G(\theta_2)$ have density functions $f(\theta_1)$ and $g(\theta_2)$ respectively, the density corresponding to $\Pi(\theta_1, \theta_2)$ is easily shown to be

$$\pi(\theta_1, \theta_2) = f(\theta_1)g(\theta_2)[1 + \lambda\{1 - 2F(\theta_1)\}\{1 - 2G(\theta_2)\}].$$

Notice that for $\lambda = 0$ we obtain $\pi(\theta_1, \theta_2) = f(\theta_1)g(\theta_2)$, that is, θ_1 and θ_2 are independent. The cases $\lambda = 1$ and $\lambda = -1$ represent the maximum degrees of positive and negative

dependence, respectively, allowed in this model. In our Bayesian analysis this translates into prior beliefs about the parameters. An advantage of the FGM system is that only moderate departures from independence are permitted. We next give two examples of this assertion (see Johnson and Kotz (1977)):

- a) If the marginals are $N(0, 1)$, the correlation is $\lambda\pi^{-1}$ (so it ranges from -0.318 to 0.318).
- b) If the marginals are uniform distributions over $(0, 1)$, the correlation is $\lambda/3$ (so it ranges from -0.333 to 0.333).

The FGM model can also be expressed as follows.

For $\lambda \in [0, 1]$:

$$\pi(\theta_1, \theta_2) = (1 - \lambda)f(\theta_1)g(\theta_2) + \lambda f(\theta_1)g(\theta_2)[1 + \{1 - 2F(\theta_1)\}\{1 - 2G(\theta_2)\}],$$

and for $\lambda \in [-1, 0]$:

$$\begin{aligned} \pi(\theta_1, \theta_2) &= (1 + \lambda)f(\theta_1)g(\theta_2) - \lambda f(\theta_1)g(\theta_2)[1 - \{1 - 2F(\theta_1)\}\{1 - 2G(\theta_2)\}] \\ &= (1 - \mu)f(\theta_1)g(\theta_2) + \mu f(\theta_1)g(\theta_2)[1 - \{1 - 2F(\theta_1)\}\{1 - 2G(\theta_2)\}] \end{aligned}$$

where $\mu = -\lambda \in [0, 1]$.

This representation naturally suggests the generalization that we describe in the next section.

3. A more general model

Let us assume that in the process of prior elicitation we come up with the marginal densities $f(\theta_1)$ and $g(\theta_2)$ but we are not entirely comfortable with the assumption of prior independence between θ_1 and θ_2 . In this case, a sensible and very simple class of prior densities is

$$\Gamma = \Gamma^+ \cup \Gamma^- \tag{3.1}$$

where

$$\begin{aligned} \Gamma^+ &= \{ \pi(\theta_1, \theta_2) = (1 - \lambda)\pi_I(\theta_1, \theta_2) + \lambda\pi^+(\theta_1, \theta_2), \lambda \in [0, 1] \}, \\ \Gamma^- &= \{ \pi(\theta_1, \theta_2) = (1 - \lambda)\pi_I(\theta_1, \theta_2) + \lambda\pi^-(\theta_1, \theta_2), \lambda \in [0, 1] \}, \end{aligned} \tag{3.2}$$

$\pi_I(\theta_1, \theta_2) = f(\theta_1)g(\theta_2)$ is the density obtained under independence and $\pi^+(\theta_1, \theta_2)$ and $\pi^-(\theta_1, \theta_2)$ are fixed densities with marginals $f(\theta_1)$ and $g(\theta_2)$, representing some degree of positive and negative dependence, respectively.

Obviously, the FGM model is a special case of this one, with

$$\begin{aligned} \pi^+(\theta_1, \theta_2) &= f(\theta_1)g(\theta_2)[1 + \{1 - 2F(\theta_1)\}\{1 - 2G(\theta_2)\}], \\ \pi^-(\theta_1, \theta_2) &= f(\theta_1)g(\theta_2)[1 - \{1 - 2F(\theta_1)\}\{1 - 2G(\theta_2)\}]. \end{aligned}$$

The class Γ^+ is quite similar to the class of distributions considered in De la Horra and Ruiz-Rivas (1988).

It is easy to see that for $\lambda = 0$ we are in the case of prior independence between θ_1 and θ_2 . As λ increases, we obtain a bigger degree of dependence, always keeping the marginal densities fixed at $f(\theta_1)$ and $g(\theta_2)$. The maximum degree of dependence allowed for is controlled by the choice of π^+ and π^- .

As we have already pointed out, the class defined in (3.1) and (3.2) is very adequate for studies of robustness to departures from the hypothesis of prior independence. Its main advantage is the simplicity of elicitation and computation. In the next section we provide analytical results for the supremum and the infimum of the posterior expectation of a function of interest, when the prior ranges over this class. This greatly alleviates the robustness analysis, which typically is computationally very demanding. We think this is an important point in favour of this class (see Example 5.1 below).

On the other hand, in some cases we may find that this class is too narrow, leaving out densities that we may not wish to discard a priori. Even when that is the case, the simplicity of its analysis still makes this class worth to be used at a first step of a robustness study. If we find absence of robustness with this class, the same conclusion would be obtained for a wider class of priors (see Example 5.2).

4. Sensitivity of posterior quantities

Consider the class of priors Γ introduced in (3.1) and (3.2). We are interested in the sensitivity of the posterior expectation of some function, $h(\theta_1, \theta_2)$, as the prior ranges over that class. We therefore need to obtain

$$\sup_{\pi \in \Gamma} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta \quad \text{and} \quad \inf_{\pi \in \Gamma} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta. \quad (4.1)$$

As Γ in (3.1) is the union of Γ^+ and Γ^- , the quantities in (4.1) are equal to

$$\begin{aligned} & \max \left\{ \sup_{\pi \in \Gamma^+} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta, \sup_{\pi \in \Gamma^-} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta \right\} \\ & \text{and} \quad \min \left\{ \inf_{\pi \in \Gamma^+} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta, \inf_{\pi \in \Gamma^-} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta \right\}, \end{aligned} \quad (4.2)$$

respectively.

Some common choices for $h(\theta)$ are $h(\theta_1, \theta_2) = \theta_1$, for studying the posterior mean of the parameter of interest, and $h(\theta_1, \theta_2) = I_C(\theta_1, \theta_2)$, for the posterior probability of a set C of interest.

In the next theorem, we give the supremum and the infimum of the posterior expectation of $h(\theta)$ when the prior $\pi(\theta)$ ranges in Γ^+ . Similar results can be obtained for Γ^- changing π^+ by π^- .

Theorem. *With Γ^+ defined as in (3.2), if*

$$\frac{\int_{\Theta} h(\theta) l(x|\theta) \pi^+(\theta) d\theta}{\int_{\Theta} l(x|\theta) \pi^+(\theta) d\theta} \geq \frac{\int_{\Theta} h(\theta) l(x|\theta) \pi_I(\theta) d\theta}{\int_{\Theta} l(x|\theta) \pi_I(\theta) d\theta},$$

then

$$\sup_{\pi \in \Gamma^+} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta = \frac{\int_{\Theta} h(\theta) l(x|\theta) \pi^+(\theta) d\theta}{\int_{\Theta} l(x|\theta) \pi^+(\theta) d\theta}$$

and

$$\inf_{\pi \in \Gamma^+} \int_{\Theta} h(\theta) \pi(\theta|x) d\theta = \frac{\int_{\Theta} h(\theta) l(x|\theta) \pi_I(\theta) d\theta}{\int_{\Theta} l(x|\theta) \pi_I(\theta) d\theta}.$$

If the inequality sign is reversed, the roles of the infimum and the supremum are interchanged.

Proof: For $\pi \in \Gamma^+$,

$$\int_{\Theta} h(\theta) \pi(\theta|x) d\theta = \frac{(1-\lambda) \int_{\Theta} h(\theta) l(x|\theta) \pi_I(\theta) d\theta + \lambda \int_{\Theta} h(\theta) l(x|\theta) \pi^+(\theta) d\theta}{(1-\lambda) \int_{\Theta} l(x|\theta) \pi_I(\theta) d\theta + \lambda \int_{\Theta} l(x|\theta) \pi^+(\theta) d\theta},$$

where $\lambda \in [0, 1]$.

Its derivative with respect to λ is

$$\frac{\int_{\Theta} h(\theta) l(x|\theta) \pi^+(\theta) d\theta \int_{\Theta} l(x|\theta) \pi_I(\theta) d\theta - \int_{\Theta} h(\theta) l(x|\theta) \pi_I(\theta) d\theta \int_{\Theta} l(x|\theta) \pi^+(\theta) d\theta}{\left\{ (1-\lambda) \int_{\Theta} l(x|\theta) \pi_I(\theta) d\theta + \lambda \int_{\Theta} l(x|\theta) \pi^+(\theta) d\theta \right\}^2}.$$

If

$$\frac{\int_{\Theta} h(\theta) l(x|\theta) \pi^+(\theta) d\theta}{\int_{\Theta} l(x|\theta) \pi^+(\theta) d\theta} \geq \frac{\int_{\Theta} h(\theta) l(x|\theta) \pi_I(\theta) d\theta}{\int_{\Theta} l(x|\theta) \pi_I(\theta) d\theta},$$

the derivative is nonnegative and the result is obtained. ♣

Needless to say, the above result could also be derived by applying the elegant linearization technique (see Lavine (1991b)), which turns the nonlinear problem of optimizing a posterior expectation into a linear one; but this technique is obviously not necessary here.

The theorem together with the statement in (4.2) imply that our calculations will be reduced to evaluating the integrals

$$\int_{\Theta} h(\theta) l(x|\theta) \pi_I(\theta) d\theta, \int_{\Theta} h(\theta) l(x|\theta) \pi^+(\theta) d\theta, \int_{\Theta} h(\theta) l(x|\theta) \pi^-(\theta) d\theta, \\ \int_{\Theta} l(x|\theta) \pi_I(\theta) d\theta, \int_{\Theta} l(x|\theta) \pi^+(\theta) d\theta \text{ and } \int_{\Theta} l(x|\theta) \pi^-(\theta) d\theta.$$

These evaluations will very often have to be done by numerical methods.

We would like to remark that the FGM model is even simpler, since it can be expressed as a function of a unique parameter

$$\pi(\theta_1, \theta_2) = f(\theta_1) g(\theta_2) [1 + \lambda \{1 - 2F(\theta_1)\} \{1 - 2G(\theta_2)\}], \quad \lambda \in [-1, 1].$$

In this case, the extrema are attained at $\lambda = -1$ and $\lambda = 1$, which correspond to π^- and π^+ , respectively, and only four integrals need to be computed.

In the following section we use this class in two robustness analyses.

5. Examples

Example 5.1: ECMO data

A clinical trial was designed to compare the performance of ECMO (extracorporeal membrane oxygenation) with that of the standard therapy in the treatment of a respiratory disease in infants (see Ware (1989)). Of nine patients that were given ECMO all nine survived, whereas of ten patients in the control group, only six survived.

Let p_1 and p_2 be the probabilities of survival under the standard treatment and ECMO, respectively. We reparameterize in terms of (δ, γ) , where $\delta = \eta_2 - \eta_1$, $\gamma = (\eta_1 + \eta_2)/2$ and $\eta_i = \log\{p_i/(1 - p_i)\}$, $i=1, 2$. The goal is then to determine whether $\delta > 0$ (or equivalently $p_2 > p_1$), γ being a nuisance parameter. Therefore we will focus on the posterior probability $P(\delta > 0|data)$.

Kass and Greenhouse (1989) carry out a robust analysis of this quantity by specifying 42 different prior distributions on (δ, η_1) and on (δ, γ) , assuming in all cases prior independence. The prior marginal distributions favoured in this study are a Cauchy(0, 1.099²) for δ and a Cauchy(0, 0.419²) for γ . This choice is made on the basis of the assumption that randomization is ethically justifiable and some historical control information. Under this independent prior, it is obtained that $P(\delta > 0|data) \approx 0.94$. Other studies of sensitivity of this posterior quantity to the specification of the prior distribution can be found in Lavine (1991a), Lavine, Wasserman and Wolpert (1991), Berger and Moreno (1992) and Moreno and Cano (1992). They all find small ranges of variation of the posterior probability of ECMO being superior to the standard therapy, as long as the class of priors is not unreasonably large.

In this section, we shall use the FGM system to check for robustness to departures from prior independence, keeping the marginal distributions fixed at Cauchy(0, 1.099²) for δ and Cauchy(0, 0.419²) for γ . According to the previous section, the extrema of $P(\delta > 0|data)$ will take the form

$$\int_{\mathbb{R}^2} h(\delta, \gamma) \pi^+(\delta, \gamma|data) d\delta d\gamma \quad \text{and} \quad \int_{\mathbb{R}^2} h(\delta, \gamma) \pi^-(\delta, \gamma|data) d\delta d\gamma,$$

where

$$\begin{aligned} h(\delta, \gamma) &= I_{(0, +\infty)}(\delta), \\ \pi^+(\delta, \gamma) &= f(\delta)g(\gamma)[1 + \{1 - 2F(\delta)\}\{1 - 2G(\gamma)\}], \\ \pi^-(\delta, \gamma) &= f(\delta)g(\gamma)[1 - \{1 - 2F(\delta)\}\{1 - 2G(\gamma)\}], \end{aligned}$$

f, g are the density functions and F, G are the distribution functions of the fixed marginal Cauchy distributions.

Evaluating these integrals through Monte Carlo importance sampling gives

$$\int_{\mathbb{R}^2} h(\delta, \gamma) \pi^+(\delta, \gamma|data) d\delta d\gamma \approx 0.96, \quad \int_{\mathbb{R}^2} h(\delta, \gamma) \pi^-(\delta, \gamma|data) d\delta d\gamma \approx 0.89,$$

from which we conclude that the posterior probability of ECMO being superior to the standard treatment is reasonably robust when the prior ranges over the FGM class. Of course this is not a surprising finding in view of the previous literature, but it is included

here with the purpose of illustrating the performance of our method with a well-known data set.

It is worth mentioning that although the FGM class is parametric, the bounds that we find here are not necessarily narrower than those obtained when using a more complicated, non-parametric class. For instance, the extreme values found in this particular example, are rather close to the ones in Lavine, Wasserman and Wolpert (1991) when ε lies in a neighborhood of 0.25.

Example 5.2:

Let $(0.52, 0)$, $(4.95, 0.03)$, $(0.1, 0.05)$, $(-3.43, -0.01)$ and $(3.15, -0.11)$ be an independent sample from a model with density

$$l(x_1, x_2 | \theta_1, \theta_2) = \frac{1}{\pi \{1 + (x_1 - \theta_1)^2\}} \frac{1}{\sqrt{0.02\pi}} e^{-(x_2 - \theta_2)^2 / 0.02},$$

that is, conditionally upon $\theta = (\theta_1, \theta_2)$, x_1 and x_2 are drawn independently from a Cauchy($\theta_1, 1$) and a Normal($\theta_2, 0.1^2$), respectively.

We wish to test the hypothesis $H_0 : \theta_1 < 1.25$ against $H_1 : \theta_1 > 1.25$. Thus we are interested in finding the posterior odds $P(H_0 | \text{data}) / P(H_1 | \text{data})$.

The marginal prior for θ_1 is a mixture of two Gamma distributions, leading to the density function

$$f(\theta_1) = \left(\frac{1}{2} \{ (0.25)^{-2} e^{-\theta_1 / 0.25} \theta_1 \} + \frac{1}{2} \left\{ \frac{(0.25)^{-10}}{9!} e^{-\theta_1 / 0.25} \theta_1^9 \right\} \right) I_{(0, \infty)}(\theta_1),$$

whereas the marginal distribution for θ_2 is Exponential with density

$$g(\theta_2) = 2 e^{-2\theta_2} I_{(0, \infty)}(\theta_2).$$

Observe that the prior chosen for θ_1 leads to $P(H_0) / P(H_1) = 1$, therefore representing indifference among the hypotheses.

Under the independent prior, $\pi_I(\theta_1, \theta_2) = f(\theta_1)g(\theta_2)$, the posterior odds are 3.17. If we now analyze sensitivity to departures from independence by embedding $\pi_I(\theta_1, \theta_2)$ in a FGM class, results in Section 4 tell us that the extrema of this posterior quantity will be attained for the priors π^+ and π^- . For π^+ it is found that $P(H_0 | \text{data}) / P(H_1 | \text{data}) \approx 6.14$, which strongly favours the null hypothesis, whereas for π^- , $P(H_0 | \text{data}) / P(H_1 | \text{data}) \approx 1.22$, which essentially does not favour any of the hypotheses. This range of variation seems too big to proceed confidently. Therefore, in this case, absence of robustness has been detected by means of a simple analysis.

Acknowledgements

This research has been supported by the D.G.I.C.Y.T. under grant number PB91-0014. We are grateful to Carmen Ruiz-Rivas, Antonio Cuevas, Mark F.J. Steel and Bas Werker

for many interesting comments. The second author acknowledges the hospitality of the Department of Econometrics, Tilburg University.

Bibliography

- Berger, J.O. and Moreno, E. (1992). "Bayesian robustness in bidimensional models: prior independence". *J. Statist. Plan. Inf.*, forthcoming.
- De la Horra, J. and Ruiz-Rivas, C. (1988). "A Bayesian method for inferring the degree of dependence for a positively quadrant dependent distribution". *Commun. Statist. A* 17, 4357-4370.
- Farlie, D.J.G. (1960). "The performance of some correlation coefficients for a general bivariate distribution". *Biometrika* 47, 307-323.
- Gumbel, E.J. (1958). "Distributions à plusieurs variables dont les marges sont donnés". *C. R. Acad. Sci.*, Paris 246, 2717-2719.
- Johnson, N.L. and Kotz, S. (1975). "On some generalized Farlie-Gumbel-Morgenstern distributions". *Commun. Statist. A* 4, 415-427.
- Johnson, N.L. and Kotz, S. (1977). "On some generalized Farlie-Gumbel-Morgenstern distributions II: regression, correlation and further generalizations". *Commun. Statist. A* 6, 485-496.
- Kass, R.E. and Greenhouse, J.B. (1989). "Investigating therapies of potentially great benefit: a Bayesian perspective". Comment on "Investigating therapies of potentially great benefit: ECMO", by J.H. Ware, *Statistical Science* 4, 310-317.
- Lavine, M. (1991a). "An approach to robust Bayesian analysis for multidimensional parameter spaces". *J.A.S.A.* 86, 400-403.
- Lavine, M. (1991b). "Sensitivity in Bayesian Statistics: the prior and the likelihood". *J.A.S.A.* 86, 396-399.
- Lavine, M., Wasserman, L. and Wolpert, R.L. (1991). "Bayesian inference with specified prior marginals". *J.A.S.A.* 86, 964-971.
- Mardia, K.V. (1970). *Families of bivariate distributions*. Charles W. Griffin, London.
- Moreno, E. and Cano, J.A. (1992). "Classes of bidimensional priors specified on a collection of sets: Bayesian robustness". Technical report, Universidad de Granada.

- Morgenstern, D. (1956). "Einfache beispiele zweidimensionaler verteilungen". *Mitt. Math. Statistik* 8, 234-235.
- Ware, J.H. (1989). "Investigating therapies of potentially great benefit: ECMO". *Statistical Science* 4, 298-306.
- Whitt, W. (1976). "Bivariate distributions with given marginals". *Ann. Statist.* 4, 1280-1289.

Discussion Paper Series, CentER, Tilburg University, The Netherlands:

(For previous papers please consult previous discussion papers.)

No.	Author(s)	Title
9310	T. Callan and A. van Soest	Female Labour Supply in Farm Households: Farm and Off-Farm Participation
9311	M. Pradhan and A. van Soest	Formal and Informal Sector Employment in Urban Areas of Bolivia
9312	Th. Nijman and E. Sentana	Marginalization and Contemporaneous Aggregation in Multivariate GARCH Processes
9313	K. Wärneryd	Communication, Complexity, and Evolutionary Stability
9314	O.P. Attanasio and M. Browning	Consumption over the Life Cycle and over the Business Cycle
9315	F. C. Drost and B. J. M. Werker	A Note on Robinson's Test of Independence
9316	H. Hamers, P. Borm and S. Tijs	On Games Corresponding to Sequencing Situations with Ready Times
9317	W. Güth	On Ultimatum Bargaining Experiments - A Personal Review
9318	M.J.G. van Eijs	On the Determination of the Control Parameters of the Optimal Can-order Policy
9319	S. Hurkens	Multi-sided Pre-play Communication by Burning Money
9320	J.J.G. Lemmen and S.C.W. Eijffinger	The Quantity Approach to Financial Integration: The Feldstein-Horioka Criterion Revisited
9321	A.L. Bovenberg and S. Smulders	Environmental Quality and Pollution-saving Technological Change in a Two-sector Endogenous Growth Model
9322	K.-E. Wärneryd	The Will to Save Money: an Essay on Economic Psychology
9323	D. Talman, Y. Yamamoto and Z. Yang	The $(2^{n+m+1} - 2)$ -Ray Algorithm: A New Variable Dimension Simplicial Algorithm For Computing Economic Equilibria on $S^n \times R_+^m$
9324	H. Huizinga	The Financing and Taxation of U.S. Direct Investment Abroad
9325	S.C.W. Eijffinger and E. Schaling	Central Bank Independence: Theory and Evidence
9326	T.C. To	Infant Industry Protection with Learning-by-Doing
9327	J.P.J.F. Scheepens	Bankruptcy Litigation and Optimal Debt Contracts

No.	Author(s)	Title
9328	T.C. To	Tariffs, Rent Extraction and Manipulation of Competition
9329	F. de Jong, T. Nijman and A. Röell	A Comparison of the Cost of Trading French Shares on the Paris Bourse and on SEAQ International
9330	H. Huizinga	The Welfare Effects of Individual Retirement Accounts
9331	H. Huizinga	Time Preference and International Tax Competition
9332	V. Feltkamp, A. Koster, A. van den Nouweland, P. Borm and S. Tijs	Linear Production with Transport of Products, Resources and Technology
9333	B. Lauterbach and U. Ben-Zion	Panic Behavior and the Performance of Circuit Breakers: Empirical Evidence
9334	B. Melenberg and A. van Soest	Semi-parametric Estimation of the Sample Selection Model
9335	A.L. Bovenberg and F. van der Ploeg	Green Policies and Public Finance in a Small Open Economy
9336	E. Schaling	On the Economic Independence of the Central Bank and the Persistence of Inflation
9337	G.-J. Otten	Characterizations of a Game Theoretical Cost Allocation Method
9338	M. Gradstein	Provision of Public Goods With Incomplete Information: Decentralization vs. Central Planning
9339	W. Güth and H. Kliemt	Competition or Co-operation
9340	T.C. To	Export Subsidies and Oligopoly with Switching Costs
9341	A. Demirgüç-Kunt and H. Huizinga	Barriers to Portfolio Investments in Emerging Stock Markets
9342	G.J. Almekinders	Theories on the Scope for Foreign Exchange Market Intervention
9343	E.R. van Dam and W.H. Haemers	Eigenvalues and the Diameter of Graphs
9344	H. Carlsson and S. Dasgupta	Noise-Proof Equilibria in Signaling Games
9345	F. van der Ploeg and A.L. Bovenberg	Environmental Policy, Public Goods and the Marginal Cost of Public Funds
9346	J.P.C. Blanc and R.D. van der Mei	The Power-series Algorithm Applied to Polling Systems with a Dormant Server
9347	J.P.C. Blanc	Performance Analysis and Optimization with the Power- series Algorithm

No.	Author(s)	Title
9348	R.M.W.J. Beetsma and F. van der Ploeg	Intramarginal Interventions, Bands and the Pattern of EMS Exchange Rate Distributions
9349	A. Simonovits	Intercohort Heterogeneity and Optimal Social Insurance Systems
9350	R.C. Douven and J.C. Engwerda	Is There Room for Convergence in the E.C.?
9351	F. Vella and M. Verbeek	Estimating and Interpreting Models with Endogenous Treatment Effects: The Relationship Between Competing Estimators of the Union Impact on Wages
9352	C. Meghir and G. Weber	Intertemporal Non-separability or Borrowing Restrictions? A Disaggregate Analysis Using the US CEX Panel
9353	V. Feltkamp	Alternative Axiomatic Characterizations of the Shapley and Banzhaf Values
9354	R.J. de Groof and M.A. van Tuijl	Aspects of Goods Market Integration. A Two-Country-Two -Sector Analysis
9355	Z. Yang	A Simplicial Algorithm for Computing Robust Stationary Points of a Continuous Function on the Unit Simplex
9356	E. van Damme and S. Hurkens	Commitment Robust Equilibria and Endogenous Timing
9357	W. Güth and B. Peleg	On Ring Formation In Auctions
9358	V. Bhaskar	Neutral Stability In Asymmetric Evolutionary Games
9359	F. Vella and M. Verbeek	Estimating and Testing Simultaneous Equation Panel Data Models with Censored Endogenous Variables
9360	W.B. van den Hout and J.P.C. Blanc	The Power-Series Algorithm Extended to the <i>BMAP/PH/1</i> Queue
9361	R. Heuts and J. de Klein	An (s,q) Inventory Model with Stochastic and Interrelated Lead Times
9362	K.-E. Wärneryd	A Closer Look at Economic Psychology
9363	P.J.-J. Herings	On the Connectedness of the Set of Constrained Equilibria
9364	P.J.-J. Herings	A Note on "Macroeconomic Policy in a Two-Party System as a Repeated Game"
9365	F. van der Ploeg and A. L. Bovenberg	Direct Crowding Out, Optimal Taxation and Pollution Abatement
9366	M. Pradhan	Sector Participation in Labour Supply Models: Preferences or Rationing?

No.	Author(s)	Title
9367	H.G. Bloemen and A. Kapteyn	The Estimation of Utility Consistent Labor Supply Models by Means of Simulated Scores
9368	M.R. Baye, D. Kovenock and C.G. de Vries	The Solution to the Tullock Rent-Seeking Game When $R > 2$: Mixed-Strategy Equilibria and Mean Dissipation Rates
9369	T. van de Klundert and S. Smulders	The Welfare Consequences of Different Regimes of Oligopolistic Competition in a Growing Economy with Firm-Specific Knowledge
9370	G. van der Laan and D. Talman	Intersection Theorems on the Simplotope
9371	S. Muto	Alternating-Move Preplays and $vN - M$ Stable Sets in Two Person Strategic Form Games
9372	S. Muto	Voters' Power in Indirect Voting Systems with Political Parties: the Square Root Effect
9373	S. Smulders and R. Gradus	Pollution Abatement and Long-term Growth
9374	C. Fernandez, J. Osiewalski and M.F.J. Steel	Marginal Equivalence in v -Spherical Models
9375	E. van Damme	Evolutionary Game Theory
9376	P.M. Kort	Pollution Control and the Dynamics of the Firm: the Effects of Market Based Instruments on Optimal Firm Investments
9377	A. L. Bovenberg and F. van der Ploeg	Optimal Taxation, Public Goods and Environmental Policy with Involuntary Unemployment
9378	F. Thuijsman, B. Peleg, M. Amitai & A. Shmida	Automata, Matching and Foraging Behavior of Bees
9379	A. Lejour and H. Verbon	Capital Mobility and Social Insurance in an Integrated Market
9380	C. Fernandez, J. Osiewalski and M. Steel	The Continuous Multivariate Location-Scale Model Revisited: A Tale of Robustness
9381	F. de Jong	Specification, Solution and Estimation of a Discrete Time Target Zone Model of EMS Exchange Rates
9401	J.P.C. Kleijnen and R.Y. Rubinstein	Monte Carlo Sampling and Variance Reduction Techniques
9402	F.C. Drost and B.J.M. Werker	Closing the Garch Gap: Continuous Time Garch Modeling
9403	A. Kapteyn	The Measurement of Household Cost Functions: Revealed Preference Versus Subjective Measures

No.	Author(s)	Title
9404	H.G. Bloemen	Job Search, Search Intensity and Labour Market Transitions: An Empirical Exercise
9405	P.W.J. De Bijl	Moral Hazard and Noisy Information Disclosure
9406	A. De Waegenaere	Redistribution of Risk Through Incomplete Markets with Trading Constraints
9407	A. van den Nouweland, P. Borm, W. van Golstein Brouwers, R. Groot Bruinderink, and S. Tijs	A Game Theoretic Approach to Problems in Telecommunication
9408	A.L. Bovenberg and F. van der Ploeg	Consequences of Environmental Tax Reform for Involuntary Unemployment and Welfare
9409	P. Smit	Arnoldi Type Methods for Eigenvalue Calculation: Theory and Experiments
9410	J. Eichberger and D. Kelsey	Non-additive Beliefs and Game Theory
9411	N. Dagan, R. Serrano and O. Volij	A Non-cooperative View of Consistent Bankruptcy Rules
9412	H. Bester and E. Petrakis	Coupons and Oligopolistic Price Discrimination
9413	G. Koop, J. Osiewalski and M.F.J. Steel	Bayesian Efficiency Analysis with a Flexible Form: The AIM Cost Function
9414	C. Kilby	World Bank-Borrower Relations and Project Supervision
9415	H. Bester	A Bargaining Model of Financial Intermediation
9416	J.J.G. Lemmen and S.C.W. Eijffinger	The Price Approach to Financial Integration: Decomposing European Money Market Interest Rate Differentials
9417	J. de la Horra and C. Fernandez	Sensitivity to Prior Independence via Farlie-Gumbel-Morgenstern Model

P.O. BOX 90153, 5000 LE TILBURG, THE NETHERLANDS

Bibliotheek K. U. Brabant

17 000 01176819 0