Tilburg University

Characterizing properties of approximate solutions for optimization problems

Norde, H.W.; Patrone, F.; Tijs, S.H.

Publication date:
1996

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Norde, H. W., Patrone, F., & Tijs, S. H. (1996). *Characterizing properties of approximate solutions for optimization problems*. (CentER Discussion Paper; Vol. 1996-101). Microeconomics.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM
8414 R
1996-101
8414
1996
NR.101

Center
for
Economic Research

Discussion paper

Tilburg University

Center
for
Economic Research

8414
1996
101

No. 96101

90

**CHARACTERIZING PROPERTIES OF
APPROXIMATE SOLUTIONS FOR
OPTIMIZATION PROBLEMS**

By Henk Norde, Fioravante Patrone and Stef Tijs

R 35

October 1996

t optimization

ISSN 0924-7815

Characterizing Properties of Approximate Solutions for Optimization Problems¹

Henk Norde

Department of Econometrics, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands.

Fioravante Patrone

Department of Mathematics, Via Dodecaneso 35, University of Genoa, 16146 Genoa, Italy.

Stef Tijs

Department of Econometrics and CentER, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands.

Abstract: Approximate solutions for optimization problems become of interest if the “true” optimum can not be found: this may happen for the simple reason that an optimum does not exist or because of the “bounded rationality” (or accuracy) of the optimizer. In this paper we characterize several approximate solutions by means of consistency and invariance properties. In particular, we prove that, besides the trivial ones, there are no consistent solutions satisfying non-emptiness, translation and multiplication invariance.

Key-words: Approximate optimization, consistency, invariance properties.

1 Introduction

In this paper we shall try to give an answer to an apparently silly question: is the concept of “approximate solution” in optimization meaningful? We will show that, moving from exact to approximate optimization, some serious problems may arise.

One reason to focus on approximate optimization can be derived from the increasing interest for the issues related with “bounded rationality” in game

¹The authors wish to thank CNR-Italy, the GNAFA group of CNR-Italy and Tilburg University for financial support and A. Rustichini, P. Wakker and T. Zolezzi for helpful suggestions for this paper.

theory: it turns out that more and more the emphasis is shifted from maximization to approximate maximization. On this point, we only refer to Fudenberg and Tirole (1991), Myerson (1991) and Radner (1980). The latter paper is interesting both, to understand the kind of results that can be achieved along this path and for the remarks about the problems that arise when approximate maximization enters the scene.

The interest for approximate optimization arising from game theory is only one particular case of a general issue. In many cases, given an optimization problem, one does not look for the maximum: this can happen for the obvious reason that a maximum does not exist or for the difficulty of finding it. In both cases, one should have some "rule" that says when the search for a maximum could stop.

Clearly, many different kinds of rules can be devised, from some "rule of thumb" to a sophisticated analysis that compares the computational costs for improving the degree of approximation and the benefits that result.

The approach that will be used in this paper is usually referred to as "axiomatic". That is, we shall state some desirable properties of an "approximate solution concept" and will analyse their consequences and mutual compatibility.

To be more specific, we shall investigate a special issue related with these rules: how they should be if one wants to behave in a consistent way across different approximation problems and, at the same time, one has to take into account some invariance properties.

The invariance requirement is due to the fact that, in many cases, the function to be maximized is only a representative of a class of equivalent functions (let us recall at least utility theory, and the fact that in hard sciences the origin or the scale of measurement quite often can be freely chosen). The remarkable result that we get is a kind of "impossibility theorem", which asserts that there are no consistent rules for choosing truly approximate solutions if one wants to take into account translation and multiplication invariance (as one should do, e.g., when dealing with expected utilities). We also investigate the cases in which one takes into account separately these invariance requirements.

Special emphasis is given to rules that take into account a reasonable monotonicity condition, that can be considered as an instance of the IIA (independence of irrelevant alternatives) principle: the main reason to consider this point of view is that we try to consider classes of optimization problems which contain both bounded and unbounded functions. This interest is an outgrowth of previous research done by the authors in the context of

semi-infinite bimatrix games (see Jurg and Tijs (1993), Lucchetti, Patrone and Tijs (1986) and Norde and Potters (1995)). We get, under appropriate assumptions, a class of approximate solutions to which belong the (ε, k) -solutions investigated in the papers quoted above. We also like to emphasize that some examples show that careless specification of the domain of the rule can give quite strange results.

The two previous paragraphs are an approximate description of the contents of sections 3 and 4 respectively, while section 2 is devoted to the setting of the problem. In section 6 we investigate rules, which make use of sequences. As it could be expected, reasonable conditions restrict the attention to maximizing sequences. So that, in some sense, the "impossibility result" is somehow circumvented: results that guarantee the existence of ε -Nash equilibria for every $\varepsilon > 0$ do have a meaning in this setting (Tijs (1981)); the same can be said, in optimization, about Tykhonov well-posedness (Dontchev-Zolezzi (1993)); see Patrone (1987) for remarks about the invariance of this property). We want to add, however, some warning about sequences. First, for practical reasons one has eventually to give one solution: so, even if sequences can be considered interesting for theoretical reasons, they do not solve the problem of finding an approximate solution for an optimization problem. Secondly, sometimes one is interested in solving (approximately) problems which are an approximate version of the "true" one: in this case, tricks like those of considering asymptotically minimizing sequences (in the terminology of Dontchev-Zolezzi), fall outside the scope of our investigation. Actually, we have completely skipped any reference to "continuity" properties of our solution maps. In our opinion, this is a topic that deserves to be thoroughly studied.

Notation Throughout this paper we denote the set $\mathbb{R} \cup \{+\infty\}$ by $\mathbb{R}^*$ and the set $\mathbb{R} \cup \{-\infty, +\infty\}$ by $\overline{\mathbb{R}}$.

2 Optimization problems

An *optimization problem* is a pair (A, u) where A is a non-empty set of alternatives and u is a real-valued function with domain A . Let $\mathcal{P}$ be a non-empty collection of optimization problems. A *solution* β on $\mathcal{P}$ is a map which assigns to every optimization problem $(A, u) \in \mathcal{P}$ a subset of A .

Example 2.1 For the following examples no special restriction is imposed upon $\mathcal{P}$.

a) The solution β_{tot} is defined by

$$\beta_{\text{tot}}(A, u) := A.$$

b) The solution β_{max} is defined by

$$\beta_{\text{max}}(A, u) := \{a \in A : u(a) \geq u(a') \text{ for every } a' \in A\}.$$

c) For $\varepsilon > 0$ the solution β_ε is defined by

$$\beta_\varepsilon(A, u) := \{a \in A : u(a) \geq u(a') - \varepsilon \text{ for every } a' \in A\}.$$

d) For $k \in \mathbb{R}$ the solution β^k is defined by

$$\beta^k(A, u) := \{a \in A : u(a) \geq k\}.$$

e) For $\varepsilon > 0, k \in \mathbb{R}$ the solution $\beta_{\varepsilon, k}$ is defined by

$$\beta_{\varepsilon, k}(A, u) := \begin{cases} \beta_{\text{max}}(A, u) & \text{if } \beta_{\text{max}}(A, u) \neq \emptyset \\ \beta_\varepsilon(A, u) & \text{if } \beta_{\text{max}}(A, u) = \emptyset \text{ and } \beta_\varepsilon(A, u) \neq \emptyset \\ \beta^k(A, u) & \text{otherwise} \end{cases}.$$

Notice that $\beta_{\text{max}}(A, u)$, $\beta_\varepsilon(A, u)$ and $\beta^k(A, u)$ can be empty; on the contrary, $\beta_{\text{tot}}(A, u) \neq \emptyset$ and $\beta_{\varepsilon, k}(A, u) \neq \emptyset$ for every $(A, u) \in \mathcal{P}$.

Two optimization problems (A, u) and (B, v) are *sup-equivalent* if

$$\sup_{x \in A} u(x) = \sup_{x \in B} v(x).$$

A solution β on $\mathcal{P}$ is *approximation consistent* if for every pair of sup-equivalent problems $(A, u), (B, v) \in \mathcal{P}$ the following statement is true:

if $b \in \beta(B, v)$ and $a \in A$ is such that $u(a) \geq v(b)$ then $a \in \beta(A, u)$.

So, if a solution β is approximation consistent, selection by β of an alternative $b \in B$ in some problem $(B, v) \in \mathcal{P}$, induces selection by β of all 'non-worse' alternatives in sup-equivalent problems. Clearly, the solutions a)-d) in example 2.1 are approximation consistent. For approximation consistent solutions we have the following proposition.

Proposition 2.1 *Let β be an approximation consistent solution on $\mathcal{P}$ and let $(A, u), (B, v) \in \mathcal{P}$ be such that $u(A) = v(B)$. Then there is a subset T of $u(A) (= v(B))$ such that $\beta(A, u) = u^{-1}(T)$ and $\beta(B, v) = v^{-1}(T)$.*

Proof Take $T := u(\beta(A, u))$.

If $a \in u^{-1}(T)$ then $u(a) = u(a')$ for some $a' \in \beta(A, u)$ and hence, by approximation consistency, $a \in \beta(A, u)$. So, $u^{-1}(T) \subseteq \beta(A, u)$. The inclusion $\beta(A, u) \subseteq u^{-1}(T)$ is obvious.

If $b \in v^{-1}(T)$ then $v(b) = u(a')$ for some $a' \in \beta(A, u)$. Since (A, u) and (B, v) are sup-equivalent we get, by approximation consistency, $b \in \beta(B, v)$. So, $v^{-1}(T) \subseteq \beta(B, v)$. If $b \in \beta(B, v)$ then, since $u(A) = v(B)$, there is an $a \in A$ such that $u(a) = v(b)$ and hence, by approximation consistency, $a \in \beta(A, u)$. Therefore, $v(b) = u(a) \in T$ and hence $b \in v^{-1}(T)$. So, $\beta(B, v) \subseteq v^{-1}(T)$. ■

The above proposition shows that, if β is approximation consistent, the set $\beta(A, u)$ only depends on the range $u(A)$ of u . So, if we are interested in approximation consistent solutions only, we may identify an optimization problem (A, u) with $u(A)$, the range of u , which is a subset of $\mathbb{R}$. In the next sections we focus on this approach.

3 Axioms and examples

Let $\mathcal{S}$ be a non-empty collection of non-empty subsets of $\mathbb{R}$. A solution σ on $\mathcal{S}$ is a map which assigns to every $S \in \mathcal{S}$ a subset $\sigma(S)$ of S .

A solution σ on $\mathcal{S}$ satisfies (AC) (*approximation consistency*) if for every $S_1, S_2 \in \mathcal{S}$ with $\sup S_1 = \sup S_2$ the following statement is true:

if $s_2 \in \sigma(S_2)$ and $s_1 \in S_1$ is such that $s_1 \geq s_2$ then $s_1 \in \sigma(S_1)$.

The reason for introducing (AC) is given by proposition 2.1: it is immediate to see that, if a solution β on $\mathcal{P}$ is approximation consistent, as defined in the previous section, then the induced solution σ on the family $\mathcal{S}$ of ranges $u(A)$ ($(A, u) \in \mathcal{P}$), satisfies (AC). Conversely, a solution σ on $\mathcal{S}$, satisfying (AC), induces, for every $\mathcal{P}$ with ranges in $\mathcal{S}$, an approximation consistent solution β .

The (AC) condition is not entirely satisfactory: one expects that $\sigma(S)$ can be described as $\{s \in S : s \geq \gamma\}$ or $\{s \in S : s > \gamma\}$ for some γ (depending on S). However, we can have strict or weak inequality, depending on the value of $\sup S$, as can be seen in the next example.

Example 3.1 Let $\mathcal{S}$ be the collection of all non-empty subsets of $\mathbb{R}$. The solution σ_{mix} on $\mathcal{S}$, defined by

$$\sigma_{\text{mix}}(S) := \begin{cases} \{s \in S : s \geq \sup S - 1\} & \text{if } \sup S \leq 0 \\ \{s \in S : s > \sup S - 1\} & \text{if } \sup S \in (0, +\infty) \\ \{s \in S : s \geq 22\} & \text{if } \sup S = +\infty \end{cases},$$

satisfies (AC).

In order to get rid of these kind of approximate solutions one would like to add the requirement that $\sigma(S)$ is a closed subset of S . However, the next example shows us that this addition does not “force” the parentheses to be closed.

Example 3.2 Consider $\mathcal{S} = \{\{0, 1\}\} \cup \{[\alpha, 1] : \alpha \in (0, 1)\}$ and let σ be the solution on $\mathcal{S}$, defined by $\sigma(\{0, 1\}) := \{1\}$ and $\sigma([\alpha, 1]) := [\alpha, 1]$ for every $\alpha \in (0, 1)$. The solution σ satisfies (AC) and $\sigma(S)$ is a closed subset of S for every $S \in \mathcal{S}$, but there is no $\gamma \in \mathbb{R}$ such that $\sigma(S) = \{s \in S : s \geq \gamma\}$ for every $S \in \mathcal{S}$.

So, we shall introduce the following axiom, an appropriate strenghtening of (AC).

A solution σ on $\mathcal{S}$ satisfies (SAC) (*strong approximation consistency*) if for every $S, S_1, S_2, \dots \in \mathcal{S}$ with $\sup S = \sup S_i$ for every $i \in \mathbb{N}$ the following statement is true:

if $s_i \in \sigma(S_i)$ for every $i \in \mathbb{N}$ and $s \in S$ is such that $s \geq \liminf_{i \rightarrow \infty} s_i$
then $s \in \sigma(S)$.

One easily verifies that (SAC) induces (AC). Moreover, (SAC) implies that $\sigma(S)$ is a closed subset of S for every $S \in \mathcal{S}$. In fact, if $\mathcal{S}$ is a collection of intervals, then σ satisfies (SAC) if and only if σ satisfies (AC) and $\sigma(S)$ is a closed subset of S for every $S \in \mathcal{S}$.

In the sequel we also make use of the following axioms.

A solution σ on $\mathcal{S}$ satisfies (NEM) (*non-emptiness*) if for every $S \in \mathcal{S}$ we have

$$\sigma(S) \neq \emptyset.$$

The collection $\mathcal{S}$ is *closed under translation* (CL+) if for every $S \in \mathcal{S}$ and $t \in \mathbb{R}$ we have $t + S := \{t + s : s \in S\} \in \mathcal{S}$. A solution σ on $\mathcal{S}$, obeying (CL+), satisfies (TI) (*translation invariance*) if for every $S \in \mathcal{S}$ and $t \in \mathbb{R}$ we have

$$\sigma(t + S) = t + \sigma(S).$$

The collection $\mathcal{S}$ is *closed under multiplication* (CL*) if for every $S \in \mathcal{S}$ and $\lambda > 0$ we have $\lambda S := \{\lambda s : s \in S\} \in \mathcal{S}$. A solution σ on $\mathcal{S}$, obeying (CL*), satisfies (MI) (*multiplication invariance*) if for every $S \in \mathcal{S}$ and $\lambda > 0$ we have

$$\sigma(\lambda S) = \lambda \sigma(S).$$

A solution σ on $\mathcal{S}$ satisfies (IIA) (*independence of irrelevant alternatives*) if for every $S, T \in \mathcal{S}$ with $S \subseteq T$ one has

$$\sigma(T) \cap S \subseteq \sigma(S).$$

So, if σ satisfies (IIA), selection by σ of an element $s \in T$, implies selection by σ of s in any subset S of T with $s \in S$. This notion of (IIA) is weaker than the one used in Kaneko (1980) and Peters (1992).

Example 3.3 For the following examples suppose that $\mathcal{S}$ is the collection of all non-empty subsets of $\mathbb{R}$.

- a) The solution σ_{mix} , defined in example 3.1, satisfies (AC) and (NEM).
- b) The solution σ_{rat} , defined by

$$\sigma_{\text{rat}}(S) := \{s \in S : s \text{ is rational}\},$$

only satisfies (IIA).

- c) The solution σ_{tot} , defined by

$$\sigma_{\text{tot}}(S) := S,$$

satisfies (SAC), (NEM), (TI), (MI) and (IIA).

- d) The solution σ_{max} , defined by

$$\sigma_{\text{max}}(S) := \{s \in S : s \geq s' \text{ for every } s' \in S\},$$

satisfies (SAC), (TI), (MI) and (IIA).

- e) The solution σ_ε (where $\varepsilon > 0$), defined by

$$\sigma_\varepsilon(S) := \{s \in S : s \geq \sup S - \varepsilon\},$$

satisfies (SAC), (TI) and (IIA).

f) The solution σ^k (where $k \in \mathbb{R}$), defined by

$$\sigma^k(S) := \{s \in S : s \geq k\},$$

satisfies (SAC) and (IIA).

g) The solution $\sigma_{\varepsilon,k}$ (where $\varepsilon > 0, k \in \mathbb{R}$), defined by

$$\sigma_{\varepsilon,k}(S) := \begin{cases} \sigma_{\max}(S) & \text{if } \sigma_{\max}(S) \neq \emptyset \\ \sigma_{\varepsilon}(S) & \text{if } \sigma_{\max}(S) = \emptyset \text{ and } \sigma_{\varepsilon}(S) \neq \emptyset \\ \sigma^k(S) & \text{otherwise} \end{cases},$$

only satisfies (NEM).

h) The solution $\hat{\sigma}_{\varepsilon,k}$ (where $\varepsilon > 0, k \in \mathbb{R}$), defined by

$$\hat{\sigma}_{\varepsilon,k}(S) := \begin{cases} \sigma_{\varepsilon}(S) & \text{if } \sup S \leq k + \varepsilon \\ \sigma^k(S) & \text{if } \sup S > k + \varepsilon \end{cases},$$

satisfies (SAC), (NEM) and (IIA). Notice that $\hat{\sigma}_{\varepsilon,k}(S) = \sigma_{\varepsilon}(S) \cup \sigma^k(S)$.

i) The solution $\sigma_{\text{pro}(\alpha,\beta)}(S)$ (where $\alpha > 1, \beta < 1$), defined by

$$\sigma_{\text{pro}(\alpha,\beta)}(S) := \begin{cases} \sigma^{\alpha s}(S) & \text{if } \sup(S) = s \in (-\infty, 0) \\ S & \text{if } \sup(S) = 0 \\ \sigma^{\beta s}(S) & \text{if } \sup(S) = s \in (0, +\infty) \\ S & \text{if } \sup(S) = +\infty \end{cases},$$

satisfies (SAC), (NEM) and (MI).

The following table summarizes the statements above.

(AC) (SAC) (NEM) (TI) (MI) (IIA)

σ_{mix}	+	-	+	-	-	-
σ_{rat}	-	-	-	-	-	+
σ_{tot}	+	+	+	+	+	+
σ_{max}	+	+	-	+	+	+
σ_{ε}	+	+	-	+	-	+
σ^k	+	+	-	-	-	+
$\sigma_{\varepsilon,k}$	-	-	+	-	-	-
$\hat{\sigma}_{\varepsilon,k}$	+	+	+	-	-	+
$\sigma_{\text{pro}(\alpha,\beta)}$	+	+	+	-	+	-

4 Characterizations for translation and multiplication invariant solutions

Let $\mathcal{S}$ be a collection of non-empty subsets of $\mathbb{R}$. We write $\mathcal{S} = \cup_{k \in \mathbb{R}^*} \mathcal{S}_k$ where $\mathcal{S}_k := \{S \in \mathcal{S} : \sup S = k\}$ for every $k \in \mathbb{R}^*$. For a function $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ we define the solution σ_a on $\mathcal{S}$ by

$$\sigma_a(S) := \{s \in S : s \geq a(\sup(S))\}.$$

So, σ_a selects, for every $S \in \mathcal{S}_k$, the elements $s \in S$ with $s \geq a(k)$. Clearly, σ_a satisfies (SAC). The following proposition shows that the solutions σ_a are completely characterized by (SAC). Note that the solutions, defined in example 3.3 c)-f),h),i), satisfy (SAC) and are, in fact, σ_a for some suitably chosen a .

Proposition 4.1 *Let $\mathcal{S}$ be a collection of non-empty subsets of $\mathbb{R}$ and let σ be a solution on $\mathcal{S}$. The solution σ satisfies (SAC) if and only if $\sigma = \sigma_a$ for some function a .*

Proof We only prove the only-if-part. So, assume that σ satisfies (SAC). First we define the function $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$. Let $k \in \mathbb{R}^*$. Define

$$a(k) := \begin{cases} \text{arbitrarily} & \text{if } S_k = \emptyset \\ \inf(\cup_{S \in S_k} \sigma(S)) & \text{otherwise} \end{cases} \quad (1)$$

(with the convention that $\inf \emptyset = +\infty$). Now we have to prove that $\sigma = \sigma_a$, i.e. we have to prove that $\sigma(S) = \sigma_a(S)$ for every $S \in \mathcal{S}$. So, let $S \in \mathcal{S}$ and let $k := \sup S$ (which trivially induces that $S_k \neq \emptyset$). For every $s \in \sigma(S)$ we have $s \geq a(k)$ by definition of $a(k)$. Therefore $\sigma(S) \subseteq \sigma_a(S)$. Note that the converse inclusion $\sigma_a(S) \subseteq \sigma(S)$ is trivial when $\sigma_a(S) = \emptyset$. So, assume $\sigma_a(S) \neq \emptyset$ and let $s \in \sigma_a(S)$. Then $s \geq a(k)$ which implies $a(k) \neq +\infty$. Therefore, $\cup_{S \in S_k} \sigma(S) \neq \emptyset$, and hence, by definition of $a(k)$, there is a sequence $S_1, S_2, \dots \in S_k$ and, for every $i \in \mathbb{N}$, an $s_i \in \sigma(S_i)$ such that $a(k) = \lim_{i \rightarrow \infty} s_i$. By (SAC) we get $s \in \sigma(S)$. Therefore $\sigma_a(S) \subseteq \sigma(S)$, which finishes the proof. ■

If we impose some feasibility condition upon the function a we get solutions which are characterized by (SAC) and (NEM).

Proposition 4.2 *Let S be a collection of non-empty subsets of $\mathbb{R}$ and let σ be a solution on S . The solution σ satisfies (SAC) and (NEM) if and only if $\sigma = \sigma_a$ for some function $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ satisfying $a(k) \leq k$ for every $k \in \mathbb{R}^*$, with strict inequality for every $k \in \mathbb{R}^*$ for which there is an $S \in S_k$ with $k \notin S$.*

Proof Let $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ be such that $a(k) \leq k$ for every $k \in \mathbb{R}^*$, with strict inequality for every $k \in \mathbb{R}^*$ for which there is an $S \in S_k$ with $k \notin S$ and let $\sigma = \sigma_a$. Clearly, σ_a satisfies (SAC). Let $S \in \mathcal{S}$ and let $k := \sup S$. If $k \in S$ then $k \in \sigma_a(S)$. If $k \notin S$ (which is, e.g., the case if $k = +\infty$) then $a(k) < k$ and we may choose an $s \in S$ with $s \geq a(k)$ and, as a consequence, $s \in \sigma_a(S)$. So, σ_a also satisfies (NEM).

In order to prove the only-if-part assume that σ satisfies (SAC) and (NEM). By (SAC) and the proof of proposition (4.1) we know that $\sigma = \sigma_a$, where a is defined by (1). Now we have to prove that a has the desired properties. So let $k \in \mathbb{R}^*$. If $S_k = \emptyset$ we choose $a(k) \leq k$. If $S_k \neq \emptyset$ there is an $S \in S_k$ and we may, by (NEM), take some element $s \in \sigma(S) = \sigma_a(S)$. We get $a(k) \leq s \leq k$, where the last inequality is strict if $k \notin S$. ■

The next theorem describes the class of solutions, which are characterized by

(SAC), (NEM) and (TI). It turns out that these solutions coincide with the collection of ' ε -optimal' solutions for some $\varepsilon \in [0, +\infty]$, if $\mathcal{S}$ only contains upper bounded sets.

Proposition 4.3 *Let $\mathcal{S}$ be a collection of non-empty subsets of $\mathbb{R}$ which satisfies (CL+) and let σ be a solution on $\mathcal{S}$. The solution σ satisfies (SAC), (NEM) and (TI) if and only if there is an $\varepsilon \geq 0$ ($\varepsilon > 0$ in case there is an upper bounded $S \in \mathcal{S}$ which has no maximum) such that $\sigma = \sigma_a$, where $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ is defined by*

$$\begin{cases} a(k) := k - \varepsilon & \text{for every } k \in \mathbb{R} \\ a(+\infty) = -\infty \end{cases} \quad (2)$$

Proof Clearly, $\sigma = \sigma_a$ satisfies (SAC), (NEM) and (TI) if a is defined by (2). In order to prove the only-if-part suppose that σ satisfies (SAC), (NEM) and (TI). Again, by (SAC) and the proof of proposition 4.1, we know that $\sigma = \sigma_a$, where a is defined by (1). Notice that, by (CL+), we have for every $k \in \mathbb{R}$

$$\mathcal{S}_k = \{k + S : S \in \mathcal{S}_0\}.$$

So, if $\mathcal{S}_k = \emptyset$ for some $k \in \mathbb{R}$, then $\mathcal{S}_k = \emptyset$ for every $k \in \mathbb{R}$ and we choose $a(k) = k - 37$ for every $k \in \mathbb{R}$. If $\mathcal{S}_k \neq \emptyset$ for some $k \in \mathbb{R}$, then $\mathcal{S}_k \neq \emptyset$ for every $k \in \mathbb{R}$. Moreover, we get by (TI),

$$\begin{aligned} a(k) &= \inf(\cup_{S \in \mathcal{S}_k} \sigma(S)) &= \inf(\cup_{S \in \mathcal{S}_0} \sigma(k + S)) &= \\ &= \inf(\cup_{S \in \mathcal{S}_0} (k + \sigma(S))) &= k + \inf(\cup_{S \in \mathcal{S}_0} \sigma(S)) &= \\ &= k + a(0). \end{aligned}$$

By (NEM) we get $a(0) \leq 0$. Moreover, if there is an upper bounded $S \in \mathcal{S}$ which has no maximum, then, by (CL+), there is an $S \in \mathcal{S}_0$ which has no maximum and we have $a(0) < 0$. So, take $\varepsilon = -a(0)$.

If $\mathcal{S}_{+\infty} = \emptyset$ we choose $a(+\infty) = -\infty$. Otherwise there is an $S \in \mathcal{S}_{+\infty}$ and, by (NEM), an $s \in \sigma(S)$. Since, by (CL+), $t + S \in \mathcal{S}_{+\infty}$ and, by (TI), $t + s \in \sigma(t + S)$ for every $t \in \mathbb{R}$ we get $a(+\infty) \leq t + s$ for every $t \in \mathbb{R}$. Hence, $a(+\infty) = -\infty$. ■

The solution $\tilde{\sigma}_{\varepsilon, k}$ of example 3.3 satisfies (SAC) and (NEM) but not (TI), the solution σ_ε satisfies (SAC) and (TI) but not (NEM) and one easily constructs a solution satisfying (NEM) and (TI) but not (SAC) (simply by defining the solution for $\mathcal{S}$ with $\sup S \in \{0, +\infty\}$ in an arbitrary but not approximation consistent way and extending this solution by translation

invariance). Therefore the axioms (SAC), (NEM) and (TI) are logically independent.

In the following proposition we characterize the 'proportional' solutions by (SAC), (NEM) and (MI).

Proposition 4.4 *Let S be a collection of non-empty subsets of $\mathbb{R}$ which satisfies (CL*) and let σ be a solution on S . The solution σ satisfies (SAC), (NEM) and (MI) if and only if there are $\alpha \geq 1$ and $\beta \leq 1$ such that $\sigma = \sigma_a$, where $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ is defined by*

$$\begin{cases} a(k) := \alpha k & \text{for every } k < 0 \\ a(0) \in \{-\infty, 0\} \\ a(k) := \beta k & \text{for every } k \in (0, \infty) \\ a(+\infty) \in \{-\infty, 0\} \end{cases} \quad (3)$$

(where $\alpha > 1$ if there is a $k < 0$ and an $S \in S_k$ with $k \notin S$, $\beta < 1$ if there is a $k > 0$ and an $S \in S_k$ with $k \notin S$ and $a(0) = -\infty$ if there is an $S \in S_0$ with $0 \notin S$).

Proof Clearly, $\sigma = \sigma_a$ satisfies (SAC), (NEM) and (MI) if a is defined by (3). In order to prove the only-if-part suppose that σ satisfies (SAC), (NEM) and (MI). Again, by (SAC) and the proof of proposition 4.1, we know that $\sigma = \sigma_a$, where a is defined by (1). Notice that, by (CL*), we have for every $k \in (0, +\infty)$

$$S_k = \{kS : S \in S_1\}.$$

So, if $S_k = \emptyset$ for some $k \in (0, +\infty)$, then $S_k = \emptyset$ for every $k \in (0, +\infty)$ and we choose $a(k) = \frac{1}{37}k$ for every $k \in \mathbb{R}$. If $S_k \neq \emptyset$ for some $k \in (0, +\infty)$, then $S_k \neq \emptyset$ for every $k \in (0, +\infty)$. Moreover, we get by (MI),

$$\begin{aligned} a(k) &= \inf(\cup_{S \in S_k} \sigma(S)) = \inf(\cup_{S \in S_1} \sigma(kS)) = \\ &= \inf(\cup_{S \in S_1} k\sigma(S)) = k \inf(\cup_{S \in S_1} \sigma(S)) = \\ &= ka(1). \end{aligned}$$

By (NEM) we get $a(1) \leq 1$. Moreover, if there is a $k \in (0, +\infty)$ and an $S \in S_k$ with $k \notin S$, then, by (CL*), there is an $S \in S_1$ with $1 \notin S$ and we have $a(1) < 1$. So, take $\beta = a(1)$.

In an analogous way we prove, in case $S_k \neq \emptyset$ for every $k \in (-\infty, 0)$, that $a(k) = k(-a(-1))$, for every $k \in (-\infty, 0)$, where $a(-1) \leq -1$ and $a(-1) < -1$ if there is an $S \in S_{-1}$ with $-1 \notin S$ (or equivalently, there is a

$k < 0$ and an $S \in \mathcal{S}_k$ with $k \notin S$).

Suppose $\mathcal{S}_0 \neq \emptyset$ and $a(0) < 0$. Then there is an $S \in \mathcal{S}$ and an $s \in \sigma(S)$ with $s < 0$. Since, by (CL*), $\lambda S \in \mathcal{S}_0$ and, by (MI), $\lambda s \in \sigma(\lambda S)$ for every $\lambda > 0$ we get $a(0) \leq \lambda s$ for every $\lambda > 0$. Hence $a(0) = -\infty$. In an analogous way one proves that $a(+\infty) \in \{-\infty, 0\}$. ■

The solution $\tilde{\sigma}_{\epsilon,k}$ of example 3.3 satisfies (SAC) and (NEM) but not (MI), the solution $\sigma_{\max}$ satisfies (SAC) and (MI) but not (NEM) and one easily constructs a solution satisfying (NEM) and (MI) but not (SAC) (simply by defining the solution for S with $\sup S \in \{-1, 0, 1, +\infty\}$ in an arbitrary but not approximation consistent way and extending this solution by multiplication invariance). Therefore the axioms (SAC), (NEM) and (MI) are logically independent.

Clearly, the trivial solution σ_{tot} satisfies (SAC), (NEM), (TI) and (MI). Moreover, if the collection $\mathcal{S}$ is such that every upper bounded $S \in \mathcal{S}$ has a maximum, then the solution, selecting the maximum for every upper bounded $S \in \mathcal{S}$ and selecting S for every $S \in \mathcal{S}$ which is not upper bounded, does either. In the following proposition we show the impossibility of finding another solution, satisfying these four properties.

Proposition 4.5 *Let $\mathcal{S}$ be a collection of non-empty subsets of $\mathbb{R}$, which satisfies (CL+) and (CL*). Suppose, moreover, that there is at least one upper bounded $S \in \mathcal{S}$ which has no maximum. Let σ be a solution on $\mathcal{S}$. The solution σ satisfies (SAC), (NEM), (TI) and (MI) if and only if $\sigma = \sigma_a$, where $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ is defined by $a(k) := -\infty$ for every $k \in \mathbb{R}^*$ (i.e. $\sigma_a = \sigma_{\text{tot}}$).*

Proof Again we only prove the only-if-part. Suppose σ satisfies (SAC), (NEM), (TI) and (MI). By (SAC) we may conclude that $\sigma = \sigma_a$ where $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ is defined by (1). Let $S \in \mathcal{S}$ be an upper bounded set without maximum and let $k := \sup S$. Since σ satisfies (NEM) there is an $s \in \sigma(S)$. Clearly $s < k$. For every $n \in \mathbb{N}$ we have, by (CL+) and (TI),

$$-(1 - n^{-1})k + S \in \mathcal{S}$$

and

$$-(1 - n^{-1})k + s \in \sigma(-(1 - n^{-1})k + S).$$

Moreover, by (CL*) and (MI), we get

$$-(n - 1)k + nS = n(-(1 - n^{-1})k + S) \in \mathcal{S}$$

and

$$-(n-1)k + ns = n(-(1-n^{-1})k + s) \in \sigma(-(n-1)k + nS).$$

Since $\sup(-(n-1)k + nS) = k$ and $\lim_{n \rightarrow \infty}(-(n-1)k + ns) = -\infty$ we get $a(k) = -\infty$. By (TI) we infer that $a(l) = -\infty$ for every $l \in \mathbb{R}$. We know already, from proposition 4.3, that $a(+\infty) = -\infty$. ■

Let us notice that, in the context of decision making under risk, u and v are von Neumann-Morgenstern utility functions representing the same preferences iff $v = cu + d$, with $c > 0$ and $d \in \mathbb{R}$. So, if one wants to stress the point of view that only preferences have a true meaning, one should use a “solution rule” for optimization problems that takes into account this fact. But proposition 4.5 just shows that it is impossible to do this in a non-trivial way. Otherwise stated: for von Neumann-Morgenstern preferences there is no sensible concept of approximate optimum! If one wants to talk in a meaningful way of approximate optimization, an escape route could be the addition of further details that allow for some “absolute” reference point (e.g.: how do we decide whether the oscillations of last week at the New York Stock Exchange were wild or not? Maybe we refer to the previous history as a benchmark). The interesting question is whether it can be done in a consistent way, without resorting to an “absolute” utility function.

5 Characterizations under IIA

Proposition 4.3 provides a nice characterization of solutions σ on collections of *upper bounded* subsets of $\mathbb{R}$: either σ selects the maximum of $S \in \mathcal{S}$ (if all upper bounded $S \in \mathcal{S}$ have a maximum) or there is an $\varepsilon \in (0, +\infty]$ such that σ selects all ‘ ε -optimal’ elements. In order to get a nice characterization, which takes also the *unbounded* subsets of $\mathcal{S}$ into account, we have to replace (TI) by (IIA). Since (IIA) deals with inclusions, an appropriate condition upon $\mathcal{S}$ would be either to be *closed under taking subsets* (CLC) or to be *closed under taking supersets* (CL $\supset$). The following example shows that a nice characterization with (IIA), is not possible if $\mathcal{S}$ satisfies (CLC).

Example 5.1 Consider the class $\mathcal{S}$ of all non-empty and upper bounded subsets S which satisfy the condition that there exists a $t \in [0, 1)$ such that $S \subseteq t + \mathbb{Z}$. Define the solution σ_a by

$$a(k) := \begin{cases} k - 22 & \text{if } k \in \mathbb{Z} \\ k - 37 & \text{otherwise} \end{cases}.$$

Clearly σ_a satisfies (SAC) and (NEM). It also satisfies (IIA): this is due to the fact that for $S, T \in \mathcal{S}$ with $S \subseteq T$ both are contained in the same $t + \mathbb{Z}$ for some $t \in [0, 1)$. In fact, one can prove that any σ_a , with a feasible function a which is non-decreasing on $t + \mathbb{Z}$ for every $t \in [0, 1)$, satisfies (SAC), (NEM) and (IIA).

The reason for strange examples as above lies in the fact that the collection $\mathcal{S}$ is too poor: $\mathcal{S}$ can be partitioned into several subcollections such that sets belonging to different subcollections are not related by inclusion. These problems do not occur if $\mathcal{S}$ satisfies (CL $\supset$).

Proposition 5.1 *Let $\mathcal{S}$ be a collection of non-empty subsets of $\mathbb{R}$ which satisfies (CL $\supset$) and let σ be a solution on $\mathcal{S}$. The solution σ satisfies (SAC), (NEM) and (IIA) if and only if $\sigma = \sigma_a$ for some non-decreasing function $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ satisfying $a(k) \leq k$ for every $k \in \mathbb{R}^*$, with strict inequality for every $k \in \mathbb{R}^*$ for which there is an $S \in \mathcal{S}_k$ with $k \notin S$.*

Proof Again we only prove the only-if-part. Suppose σ satisfies (SAC), (NEM) and (IIA). By (SAC) we may conclude that $\sigma = \sigma_a$, where a is defined by (1). The only thing left to prove is that a is non-decreasing. Choose $a(k) = -\infty$ if $\mathcal{S}_k = \emptyset$. Notice that, if $\mathcal{S}_k = \emptyset$ for some $k \in \mathbb{R}^*$, (CL $\supset$) implies that $\mathcal{S}_l = \emptyset$ for every $l < k$. So let $k, l \in \mathbb{R}^*$ with $k < l$ and suppose that $\mathcal{S}_k \neq \emptyset$ and $\mathcal{S}_l \neq \emptyset$. By (CL $\supset$) we get $S := (-\infty, k] \in \mathcal{S}_k$ and $T := (-\infty, l] \in \mathcal{S}_l$. Therefore $\sigma(S) = [a(k), k]$ and $\sigma(T) = [a(l), l]$. As a consequence we get, by (IIA),

$$[a(l), k] = \sigma(T) \cap S \subseteq \sigma(S) = [a(k), k].$$

Therefore $a(k) \leq a(l)$ which finishes the proof. ■

An example of a solution, satisfying the requirements above, is given by $\hat{\sigma}_{\varepsilon, k}$, described in h) of example 3.3. The proposition above is not completely satisfactory, due to the fact that (CL $\supset$) is a very strong requirement on the class $\mathcal{S}$. However, example 5.1 showed that it is not easy to get rid of it. Another approach is that one asks for some strengthening of (SAC), instead of looking for too special classes $\mathcal{S}$.

A solution σ on $\mathcal{S}$ satisfies (SMAC) (*strong monotonic approximation consistency*) if for every $S, S_1, S_2, \dots \in \mathcal{S}$ with $\sup S \leq \sup S_i$ for every $i \in \mathbb{N}$ the following statement is true:

if $s_i \in \sigma(S_i)$ for every $i \in \mathbb{N}$ and $s \in S$ is such that $s \geq \liminf_{i \rightarrow \infty} s_i$ then $s \in \sigma(S)$.

It is obvious from the definitions that (SMAC) implies (SAC). Conversely, if $S_\infty = \emptyset$, we have that (SAC) and (TI) imply (SMAC).

With respect to what has been said at the beginning of this section, about an unsatisfactory characterization of solutions for unbounded subsets, we want to point out the following: if we replace (SAC) and (TI) by (SMAC), which is a weaker condition on *bounded* subsets, it is possible to give a nice characterization which takes also *unbounded* sets into account.

Proposition 5.2 *Let S be a collection of non-empty subsets of $\mathbb{R}$ and let σ be a solution on S . The solution σ satisfies (SMAC) and (NEM) if and only if $\sigma = \sigma_a$ for some non-decreasing function $a : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ satisfying $a(k) \leq k$ for every $k \in \mathbb{R}^*$, with strict inequality for every $k \in \mathbb{R}^*$ for which there is an $S \in S_k$ with $k \notin S$.*

Proof First we prove the if-part: suppose $\sigma = \sigma_a$ for some a as mentioned above. By proposition 4.2 we infer that σ satisfies ((SAC) and) (NEM). Suppose $S, S_1, S_2, \dots \in S$ with $\sup S \leq \sup S_i$ for every $i \in \mathbb{N}$ and let $s \in S$, $s_i \in \sigma(S_i)$ for every $i \in \mathbb{N}$, be such that $s \geq \liminf_{i \rightarrow \infty} s_i$. Since $s_i \geq a(\sup S_i) \geq a(\sup S)$ for every $i \in \mathbb{N}$, we get $s \geq a(\sup S)$. Hence $s \in \sigma(S)$. So, σ satisfies (SMAC).

For the proof of the only-if-part suppose that σ satisfies (SMAC) and (NEM). Since (SMAC) induces (SAC) we get $\sigma = \sigma_a$, where a is defined by (1). Notice however, that a needs not be non-decreasing. Now define $\bar{a} : \mathbb{R}^* \rightarrow \overline{\mathbb{R}}$ by

$$\bar{a}(k) = \inf\{a(l) : l \text{ is such that } S_l \neq \emptyset \text{ and } l \geq k\}$$

(with the usual convention that $\inf \emptyset = +\infty$). Clearly, $\bar{a}$ is non-decreasing. We prove that $\sigma_a = \sigma_{\bar{a}}$. So, let $S \in S$ and $k := \sup S$ (which induces $S_k \neq \emptyset$). Since $\bar{a}(k) \leq a(k)$ we have $\sigma_a(S) \subseteq \sigma_{\bar{a}}(S)$. Now let $s \in \sigma_{\bar{a}}(S)$. By definition of $\bar{a}(k)$ there is a sequence $l_1, l_2, l_3, \dots$ with $S_{l_i} \neq \emptyset$ and $l_i \geq k$ for every $i \in \mathbb{N}$ and $\bar{a}(k) = \lim_{i \rightarrow \infty} a(l_i)$. For every $i \in \mathbb{N}$ there is, by definition of $a(l_i)$, an $S_i \in S_{l_i}$ and $s_i \in \sigma(S_i)$ such that $\lim_{i \rightarrow \infty} a(l_i) = \lim_{i \rightarrow \infty} s_i$. Since $s \geq \bar{a}(k) = \lim_{i \rightarrow \infty} a(l_i) = \lim_{i \rightarrow \infty} s_i$ we get by (SMAC): $s \in \sigma(S) = \sigma_a(S)$. So, $\sigma_{\bar{a}}(S) \subseteq \sigma_a(S)$. ■

6 Approximation with sequences

A remarkable result of section 4 was that there was no solution, besides the trivial σ_{tot} , satisfying (SAC), (NEM), (TI) and (MI), in case $\mathcal{S}$ contains at least one upper bounded set without maximum. In this section we will get a positive result by considering generalized solutions. In order to do so we need some definitions.

Let $\mathcal{S}$ be a non-empty collection of non-empty subsets of $\mathbb{R}$. A *generalized solution* Σ on $\mathcal{S}$ is a map which assigns to every $S \in \mathcal{S}$ a subset $\Sigma(S)$ of S_{inc}^N , the collection of non-decreasing sequences in S . A sequence $\underline{s} = s_1, s_2, \dots$ in S_{inc}^N can be interpreted as a sequence of approximate optimal elements, where the degree of approximation gets better when indices are increasing. For generalized solutions we use the following axioms.

A solution Σ on $\mathcal{S}$ satisfies (AC) (*approximation consistency*) if for every $S_1, S_2 \in \mathcal{S}$ with $\sup S_1 = \sup S_2$ the following statement is true:

if $\underline{s}_2 \in \Sigma(S_2)$ and $\underline{s}_1 \in S_1$ is such that $\lim \underline{s}_1 \geq \lim \underline{s}_2$ then $\underline{s}_1 \in \Sigma(S_1)$.

The definitions of (NEM), (TI) and (MI) for generalized solutions are obvious.

Example 6.1 The following two generalized solutions, defined on a collection $\mathcal{S}$ obeying (CL+) and (CL*), all satisfy (AC), (NEM), (TI) and (MI).

- a) Define Σ_{opt} by $\Sigma_{opt}(S) := \{\underline{s} \in S_{inc}^N : \lim \underline{s} = \sup S\}$.
- b) Define Σ_{tot} by $\Sigma_{tot}(S) := S_{inc}^N$.

The following proposition shows that all generalized solutions, satisfying (AC), (NEM), (TI) and (MI), are mixtures of the solutions in example 6.1.

Proposition 6.1 *Let $\mathcal{S}$ be a collection of non-empty subsets of $\mathbb{R}$ and let Σ be a generalized solution on $\mathcal{S}$. The solution Σ satisfies (AC), (NEM), (TI) and (MI) if and only if Σ coincides with one of the solutions Σ_{opt} or Σ_{tot} on the collection of upper bounded subsets in $\mathcal{S}$ and if Σ coincides with one of these two solutions (but not necessarily the same) on $\mathcal{S}_{+\infty}$.*

Proof The if-part of the proof is left to the reader. For the only-if-part assume that Σ satisfies (AC), (NEM), (TI) and (MI). We only prove that Σ coincides with one of the solutions Σ_{opt} or Σ_{tot} on the collection of upper bounded subsets. The proof for $\mathcal{S}_{+\infty}$ is similar. We distinguish between two

cases.

Case 1: $\Sigma(S) \subseteq \Sigma_{opt}(S)$ for every upper bounded $S \in \mathcal{S}$.

Let $S \in \mathcal{S}$ be upper bounded. By (NEM), there is a sequence $\underline{s} \in \Sigma(S)$. By (AC), comparison of every sequence $\underline{s}' \in \Sigma_{opt}(S)$ with $\underline{s}$, implies $\Sigma_{opt}(S) \subseteq \Sigma(S)$. Therefore, $\Sigma(S) = \Sigma_{opt}(S)$.

Case 2: there is an upper bounded $S' \in \mathcal{S}$ with $\Sigma(S') \not\subseteq \Sigma_{opt}(S')$.

Let $S' \in \mathcal{S}$ and $\underline{s}' \in \Sigma(S')$ be such that $\underline{s}' \notin \Sigma_{opt}(S')$, let $k' := \sup S'$ and $k'' := \lim \underline{s}'$. Since $\underline{s}' \notin \Sigma_{opt}(S')$ we have $k'' < k'$. Moreover, we have for every $n \in \mathbb{N}$ by (CL+) and (TI),

$$-(1 - n^{-1})k' + S' \in \mathcal{S}$$

and

$$-(1 - n^{-1})k' + \underline{s}' \in \Sigma(-(1 - n^{-1})k' + S').$$

Moreover, by (CL*) and (MI), we get

$$-(n - 1)k' + nS' = n(-(1 - n^{-1})k' + S') \in \mathcal{S}$$

and

$$-(n - 1)k' + n\underline{s}' = n(-(1 - n^{-1})k' + \underline{s}') \in \Sigma(-(n - 1)k' + nS').$$

Now let $\underline{s} \in \Sigma_{tot}(S')$ and let $l := \lim \underline{s}$. Choose $n \in \mathbb{N}$ such that $-(n - 1)k' + nk'' < l$. By (AC), comparison of the sequences $\underline{s}$ and $-(n - 1)k' + n\underline{s}'$ yields $\underline{s} \in \Sigma(S')$. Therefore, $\Sigma(S') = \Sigma_{tot}(S')$. By (TI) one infers that $\Sigma(S) = \Sigma_{tot}(S)$ for every upper bounded $S \in \mathcal{S}$. ■

References

- Dontchev A. and Zolezzi T. (1993), Well-posed Optimization Problems. Lecture Notes in Mathematics, **1543**, Springer, Berlin.
- Fudenberg D. and Tirole J. (1991), Game Theory. MIT Press, Cambridge (Massachusetts).
- Jurg P. and Tijs S. (1993), On the determinateness of semi-infinite bimatrix games. International Journal of Game Theory **21**: 361-369.
- Kaneko M. (1980), An extension of the Nash bargaining problem and the Nash social welfare function. Theory and Decision, **12**, 135-148.

- Lucchetti R., Patrone F., Tijs S.H. (1986), Determinateness of two-person games. *Bollettino U.M.I.* **6**: 907-924.
- Myerson R.B. (1991), *Game Theory: Analysis of Conflict*. Harvard University Press, Cambridge (Massachusetts).
- Norde H. and Potters J. (1995), On the determinateness of $m \times \infty$ -bimatrix games. Report 9515, Department of Mathematics, University of Nijmegen.
- Patrone F. (1987), Well-Posedness as an Ordinal Property. *Rivista di Matematica pura ed applicata* **1**: 95-104.
- Peters H. (1992), *Axiomatic bargaining game theory*. Kluwer Academic Publishers, Dordrecht.
- Radner R. (1980), Collusive behavior in non-cooperative epsilon equilibria of oligopolies with long but finite lives, *Journal of Economic Theory*, **22**: 121-157.
- Tijs S. (1981), Nash equilibria for noncooperative n-person games in normal form. *SIAM Review* **23**: 225-237.

No.	Author(s)	Title
9616	G. Asheim and M. Dufwenberg	Admissibility and Common Knowledge
9617	C. Fernández, J. Osiewalski and M. Steel	On the Use of Panel Data in Bayesian Stochastic Frontier Models
9618	H. Huizinga	Intrafirm Information Transfers and Wages
9619	B. Melenberg and B. Werker	On the Pricing of Options in Incomplete Markets
9620	F. Kleibergen	Reduced Rank Regression using Generalized Method of Moments Estimators
9621	F. Janssen, R. Heuts and T. de Kok	The Value of Information in an (R,s,Q) Inventory Model
9622	F. Groot, C. Withagen and A. de Zeeuw	Strong Time-Consistency in the Cartel-Versus-Fringe Model
9623	R. Euwals and A. van Soest	Desired and Actual Labour Supply of Unmarried Men and Women in the Netherlands
9624	M. Khanman, M. Perry and P.J. Reny	An Ex-Post Envy-Free and Efficient Allocation Mechanism: Imperfect Information Without Common Priors
9625	C. Eaves, G. van der Laan D. Talman and Z. Yang	Balanced Simplices on Polytopes
9626	H. Bloemen and A. Kalwij	Female Employment and Timing of Births Decisions: a Multiple State Transition Model
9627	P. Bolton	Strategic Experimentation: a Revision
9628	C. Fershtman and Y. Weiss	Social Rewards, Externalities and Stable Preferences
9629	P.M. Kort, G. Feichtinger R.F. Hartl and J.L. Haunschmied	Optimal Enforcement Policies (Crackdowns) on a Drug Market
9630	C. Fershtman and A. de Zeeuw	Tradeable Emission Permits in Oligopoly
9631	A. Cukierman	The Economics of Central Banking
9632	A. Cukierman	Targeting Monetary Aggregates and Inflation in Europe
9633	F. de Roon, C. Veld and J. Wei	A Study on the Efficiency of the Market for Dutch Long Term Call Options
9634	B. van Aarle, L. Bovenberg M. Raith	Money, Fiscal Deficits and Government Debt in a Monetary Union

No.	Author(s)	Title
9635	G. van der Laan, D. Talman and Z. Yang	Existence of and Equilibrium in a Competitive Economy with Indivisibilities and Money
9636	I. Brouwer, J. van der Put and C. Veld	Contrarian Investment Strategies in a European Context
9637	M. Berg, F. van der Duyn Schouten and J. Jansen	Optimal Service Policies to Remote Customers with Delay-Limits
9638	F.C. Drost and C.A.J. Klaassen	Efficient Estimation in Semiparametric GARCH Models
9639	A.G. de Kok and F.B.S.L.P. Janssen	Demand Management in Multi-Stage Distribution Chain
9640	D.A. Freund, T.J. Kniesner and A.T. LoSasso	How Managed Care Affects Medicaid Utilization A Synthetic Difference-in-Differences Zero-Inflated Count Model
9641	M.J. Lee	Instrumental Variable Estimation For Linear Panel Data Models
9642	W. Härdle, E. Mammen and M. Müller	Testing Parametric versus Semiparametric Modelling in Generalized Linear Models
9643	C. Dustmann, N. Rajah and A. van Soest	Part-Time Work, School Success and School Leaving
9644	S.C.W. Eijffinger and M. Hoeberichts	The Trade Off Between Central Bank Independence and Conservativeness
9645	R. Sarin and P. Wakker	A Single-Stage Approach to Anscombe and Aumann's Expected Utility
9646	J.P. Ziliak and T.J. Kniesner	The Importance of Sample Attrition in Life Cycle Labor Supply Estimation
9647	P.M. Kort	Optimal R&D Investments of the Firm
9648	M.P. Berg	Performance Comparisons for Maintained Items
9649	H. Uhlig and Y. Xu	Effort and the Cycle: Cyclical Implications of Efficiency Wages
9650	M. Slikker and A. van den Nouweland	Communication Situations with a Hierarchical Player Partition
9651	H.L.F. de Groot	The Struggle for Rents in a Schumpeterian Economy
9652	R.M. de Jong and J. Davidson	Consistency of Kernel Estimators of heteroscedastic and Autocorrelated Covariance Matrices

No.	Author(s)	Title
9653	J. Suijs, A. De Waegenare and P. Borm	Stochastic Cooperative Games in Insurance and Reinsurance
9654	A.N. Banerjee and J.R. Magnus	Testing the Sensitivity of OLS when the Variance Matrix is (Partially) Unknown
9655	A. Kalwij	Estimating the Economic Return to Schooling on the basis of Panel Data
9656	M. Lind and F. van Megen	Order Based Cost Allocation Rules
9657	A. van Soest, P. Fontein and Rob Euwals	Earnings Capacity and Labour Market Participation
9658	C. Fernández and M.F.J. Steel	On Bayesian Modelling of Fat Tails and Skewness
9659	R. Sarin and P. Wakker	Revealed Likelihood and Knightian Uncertainty
9660	J.R. Magnus and J. Durbin	A Classical Problem in Linear Regression or How to Estimate the Mean of a Univariate Normal Distribution with Known Variance
9661	U. Gneezy and J. Potters	An Experiment on Risk Taking and Evaluation Periods
9662	H.J. Bierens	Nonparametric Nonlinear Co-Trending Analysis, with an Application to Interest and Inflation in the U.S.
9663	J.P.C. Blanc	Optimization of Periodic Polling Systems with Non-Preemptive, Time-Limited Service
9664	M.J. Lee	A Root-N Consistent Semiparametric Estimator for Fixed Effect Binary Response Panel Data
9665	C. Fernández, J. Osiewalski and M.F.J. Steel	Robust Bayesian Inference on Scale Parameters
9666	X. Han and H. Webers	A Comment on Shaked and Sutton's Model of Vertical Product Differentiation
9667	R. Kollmann	The Exchange Rate in a Dynamic-Optimizing Current Account Model with Nominal Rigidities: A Quantitative Investigation
9668	R.C.H. Cheng and J.P.C. Kleijnen	Improved Design of Queueing Simulation Experiments with Highly Heteroscedastic Responses
9669	E. van Heck and P.M.A. Ribbers	Economic Effects of Electronic Markets
9670	F.Y. Kumah	The Effect of Monetary Policy on Exchange Rates: How to Solve the Puzzles

No.	Author(s)	Title
9671	J. Jansen	On the First Entrance Time Distribution of the $M/D/\infty$ Queue: a Combinatorial Approach
9672	Y.H. Farzin, K.J.M. Huisman and P.M. Kort	Optimal Timing of Technology Adoption
9673	J.R. Magnus and F.J.G.M. Klaassen	Testing Some Common Tennis Hypotheses: Four Years at Wimbledon
9674	J. Fidrmuc	Political Sustainability of Economic Reforms: Dynamics and Analysis of Regional Economic Factors
9675	M. Das and A. van Soest	A Panel Data Model for Subjective Information on Household Income Growth
9676	A.M. Lejour and H.A.A. Verbon	Fiscal Policies and Endogenous Growth in Integrated Capital Markets
9677	B. van Aarle and S.-E. Hougaard Jensen	Output Stabilization in EMU: Is There a Case for an EFTS?
9678	Th.E. Nijman, F.A. de Roon and C. Veld	Pricing Term Structure Risk in Futures Markets
9679	M. Dufwenberg and U. Gneezy	Efficiency, Reciprocity, and Expectations in an Experimental Game
9680	P. Bolton and E.-L. von Thadden	Blocks, Liquidity, and Corporate Control
9681	T. ten Raa and P. Mohnen	The Location of Comparative Advantages on the Basis of Fundamentals only
9682	S. Hochguertel and A. van Soest	The Relation between Financial and Housing Wealth of Dutch Households
9683	F.A. de Roon, Th.E. Nijman and B.J.M. Werker	Testing for Spanning with Futures Contracts and Nontraded Assets: A General Approach
9684	F.Y. Kumah	Common Stochastic Trends in the Current Account
9685	U.Gneezy and M. Das	Experimental Investigation of Perceived Risk in Finite Random Walk Processes
9686	B. von Stengel, A. van den Elzen and D. Talman	Tracing Equilibria in Extensive Games by Complementary Pivoting
9687	S.Tijs and M. Koster	General Aggregation of Demand and Cost Sharing Methods
9688	S.C.W. Eijffinger, H.P. Huizinga and J.J.G. Lemmen	Short-Term and Long-Term Government Debt and Nonresident Interest Withholding Taxes

No.	Author(s)	Title
9689	T. ten Raa and E.N. Wolff	Outsourcing of Services and the Productivity Recovery in U.S. Manufacturing in the 1980s
9690	J. Suijs	A Nucleolus for Stochastic Cooperative Games
9691	C. Seidl and S.Traub	Rational Choice and the Relevance of Irrelevant Alternatives
9692	C. Seidl and S.Traub	Testing Decision Rules for Multiattribute Decision Making
9693	R.M.W.J. Beetsma and H. Jensen	Inflation Targets and Contracts with Uncertain Central Banker Preferences
9694	M. Voorneveld	Equilibria and Approximate Equilibria in Infinite Potential Games
9695	F.B.S.L.P. Janssen and A.G. de Kok	A Two-Supplier Inventory Model
9696	L. Ljungqvist and H. Uhlig	Catching up with the Keynesians
9697	A. Rustichini	Dynamic Programming Solution of Incentive Constrained Problems
9698	G.Gürkan and A.Y. Özge	Sample-Path Optimization of Buffer Allocations in a Tandem Queue - Part I: Theoretical Issues
9699	H. Huizinga	The Dual Role of Money and Optimal Financial Taxes
96100	H. Huizinga	The Taxation Implicit in Two-Tiered Exchange Rate Systems
96101	H. Norde, F. Patrone and S. Tijs	Characterizing Properties of Approximate Solutions for Optimization Problems

P.O. BOX 99150 5000 LE TILBURG 1 NL
LANDS

Bibliotheek K. U. Brabant

17 000 01254673 6