Tilburg University

Cost allocation in the Chinese postman problem

Hamers, H.J.M.; Borm, P.E.M.; van den Leensel, A.; Tijs, S.H.

Published in:
European Journal of Operational Research

Publication date:
1999

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Hamers, H. J. M., Borm, P. E. M., van den Leensel, A., & Tijs, S. H. (1999). Cost allocation in the Chinese postman problem. *European Journal of Operational Research*, 118(1), 153-163.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Reprinted from

EUROPEAN JOURNAL OF OPERATIONAL RESEARCH

European Journal of Operational Research 118 (1999) 153–163

Theory and Methodology

Cost allocation in the Chinese postman problem

Herbert Hamers ^{a,*}, Peter Borm ^a, Robert van de Leensel ^b, Stef Tijs ^a

^a *CentER and Department of Econometrics, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands*

^b *Department of Quantitative Economics, University of Limburg, P.O. Box 616, 6200 MD Maastricht, The Netherlands*

Received 21 July 1997; accepted 3 August 1998

ELSEVIER

EUROPEAN JOURNAL OF OPERATIONAL RESEARCH

www.elsevier.com/locate/ejor

Editors

Roman Slowinski
IDSS/EJOR
Institute of Computing Science
Poznan University of Technology
Piotrowo 3a
60-965 Poznan, Poland
tel.: +48-61-8790790
fax: +48-61-8771525
email: slowinsk@man.poznan.pl

Jacques Teghem
MATHRO/EJOR
Faculté Polytechnique de Mons
9, Rue de Houdain
7000 Mons, Belgium
tel.: +32-65-374680
fax: +32-65-374689
email: teghem.ejor@mathro.fpms.ac.be

Jyrki Wallenius
Helsinki School of Economics
P.O. Box 1210
00101 Helsinki, Finland
tel.: +358-9-43131
fax: +358-9-43138613
email: walleniu@hkkk.fi

Editors OR-Software Section

Hervé Thiriez, Centre HEC-ISA, 1, rue de la Libération, 78350 Jouy-en-Josas, France
(ORSEP) Horst W. Hamacher, FB Mathematik, Universität Kaiserslautern, Postfach 3049, 67618 Kaiserslautern, Germany

Editor Book Review Section

H. Fleuren, Centre for Quantitative Methods CQM B.V., P.O. Box 414, 5600 AK Eindhoven, Netherlands

Editorial board

Sven Axäster, Lund University, Lund, Sweden
Achim Bachem, German Aerospace Research Establishment,
Cologne, Germany
Egon Balas, Carnegie-Mellon University, Pittsburgh, USA
Paul van Beek, Agricultural University Wageningen, Wageningen,
Netherlands
M. Bielli, Consiglio Nazionale delle Ricerche, Roma, Italy
Jean-Pierre Brans, Vrije Universiteit Brussel, Brussels, Belgium
C.M. Brugga, University College Dublin, Dublin, Ireland
Derek Bunn, London University, London, UK
Rainer Burkard, Technische Universität Graz, Graz, Austria
Christer Carlson, Åbo University, Åbo, Finland
William Cooper, The University of Texas at Austin, Austin, USA
Jehoshua Eliashberg, University of Pennsylvania, Philadelphia, USA
Laureano Escudero, University of Madrid, Madrid, Spain
A.A. Farley, Monash University, Clayton, Australia
Ludo Gelders, Katholieke Universiteit Leuven, Heverlee-Leuven,
Belgium
Brian Haley, University of Birmingham, Birmingham, UK
Masao Iri, Chuo University, Tokyo, Japan
Søren Kruse Jacobsen, The Technical University of Denmark, Lyngby,
Denmark

Jack Kleijnen, Tilburg University, Tilburg, Netherlands
Istvan Maros, Imperial College, London, UK
Heiner Müller-Merbach, Universität Kaiserslautern, Kaiserslautern,
Germany
Toshio Nakagawa, Aichi Institute of Technology, Tokyo, Japan
Giacomo Patrizi, University of Rome "La Sapienza", Rome, Italy
Charles ReVelle, The Johns Hopkins University, Baltimore, USA
Marc Roubens, University of Liège, Liège, Belgium
B. Roy, Université de Paris-IX-Dauphine, Paris, France
Y. Siskos, Technical University of Crete, Chania, Greece
Yukio Takahashi, Tokyo Institute of Technology, Tokyo, Japan
Luis Tavares, Universidade Técnica de Lisboa, Lisboa, Portugal
Paolo Toth, University of Bologna, Bologna, Italy
John Ulhøi, The Århus School of Economics and Business Adminis-
tration, Århus, Denmark
Gunduz Ulusoy, Bogazici University, Istanbul, Turkey
Gerhard Wäscher, Martin-Luther-Universität, Halle (Saale), Germany
Jan Weglarz, Polytechnika Poznanska Instytut Informatyki, Poznan,
Poland
Dominique de Werra, Ecole Polytechnique Fédérale de Lausanne,
Lausanne, Switzerland
Uri Yechiali, Tel Aviv University, Tel Aviv, Israel

Editorial policy

Operational Research (OR) involves the application of scientific methods to the management of complex systems of people, machinery, materials, money and information. It seeks to produce an understanding of managerial problems and to develop models which will enable the consequences of decisions to be investigated. OR methods have already had widespread application and success in many areas of business, industry and government, and their use is rapidly increasing in social systems and in the international arena. The European Journal of Operational Research (EJOR) is intended to strengthen these advances by publishing high-quality, original papers that contribute to the practice of decision making, within and beyond Europe, irrespective of whether their content describes an application or a theoretical development. A special call is made for case study papers showing OR in action in firms and other organizations. Mathematical or computer-science contributions should be capable of being used in the foreseeable future.

EJOR has regular and feature issues. Almost all issues contain a leading invited review explaining the developments in an OR topic over the past five years to the general OR reader. In addition to the other five types of contribution described in the Instructions to Authors at the end of this issue, EJOR includes OR software news (including ORSEP) and reviews of books published recently. For further information, see the Instructions to Authors at the end of each issue. Authors of contributions are asked to read these Instructions before sending four copies to any of the editors at the address above.

Theory and Methodology

Cost allocation in the Chinese postman problem

Herbert Hamers ^{a,*}, Peter Borm ^a, Robert van de Leensel ^b, Stef Tijs ^a^a CentER and Department of Econometrics, Tilburg University, P.O. Box 90153, 5000 LE Tilburg, The Netherlands^b Department of Quantitative Economics, University of Limburg, P.O. Box 616, 6200 MD Maastricht, The Netherlands

Received 21 July 1997; accepted 3 August 1998

Abstract

This paper considers a cost allocation problem that arises from a delivery problem associated with the Chinese postman problem (CPP). A delivery problem is described by a connected undirected graph in which each edge belongs to a different player, a cost function on the edges of this graph and a fixed vertex which is referred to as the post office. Assume that the post office is providing some service to the players. The nature of this service, which can be thought of as mail delivery, requires that a server will travel along the edges of the graph and returns to the post office. The cost allocation problem is concerned with the cost of providing the service to all players. A specific cost allocation rule is introduced and characterized. Further, the class of delivery problems gives rise to a new class of cooperative combinatorial optimization games called delivery games. It is shown that the outcome of the allocation rule with respect to a bridge-connected Euler graph is a core element of the corresponding delivery game. © 1999 Elsevier Science B.V. All rights reserved.

Keywords: Chinese postman problem; Cost allocation; Delivery game

1. Introduction

Since the early days of operations research and non-cooperative game theory there has been interaction between the two fields. We mention:

- (i) Duality result in mathematical programming and minimax results in zero sum game theory (Dantzig, 1963).
- (ii) Linear complementarity problems and the theory of bimatrix games (Lemke and Howson, 1964).
- (iii) Markov decision theory and the theory of stochastic games (Shapley, 1953).
- (iv) Optimal control theory and the theory of differential games (Isaacs, 1965).

* Corresponding author. Tel.: 013-466-2660; e-mail: h.j.m.hamers@kub.nl

In the last two decades one also sees many interesting interrelations between operations research and cooperative game theory, where often two problems come together: on the one hand an optimization problem and on the other hand a cost allocation problem. We mention

- (i) Minimum spanning tree games and spanning network games (Claus and Kleitman, 1973; Bird, 1976; Megiddo, 1978; Granot and Huberman, 1981).
- (ii) Linear production games and mathematical programming games (Owen, 1975; Granot, 1986).
- (iii) Flow games (Kalai and Zemel, 1982; Derks and Tijs, 1986).
- (iv) Traveling salesman games (Potters et al., 1992; Tamir, 1989).
- (v) Sequencing games (Curiel et al. (1989)).

This paper will focus on a new interrelation between operations research and cooperative game theory. We consider delivery problems corresponding to situations where from a central depot (post office) deliveries are to be made to customers and where on the one hand the central depot wants to deliver as cheaply as possible and on the other hand there is the problem of allocating a fair share of the overall costs to each customer. This problem is related to the Chinese postman problem (CPP), introduced by Mei-Ko Kwan (1962). Here an undirected connected graph and a cost function on the edges are involved. The problem is to find a route of minimal costs that visits each edge (street) at least once. Edmonds and Johnson (1973) provide a polynomial algorithm that solves this problem.

In fact, this paper considers a cost allocation problem that arises from the CPP. Let there be an undirected connected graph (street plan). In the model we identify each edge (street) with a player of the player set N and there is a fixed vertex called the post office, where all routes of the postman will start and finish. The postman has to make an S -tour to deliver a coalition $S \subseteq N$. An S -tour is a tour of the postman in which the mail can be delivered in each street of coalition S . So, an S -tour visits each edge of S at least once, but may use streets outside S to reach the streets of S . For an S -tour we have that the postman charges for each time a specific street is visited the corresponding travel costs. We assume that in any S -tour each street of S pays once his own specific travel costs itself. Hence, each player of S has individually fixed costs. The sum over the members of S of these fixed costs in an S -tour is called the separable costs of an S -tour. Note that these separable costs are independent of the chosen S -tour. The remaining travel costs corresponding to an S -tour we will call the non-separable costs. These have to be paid in some way by the members of S . Obviously, a coalition S wants to be served by an S -tour which minimizes the non-separable costs. Such an S -tour is called a minimal S -tour. Note that the grand coalition N , which consists of all edges, faces a CPP to obtain a minimal N -tour. The general problem is whether one can find an allocation of the non-separable costs of a minimal N -tour that is stable in the sense that no (sub)coalition has an incentive to split off and to have the mail delivered separately. For this purpose we investigate delivery games, a class of cooperative combinatorial optimization games that arise from delivery problems. The costs of a coalition S in a delivery game is given by the non-separable costs of a minimal S -tour. The core of this game consists of all allocations which are stable in the way described above.

Section 2 provides the formal description of the delivery problem. In Section 3 we consider delivery problems corresponding to bridge-connected Euler graphs, i.e. graphs in which all components that arise by removing the bridges are Euler graphs. For this class of delivery models we introduce and characterize a specific rule γ that divides the non-separable costs of a minimal N -tour among all players. Cooperative delivery games related to delivery models are introduced in Section 4. It is shown that, in general, delivery games may have an empty core, i.e. no stable allocation exists. For bridge-connected Euler graphs, however, it can be shown that the outcome of γ is always a core element. Finally, in Section 5, an extension of the rule γ is provided to a more general class of graphs. It turns out that this extension can be characterized along the same lines as γ but need not be in the core of the corresponding delivery game.

2. The delivery problem

For the formal description of a delivery problem we need some preliminary definitions. Let $G = (V, E, i)$ be an undirected connected graph where V is the set of vertices, E the set of edges and i the incidence mapping that assigns to each edge of E an unordered pair of not necessarily different vertices of V . A *walk* in G is a finite sequence of edges and vertices of the form $v_1, e_1, v_2, \dots, e_k, v_{k+1}$ with $k \geq 0, v_1, \dots, v_{k+1} \in V, e_1, \dots, e_k \in E$ such that $i(e_j) = \{v_j, v_{j+1}\}$ for all $j \in \{1, \dots, k\}$. Such a walk is a *closed walk* if $v_1 = v_{k+1}$. A closed walk in which all edges are distinct is called a *closed trail*. A *path* in G is a walk in which all vertices (except, possibly $v_1 = v_{k+1}$) and edges are distinct. A closed path, i.e. $v_1 = v_{k+1}$, containing at least one edge is called a *circuit*.

We assume that each edge corresponds to one player. Formally, there is a one-to-one function $g: E \rightarrow N$, where $N = \{1, \dots, |E|\}$ is the set of players. Let $v_0 \in V$ and $S \subseteq N$. An S -tour w.r.t. v_0 is a closed walk $v_0, e_1, v_1, \dots, v_{k-1}, e_k, v_0$ in G such that $S \subset \{g(e_j) \mid j \in \{1, \dots, k\}\}$.

An S -tour visits each edge that corresponds to a player of S and starts and finishes in the same vertex v_0 . Note that an edge can be visited more than once by an S -tour. Note that an S -tour may also use edges outside the coalition S . We will assume that any S -tour for all $S \subseteq N$ will start and finish in the same vertex. This vertex is denoted by v_0 and we will refer to it as the post office. Further, we denote the set of all S -tours by $D(S)$.

To the edges of the graph G we assign travel costs $t: E \rightarrow [0, \infty)$. Suppose a coalition S is served according to the S -tour $v_0, e_1, v_1, \dots, v_{k-1}, e_k, v_0 \in D(S)$. The total costs of this tour are $\sum_{j=1}^k t(e_j)$. We will assume that each player $g(e) \in S$ (once) pays the travel costs $t(e)$ by himself. In this way we already allocated the *separable costs* $\sum_{e: g(e) \in S} t(e)$ of an S -tour. Note that these separable costs are independent of the chosen S -tour. The remaining *non-separable costs* C_S for coalition S ,

$$C_S(v_0, e_1, \dots, e_k, v_0) := \sum_{j=1}^k t(e_j) - \sum_{e: g(e) \in S} t(e), \quad (1)$$

have to be allocated in some way to the members of S . This implies that each time the postman revisits an edge of S or visits an edge that corresponds to a player outside S , the coalition S has to pay the travel costs of this edge. Since each coalition S wants to be delivered as cheaply as possible, it will choose an S -tour in $D(S)$ that minimizes the non-separable costs as given in Eq. (1). Such an S -tour is called *minimal*.

As an illustration we consider the following example.

Example 1. Consider the delivery problem with $N = \{1, 2, 3, 4, 5\}$ as described in Fig. 1. Here $g(e_j) = j$ for all $j \in N$ and all edges have travel costs 1.

Fig. 1.

Each N -tour will contain e_1 at least twice. The N -tour $v_0, e_1, v_1, e_2, v_2, e_3, v_3, e_4, v_4, e_5, v_1, e_1, v_0$ visits the edges e_2, e_3, e_4, e_5 once and e_1 is visited twice. Hence, this is a minimal N -tour and the non-separable costs for coalition N are equal to $t(e_1) = 1$. Note that this minimal N -tour is also a $\{2\}$ -tour for coalition $\{2\}$. Then the non-separable costs for coalition $\{2\}$ of this $\{2\}$ -tour are equal to $t(e_1) + t(e_3) + t(e_4) + t(e_5) + t(e_1) = 5$. A minimal $\{2\}$ -tour is $v_0, e_1, v_1, e_2, v_2, e_2, v_1, e_1, v_0$ where the non-separable costs are equal to $t(e_1) + t(e_2) + t(e_1) = 3$.

In the following a delivery problem is denoted by $\Gamma = (N, (V, E, i, v_0), t, g)$. Here, N is the set of players, (V, E, i, v_0) is a connected graph in which v_0 is the post office, $t : E \rightarrow [0, \infty)$ assigns the travel costs to the edges and g gives the one-one correspondence between players and edges. The class of delivery problems corresponding to the player set N is denoted by $DP(N)$. We will study the cost allocation problem of the non-separable costs of a minimal N -tour.

3. A cost allocation rule for bridge-connected Euler graphs

This section considers delivery problems $\Gamma = (N, (V, E, i, v_0), t, g)$ where $G = (V, E, i)$ is a bridge-connected Euler graph. For this class of delivery problems we introduce and characterize a specific cost allocation rule γ that divides the non-separable costs of a minimal N -tour.

First we recall the definition of an Euler graph. A connected graph G is called an *Euler graph* if there exists a closed trail that includes every edge of G . This implies that for a delivery problem in which the underlying graph is an Euler graph there exists an N -tour that visits each edge of E exactly once. Obviously, this tour is a minimal N -tour since the non-separable costs equal zero. Hence, an (obvious) division of the non-separable costs in this case is the vector $(0, 0, \dots, 0) \in \mathbb{R}^V$.

Let $G = (V, E, i)$ be an arbitrary connected graph. Then the set of *bridges* $B(G)$ in G is defined by

$$B(G) := \{b \in E \mid (V, E - \{b\}, i_{|E - \{b\}}) \text{ is not connected}\}.$$

Hence, in case $i(b) = \{v, w\}$ with $v \neq w$ we have that $b \in B(G)$ if and only if there is no path between v and w in the graph $(V, E - \{b\}, i_{|E - \{b\}})$. Now, we can introduce the class of bridge-connected Euler graphs. A connected graph $G = (V, E, i)$ is called a *bridge-connected Euler graph* if all the components of the graph $\hat{G} = (V, E - B(G), i_{|E - B(G)})$ are Euler graphs or singletons.

Note that the graph $\hat{G}$ is derived from G by removing all bridges in G . Since all components of $\hat{G}$ are Euler graphs there exists in each component a closed trail. The components are connected by the elements of $B(G)$. Hence, we can make a closed walk in G that visits each bridge twice and all other edges in G once. Obviously, this closed walk is an N -tour in the delivery problem $(N, (G, v_0), t, g)$. Since each N -tour starts and finishes in v_0 it follows from the definition of a bridge that each bridge is visited at least twice in any N -tour. This implies that this specific closed walk is a minimal N -tour for the delivery problem $(N, (G, v_0), t, g)$. Moreover, it readily follows that the non-separable costs of this minimal N -tour equal $\sum_{b \in B(G)} t(b)$ (cf. (1)).

The following example illustrates a bridge-connected Euler graph.

Example 2. Let G be the graph as shown in Fig. 2. The bridges of G are the edges b_1, b_2 and b_3 . The graph $\hat{G}$ is derived from G by removing all bridges of G . Since the components E_0, E_1, E_2 and E_3 of $\hat{G}$ are Euler graphs or singletons, we have that G is a bridge-connected Euler graph.

The class of delivery problems corresponding to bridge-connected Euler graphs and player set N is denoted by $DE(N)$. For introducing a rather intuitive rule on $DE(N)$ that divides for each $\Gamma \in DE(N)$ the non-separable costs of a minimal N -tour among the members of N we need the notion of followers of a bridge with respect to v_0 in a bridge-connected Euler graph $G = (V, E, i)$. Let $v_0 \in V$ be the post office and

Fig. 2.

let $b \in B(G)$ be a bridge. Then $e \in E$ is a *follower* of b with respect to v_0 if and only if each path $v_0, e_1, \dots, e_k, v_{k+1}$ with $e_k = e$ also contains b . The set of followers of b will be denoted by $F_b(G, v_0)$. Note that $b \in F_b(G, v_0)$ and that the set of followers depends on the location of v_0 in the graph. Let $b \in B(G)$. Clearly the postman needs this bridge to visit each edge in $F_b(G, v_0)$. We propose that each player in $g(F_b(G, v_0))$ will pay an equal share of the travel costs of b . So, if a player needs several bridges to be visited, he has to contribute a fair share of the costs of all these bridges. Formally, the division rule $\gamma : \text{DE}(N) \rightarrow \mathbb{R}^N$ is defined for all $\Gamma = (N, (V, E, i, v_0), t, g) \in \text{DE}(N)$ with $G = (V, E, i)$ by

$$\gamma_{g(e)}(\Gamma) = \sum_{b \in B(G), e \in F_b(G, v_0)} \frac{t(b)}{|F_b(G, v_0)|} \quad \text{for all } e \in E. \quad (2)$$

Note that $\gamma_{g(e)}(\Gamma)$ represents the costs of player $g(e)$.

Now we will provide a characterization of the division rule γ . For this we need the notions of bridge-clusters of a bridge-connected Euler graph and the condensation of a graph with respect to an extreme bridge. Let $\Gamma = (N, (V, E, i, v_0), t, g) \in \text{DE}(N)$. Let $G = (V, E, i)$. A *bridge-cluster* is a set of edges that needs the same set of bridges to be connected to the post office. So, formally, we have at most $|B(G) + 1|$ bridge clusters $C_0(G, v_0)$ and $\{C_b(G, v_0)\}_{b \in B(G)}$. Here,

$$C_0(G, v_0) = E - \bigcup_{b \in B(G)} F_b(G, v_0), \quad (3)$$

is the set of edges that do not need any bridge to be connected to v_0 . Note that this set may be empty. For $b \in B(G)$,

$$C_b(G, v_0) = F_b(G, v_0) - \bigcup_{\{b' \in B(G) \cap F_b(G, v_0), b' \neq b\}} F_{b'}(G, v_0), \quad (4)$$

is the cluster of edges that needs the set $\{b^* \in B(G) \mid b \in F_{b^*}(G, v_0)\}$ to be connected to v_0 . A bridge $b \in B(G)$ is called an *extreme bridge* if it has no other bridge as a follower, or equivalently if $C_b(G, v_0) = F_b(G, v_0)$. The following example gives the bridge-clusters and extreme bridges of a bridge-connected Euler graph.

Example 3. Take the graph G from Example 2. Then $C_0(G, v_0) = \emptyset$ and for each $j \in \{1, 2, 3\}$ the bridge-cluster $C_j(G, v_0)$ consists of b_j and the edges of component E_j . The extreme bridges are b_2 and b_3 .

In the following we describe the procedure to construct a so-called *condensed graph* of a bridge-connected Euler graph with respect to an extreme bridge. Let $\Gamma = (N, (G, v_0), t, g)$ and $G = (V, E, i)$. Let $b \in B(G)$ be an extreme bridge of G . Let $v_1^* \in i(b)$ be such that there exists a path in the graph $(V, E - \{b\}, i_{E-\{b\}})$ between v_0 and v_1^* . Let $F_b(G, v_0)$ be the set of followers of b and let $V(F_b(G, v_0))$ be the vertices incident with the edges of $F_b(G, v_0)$. The graph $\underline{G}$ is derived from G by removing all edges in $F_b(G, v_0)$ and vertices $V(F_b(G, v_0)) \setminus \{v_1^*\}$. With $|F_b(G, v_0)| = m$, the graph G^* arises from the graph $\underline{G}$ by constructing a circuit defined by $v_1^*, e_1^*, v_2^*, \dots, v_m^*, e_m^*, v_1^*$ where $\{v_1^*, v_2^*, \dots, v_m^*\} \cap V = v_1^*$ and $\{e_1^*, \dots, e_m^*\} \cap E = \emptyset$ to the vertex v_1^* . The graph G^* is called the *condensed graph* of G with respect to the extreme bridge b . Note that G^* is also a bridge-connected Euler graph. Moreover, the number of edges in G and in G^* coincide and $|B(G^*)| = |B(G)| - 1$.

Example 4. Take the graph G of example 2. Fig. 3 shows the graph G^* that arises from G by condensation with respect to the extreme bridge b_2 . In an obvious way we can now define the *condensed delivery problem* $\Gamma_b = (N, (V^*, E^*, i^*, v_0), t^*, g^*)$ with respect to an extreme bridge $b \in B(G)$. Here, $G^* = (V^*, E^*, i^*)$ is the condensed graph of G with respect to b . Further, the travel costs $t^* : E^* \rightarrow [0, \infty)$ are defined by

$$t^*(e) = \begin{cases} t(e) & \text{if } e \in E - F_b(G), \\ 0 & \text{otherwise,} \end{cases}$$

and for the one-one map $g^* : E^* \rightarrow N$ it holds that $g^*(e) = g(e)$ if $e \in E - F_b(G)$ and $g^*({e_1^*, \dots, e_m^*}) = g(F_b(G))$. So, in fact we have defined a whole class of condensed delivery problems.

Now consider the following three properties for a division rule $f : \text{DE}(N) \rightarrow \mathbb{R}^N$.

Efficiency: Let $G = (V, E, i)$ and let $\Gamma = (N, (G, v_0), t, g) \in \text{DE}(N)$. Then it holds that $\sum_{i \in N} f_i(\Gamma) = \sum_{b \in B(G)} t(b)$.

Bridge-cluster symmetry: Let $G = (V, E, i)$ and let $\Gamma = (N, (G, v_0), t, g) \in \text{DE}(N)$. Then $f_{g(e_1)}(\Gamma) = f_{g(e_2)}(\Gamma)$ for all $e_1, e_2 \in C_0(G, v_0)$ and for any $b \in B(G)$ it holds that $f_{g(e_1)}(\Gamma) = f_{g(e_2)}(\Gamma)$ for all $e_1, e_2 \in C_b(G, v_0)$.

Fig. 3.

Condensation independence: Let $G = (V, E, i)$ and let $\Gamma = (N, (V, E, i, v_0), t, g) \in \text{DE}(N)$. If b is an extreme bridge of G and $\Gamma_b = (N, (V^*, E^*, i^*, v_0), t^*, g^*)$ is a condensed problem of Γ with respect to b , then $f_{g(e)}(\Gamma) = f_{g^*(e)}(\Gamma_b)$ for all $e \in E - F_b(G, v_0)$.

Efficiency states that the non-separable costs of a minimal N -tour are fully accounted for. Bridge-cluster symmetry states that each group of players that need the same set of bridges to be connected with the post office will contribute the same part to the non-separable costs. Condensation independence compares cost shares in a delivery problem and a corresponding condensed delivery problem. All players who are not in the bridge-cluster corresponding to the removed extreme bridge face in this reduced graph the same problem to reach the post office as in the original graph. The condensation independence claims that in both situations this rule assigns to each member of this group of players the same cost share.

Theorem 1. *The allocation rule $\gamma : \text{DE}(N) \rightarrow \mathbb{R}^N$ is the unique rule that satisfies efficiency, bridge-cluster symmetry and the condensation property.*

Proof. It is easy to check that γ satisfies the three properties. Let $f : \text{DE}(N) \rightarrow \mathbb{R}^N$ be a rule that satisfies the three properties. We show that $f = \gamma$ by induction to the number of bridges of $G = (V, E, i)$. First, let $|B(G)| = 0$. Then efficiency and bridge-cluster symmetry implies that $f(\Gamma) = (0, \dots, 0) = \gamma(\Gamma)$. Assume that if $|B(G)| = q$ then $f(\Gamma) = \gamma(\Gamma)$. Let $|B(G)| = q + 1$. Let b be an extreme bridge of G . Let $|F_b(G, v_0)| = m$ and let $\Gamma_b = (N, (V^*, E^*, i^*, v_0), t^*, g^*)$ be a condensed problem of Γ with respect to b . Obviously, the underlying graph of Γ_b has q bridges. Then, by induction we have that

$$\gamma_{g(e)}(\Gamma_b) = f_{g^*(e)}(\Gamma_b) \quad \text{for all } e \in E^*. \quad (5)$$

The condensation property yields that

$$\gamma_{g(e)}(\Gamma) = \gamma_{g(e)}(\Gamma_b) \quad \text{for all } e \in E - F_b(G, v_0), \quad (6)$$

$$f_{g(e)}(\Gamma) = f_{g(e)}(\Gamma_b) \quad \text{for all } e \in E - F_b(G, v_0). \quad (7)$$

From Eqs. (5)–(7) it follows immediately that

$$\gamma_{g(e)}(\Gamma) = f_{g(e)}(\Gamma) \quad \text{for all } e \in E - F_b(G, v_0). \quad (8)$$

Efficiency and Eq. (8) gives that

$$\sum_{e \in F_b(G, v_0)} \gamma_{g(e)}(\Gamma) = \sum_{e \in F_b(G, v_0)} f_{g(e)}(\Gamma). \quad (9)$$

Since b is an extreme bridge we have that $C_b(G, v_0) = F_b(G, v_0)$. Bridge-cluster symmetry implies that

$$\gamma_{g(e)}(\Gamma) = \gamma_{g(b)}(\Gamma) \quad \text{for all } e \in F_b(G, v_0), \quad (10)$$

$$f_{g(e)}(\Gamma) = f_{g(b)}(\Gamma) \quad \text{for all } e \in F_b(G, v_0). \quad (11)$$

Substitution of Eqs. (10) and (11) in Eq. (9) and the fact that $|F_b(G, v_0)| = m$ gives

$$m\gamma_{g(b)}(\Gamma) = mf_{g(b)}(\Gamma)$$

and, consequently, $\gamma_{g(b)}(\Gamma) = f_{g(b)}(\Gamma)$. From Eqs. (10) and (11) it then follows that

$$\gamma_{g(e)}(\Gamma) = f_{g(e)}(\Gamma) \quad \text{for all } e \in F_b(G, v_0). \quad (12)$$

This finishes the proof. $\square$

4. Delivery games

In this section we introduce to each delivery problem a corresponding cooperative delivery game. First, we recall some well known facts concerning cooperative games. A *cooperative cost game* is a pair (N, c) where N is a finite set of players and c is a mapping $c : 2^N \rightarrow \mathbb{R}$ with $c(\emptyset) = 0$. Here, 2^N is the collection of all subsets of N (coalitions).

Cooperative game theory focuses on ‘fair’ and/or ‘stable’ division rules from the value $c(N)$ of the grand coalition. A core element $x = (x_i)_{i \in N} \in \mathbb{R}^N$ is such that no coalition $S \subset N$ has an incentive to split off, i.e.

$$\sum_{i \in N} x_i = c(N) \text{ and } \sum_{i \in S} x_i \leq c(S) \text{ for all } S \in 2^N.$$

The core $\text{Core}(c)$ consists of all core elements. A game is called balanced if its core is non-empty.

In Section 2 we defined a minimal S -tour of a coalition S . By defining the cost value of a coalition S as the non-separable costs of a minimal S -tour (cf. (1)), we obtain the delivery game corresponding to the delivery problem.

Definition 1. The cooperative delivery game (N, c) corresponding to $\Gamma = (N, (V, E, i, v_0), t, g) \in \text{DP}(N)$ is defined for all $S \subseteq N$ by

$$c(S) := \min_{t_0, e_1, \dots, e_k, t_k \in D(S)} \left(\sum_{j=1}^k t(e_j) - \sum_{e: g(e) \in S} t(e) \right).$$

In the following example a delivery game is considered.

Example 5. Take the delivery problem of Example 1. In Example 1 it is shown that the non-separable costs of a minimal N -tour for coalition N equal 1 and that the non-separable costs of the minimal $\{2\}$ -tour for coalition $\{2\}$ equal 3. Hence, $c(N) = 1$ and $c(\{2\}) = 3$. Since for each coalition S the edge e_1 is contained two times in all S -tours, we have that $c(S) \geq t(e_1) = 1$. Since $c(N) = 1$ we have that e.g. $(\frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}, \frac{1}{5}) \in \text{Core}(c)$.

The following example shows that a delivery game is not necessarily balanced.

Example 6. Consider the delivery problem with $N = \{1, 2, 3, 4, 5\}$ as described in Fig. 4 in which $g(e_j) = j$ and $t(e_j) = 1$ for all $j \in \{1, \dots, 5\}$.

Then $c(N) = 1$ and $c(S) = 0$ if $S \in \mathcal{A} := \{\{1, 2, 5\}, \{3, 4, 5\}, \{1, 2, 3, 4\}\}$. Suppose $x \in \text{Core}(c)$. Then

Fig. 4.

$$0 < 2c(N) = \sum_{S \in \mathcal{S}} \sum_{i \in S} x_i \leq \sum_{S \in \mathcal{S}} c(S) = 0.$$

Contradiction, so no core elements exists.

For $\Gamma \in \text{DE}(N)$, i.e. a delivery problem corresponding to bridge-connected Euler graph, it will be shown that $\gamma(\Gamma)$ is an element of the core of the corresponding delivery game. Consequently, each delivery game that corresponds to a delivery problem of $\text{DE}(N)$ is balanced.

Theorem 2. Let $\Gamma = (N, (G, v_0), t, g) \in \text{DE}(N)$ and let (N, c) be the corresponding delivery game. Then $\gamma(\Gamma) \in \text{Core}(c)$.

Proof. Obviously, $\sum_{e \in E} \gamma_{g(e)}(\Gamma) = c(N)$. Let $S \subset N$ and $g^{-1}(S) = A \subset E$. Then A can be partitioned into at most $|B(G)| + 1$ sets $A_0, \{A_b\}_{b \in B(G)}$ with $A_0 = A \cap C_0(G, v_0)$ and $A_b = A \cap C_b(G, v_0)$. Then

$$\begin{aligned} \sum_{i \in S} \gamma_i(\Gamma) &= \sum_{e \in A} \sum_{b \in B(G): e \in F_b(G, v_0)} \frac{t(b)}{|F_b(G, v_0)|} \\ &= \sum_{b' \in B(G)} \sum_{e \in A_{b'}} \sum_{b \in B(G): e \in F_b(G, v_0)} \frac{t(b)}{|F_b(G, v_0)|} = \sum_{b' \in B(G)} \sum_{b \in B(G): F_b(G, v_0) \cap A_{b'} \neq \emptyset} \frac{|A_{b'}| t(b)}{|F_b(G, v_0)|} \\ &= \sum_{b \in B(G): F_b(G, v_0) \cap A \neq \emptyset} \sum_{b' \in B(G): F_b(G, v_0) \cap A_{b'} \neq \emptyset} \frac{|A_{b'}| t(b)}{|F_b(G, v_0)|} \leq \sum_{b \in B(G): F_b(G, v_0) \cap A \neq \emptyset} t(b) \\ &\leq c(S). \quad \square \end{aligned}$$

Hamers (1997) shows that delivery games corresponding to bridge-connected Euler graphs need not be concave, but that delivery games corresponding to bridge-connected cyclic graphs are concave. Here, a connected graph $G = (V, E, i)$ is called a bridge-connected cyclic graph if the components of the graph $\hat{G} = (V, E - B(G), i|_{E - B(G)})$ are cycles or single vertices. A cycle is the union of edge-disjoint circuits. Note that the class of bridge-connected cyclic graphs is contained in the class of bridge-connected Euler graphs. Granot et al. (1996) imposed necessary and sufficient conditions on a graph such that the associated delivery game is balanced, totally balanced and concave, respectively.

5. Concluding remarks

This section discusses an extension of the division rule γ to a more general class of delivery problems.

For this, we need the observation that any connected graph $G = (V, E, i)$ can be considered as a generalization of a bridge-connected Euler graph. This follows from the fact that the set of edges E can be partitioned into two subsets: the edges that are contained in a circuit and the edges that are a bridge. Hence, after the removal of all bridges in an arbitrary connected graph we get components which either consists of a single vertex or consists of a connected (sub)graph in which each edge is contained in at least one circuit. Let $v_0 \in V$ be the post office in G , then $B(G)$ is the set of bridges of G and the set of followers $F_b(G, v_0)$ of $b \in B(G)$ is defined in the same way as in Section 3. Moreover, a bridge-cluster is again a set of edges that need the same set of bridges to be connected to the post office (cf. (3) and (4)). We restrict our attention to graphs in which v_0 is connected only to bridges, which implies that $C_0(G, v_0) = \emptyset$, or v_0 is not connected to any bridge of G . For this class of graphs we have that, whenever $C_0(G, v_0) \neq \emptyset$, each edge needs at least one edge of $C_0(G, v_0)$ to be connected to v_0 .

ASSOCIATION OF EUROPEAN OPERATIONAL RESEARCH SOCIETIES (EURO)
within the International Federation of Operational Research Societies (IFORS)

EURO-goals

In the belief that the purpose of Operational Research is to improve the well-being of people by improving the relevance and effectiveness of the institutions and organisations which serve them; that it seeks to do so by means of the rational methods of science exercised by representatives of diverse disciplines working together, each supporting the others; that this co-operation which respects no boundaries between disciplines should respect no boundaries between peoples; and that the Operational Research Societies of Europe should more closely co-operate one with another to further the theory and practice of Operational Research, the European Operational Research Societies established in 1975 the Association of European Operational Research Societies (EURO) within IFORS. The members of EURO agree to grant any fully paid-up member of any signatory body all rights and privileges as are offered by them to their own members, to exchange all appropriate information, e.g. bulletins, etc., to include relevant information from other signatories in their bulletin, etc., to inform other signatories of existing working groups and the dates and locations of their meetings, to open such working groups to individual members of other signatories, to organise European conferences on Operational Research and European working groups on topics which are felt to be important by the signatories, to encourage either individually or in concert the formation of Operational Research Societies in other European countries and to give such new bodies any possible help they may require.

EURO-officers

President: Jan Weglarz, Institute of Computing Science, Technical University of Poznan, Piotrowo 3A, PL-60965 Poznan, Poland
Past-President: Paolo Toth, D.E.I.S., University of Bologna, Viale Risorgimento 2, I-40136 Bologna, Italy
Vice-President 1: Valerie Belton, Department of Management Science, University of Strathclyde, 40 George Street, Glasgow G1 1QE, UK
Vice-President 2: Raymond Bisdorff, CRP - Centre Universitaire, Cellule Stade, 13, rue de Braganca, L-1255 Luxembourg, Luxembourg
Secretary: Denis Bouyssou, ESSEC, Avenue Bernard Hirsch, B.P. 105, F-95021 Cergy-Pontoise Cedex, France
Treasurer: Marino Widmer, Institut d'Informatique, Université de Fribourg, Site de Regina Mundi, Rue Faucigny 2, CH-1700 Fribourg, Suisse
Past Presidents: H. Zimmermann, B. Rapp, R. Tomlinson, J. Brans, B. Roy, D. de Werra, J. Krarup, J. Spronk, M.F. Shuttler, P. Toth

EURO-members

Austria: ÖGOR, c/o Institut für Ökonometrie, OR und Systemtheorie, Technische Universität Wien, Argentinierstraße 8, A-1040 Wien
Belgium: SOGESCI/BVWB, Renaissance laan 30, Av. de la Renaissance, B-1040 Brussels
Bulgaria: BORS, Institute of Informatics, Bulgarian Academy of Sciences, Acad. G. Bonchev Str. Bl. 29A, 1113 Sofia
Croatia: CRORS, Ekonomski fakultet, Kennedyjev trg. 6, 41000 Zagreb

Czech Republic: CSOV/CSOR, Department of Econometrics, University of Economics, Nam. W. Churchilla 4, CZ 130 67 Praha 3
Denmark: DORS, c/o IMM, bldg. 321, Technical University of Denmark, DK-2800 Lyngby
Finland: FORS, c/o Rauni Seppola, Helsinki School of Economics, Runeberginkatu 14-16, FIN-00100 Helsinki
France: AFCET, 156 Bld. Péreire, F-75017 Paris
Germany: DGOR, c/o Mrs. Bärbel Niedzwetzki, Am Steinknapp 14b, D-44795 Bochum
Greece: HELORS, Solomou 20, 5th floor, att. Kath. Papanastassatou, 106 82 Athens
Hungary: HORS, MTA SZTAKI, Kende u. 13-17, H-1111 Budapest
Iceland: ICORS, University of Iceland, Faculty of Economics and Business Administration, IS-101 Reykjavik
Ireland: ORMSI, c/o Mr D.P. Darcy, Department of Management Information Systems, Arts-Commerce Building, University College Dublin, Belfield, IRL-Dublin 4
Israel: ORSIS, c/o Faculty of Industrial Engineering and Management, Technion, Haifa 32000
Italy: AIRO, c/o Signora Agnese Martinoli-Bagnasco, Via Serretto 1/4, I-16131 Genova
Netherlands: NSOR, c/o Dr. P. van Laarhoven, c/o McKinsey & Company, Amstel 344, NL-1017 AS Amsterdam, The Netherlands
Norway: NORS, c/o Polyteknisk Forening, Rosenkrantzgt. 7, N-0157 Oslo 1
Poland: Polish Cybernetical Society, Technical University of Poznan, Inst. Automatyki, Piotrowo 3A, PL-60-965 Poznan or ASPORS, IBS Pan, Nowelska 6, PL-01-447 Warszawa
Portugal: APDIO, Cesur-Instituto Superior Tecnico, Av. Rovisco Pais, P-1000 Lisboa
Slovak Republic: SSOR, att. Ivan Brezina, Dept. of Operations Research and Econometrics, Odbojarov 10, 832 20 Bratislava
South Africa: ORSSA, P.O. Box 850, ZA-Groenkloof 0027
Spain: SEIO, Hortaleza 104-2 Izqda, E-28004 Madrid
Sweden: SOAF/SORA, FOA, S-172 90 Sundbyberg
Switzerland: SVOR/ASRO, c/o Ms T. Häring, Grossmattstrasse 38, CH-4133 Pratteln
Turkey: ORST, Department of Industrial Engineering, Middle East Technical University, TR-06531 Ankara
United Kingdom: ORS, Seymour House, 12 Edward Street, Birmingham B1 2RX
Yugoslavia: ETAN, Mihajlo Pupin Institute, P.O. Box 15, Volgina 15, 11000 Beograd

EURO Office and Bulletin

Please submit conference announcements and other news items to the EURO Bulletin (not to EJORI):
Prof. Ph. Van Asbroeck, EURO Office, Université Libre de Bruxelles, Service de mathématiques de la gestion, Bld. du Triomphe CP210/01, B-1050 Brussels, Belgium, Fax: +32 2 650 5970, E-mail: euro@ulb.ac.be
EURO Bulletin is available through the member societies.

EURO WWW pages: <http://www.ulb.ac.be/euro>

Publication information. *European Journal of Operational Research* (ISSN 0377-2217) is sponsored by the Association of European Operational Research Societies within IFORS. For 1999 volumes 112-119 are scheduled for publication. Subscription prices are available upon request from the publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where air delivery via SAL is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, P.R. China, Singapore, South Africa, South Korea, Taiwan, Thailand, U.S.A. For all other countries airmail rates are available upon request. Claims for missing issues must be made within six months of our publication (mailing) date. For orders, claims, product enquiries (no manuscript enquiries) please contact the Customer Support Department at the Regional Sales Office nearest to you:

New York: Elsevier Science, P.O. Box 945, New York, NY 10159-0945, USA. Tel: (+1) 212-633-3730, [Toll Free number for North American Customers: 1-888-4ES-INFO (437-4636)], Fax: (+1) 212-633-3680, E-mail: usinfo-f@elsevier.com
Amsterdam: Elsevier Science, P.O. Box 211, 1000 AE Amsterdam, The Netherlands. Tel: (+31) 20-485-3757, Fax: (+31) 20-485-3432, E-mail: nlinfo-f@elsevier.nl
Tokyo: Elsevier Science K.K., 9-15, Higashi-Azabu 1-chome, Minato-ku, Tokyo 106, Japan. Tel: (+81) 3-5561-5033, Fax: (+81) 3-5561-5047, E-mail: info@elsevier.co.jp
Singapore: Elsevier Science, No. 1 Temasek Avenue, #17-01 Millenia Tower, Singapore 039192. Tel: (+65) 434-3727, Fax (+65) 337-2230, E-mail: asiainfo@elsevier.com.sg
Rio de Janeiro: Elsevier Science, Rua Sete de Setembro 111/16 Andar, 20050-002 Centro, Rio de Janeiro - RJ, Brazil; phone: (+55) (21) 509 5340; fax: (+55) (21) 507 1991; e-mail: elsevier@campus.com.br [Note (Latin America): for orders, claims and help desk information, please contact the Regional Sales Office in New York as listed above]

US MAILING NOTICE - *European Journal of Operational Research* (0377-2217) is published semi-monthly by Elsevier Science B.V. (Molenwerf 1, Postbus 211, 1000 AE Amsterdam). Annual subscription price in the USA US\$ 3249 (US\$ price valid in North, Central and South America only), including air speed delivery. Publications postage paid at Jamaica, NY 11431.

USA POSTMASTERS: Send address changes to: *European Journal of Operational Research*, Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

© The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

Published semimonthly

Printed in the Netherlands