Tilburg University

Health related quality of life among cancer patients in their last year of life

Raijmakers, N.J.H.; Zijlstra, M.; van Roij, J.; Husson, O.; Oerlemans, S.; van de Poll-Franse, L.V.

Published in:
Supportive Care in Cancer

DOI:
[10.1007/s00520-018-4181-6](https://doi.org/10.1007/s00520-018-4181-6)

Publication date:
2018

Document Version
Peer reviewed version

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Raijmakers, N. J. H., Zijlstra, M., van Roij, J., Husson, O., Oerlemans, S., & van de Poll-Franse, L. V. (2018). Health related quality of life among cancer patients in their last year of life: Results from the PROFILES registry. *Supportive Care in Cancer*, 26(10), 3397–3404. <https://doi.org/10.1007/s00520-018-4181-6>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Original Article

Title: Health-related quality of life among cancer patients in their last year of life. Results from the PROFILES registry

Authors

N.J.H. Raijmakers¹, M. Zijlstra^{1,2}, J. van Roij¹, O. Husson^{3,4}, S. Oerlemans¹, L.V. van de Poll-Franse^{1,5,6}

Affiliations:

¹The Netherlands Comprehensive Cancer Organisation, Utrecht, The Netherlands

² Department of Medical Oncology, Radboud MC, Nijmegen, The Netherlands

³ Department of Medical Psychology, Radboud University MC, Nijmegen, The Netherlands

⁴ Institute of Cancer Research and Royal Marsden NHS Foundation Trust, London, UK

⁵ CoRPS - Center of Research on Psychology in Somatic diseases, Department of Medical and Clinical Psychology, Tilburg University, Tilburg, The Netherlands

⁶Division of Psychosocial Research and Epidemiology, The Netherlands Cancer Institute, Amsterdam, The Netherlands

Corresponding author:

Dr. Natasja JH Raijmakers

Netherlands Comprehensive Cancer Organisation (IKNL)

PO box 19079, 3501 DB Utrecht, The Netherlands.

Telephone: +31 088 234 6156

E-mail: n.raijmakers@iknl.nl

Abstract word count 249

Main text word count 3097

ABSTRACT

Purpose

The aim of this study was to assess health-related quality of life (HRQoL) in the last year of life of cancer patients stratified by four periods of time before death.

Patients and methods

Between 2008 and 2015, cancer patients were invited to participate in PROFILES(Patient Reported Outcomes Following Initial treatment and Long term Evaluation of Survivorship) registry studies.

Patients were eligible for inclusion in this secondary analysis if they had been invited to complete the European Organization for Research and Treatment of Cancer Quality of Life Questionnaire(EORTC QLQ-C30) in their last year of life($N=892$). 458 patients (51%) responded. Descriptive statistics were used to describe the HRQoL of cancer patients in the last 3 months of life($N=61$), last 3-6 months($N=110$), last 6-9 months($N=138$), or last 9-12 months of their life($N=129$)

Results

Patients in the last 3 months report a significant lower HRQoL, lower functioning and higher symptom burden of fatigue and appetite loss compared to patients in different time periods before death($p<0.008$). Clinical relevance of the differences for global QoL, cognitive and social functioning were large. Patients' HRQoL in the last year of life was significantly lower than of the normative population ($p<0,001$).

Conclusions

All aspects of HRQoL are considerably impaired in patients with advanced cancer, with a marked lower HRQoL in the final months of life. This marked decline of HRQoL in the final months of life may be an indicator of approaching death and serve as an important trigger for end-of- life communication and decision-making about subsequent treatment and supportive care.

Key words: quality of life, advanced cancer, palliative care, population-based cohort

INTRODUCTION

Treatment of cancer is evolving rapidly and in recent decades cancer survival has improved partly attributed to screening (early detection) and better treatment [1]. Trends in the United States show that five year relative survival in adults with solid cancer has increased from 49% to 68% over the last 40 years [2]. Nevertheless, despite these advances, cancer is still the second leading cause of death worldwide and more than 8 million people die of cancer every year [3]. Ergo, a large number of cancer patients experience a phase of their illness in which they might need palliative care. Palliative care aims to improve or maintain the quality of life (QoL) of patients and their relatives facing problems associated with a life-threatening disease, such as cancer. In their landmark paper, Temel et al showed that early palliative care in fact leads to significant improvements in both QoL and mood [4], as confirmed by a recent meta-analysis [5].

QoL is the perceived quality of an individual's life, that is, an assessment of their well-being and includes multiple domains, including the physical, psychological, social, and spiritual domain. QoL is subjective and dynamic over time. Lynn and Anderson [6] have distinguished three common illness trajectories in patients potentially in need of palliative care, including the cancer trajectory. The 'cancer-trajectory' consists of a short period of marked decline of function and a foreseen death. In line with this trajectory, several studies showed that cancer patients experience a steep decline of function and QoL in the last months of life. Giesinger et al [7] used routinely collected clinical practice data of 85 advanced cancer patients and showed that during the last 3 months of life HRQoL worsened sharply. Hwang et al [8] reported a fast deterioration in the last two months of life of 67 advanced cancer patients admitted to a US tertiary care teaching hospital. Furthermore, Elmqvist et al [9] combined data of two clinical trials from Norway and Sweden and showed that advanced patients' functioning (n=116) deteriorated and the most marked changes occurred in the last 2 months of life.

Insight in the course of advanced cancer patients' quality of life (QoL) during the final year of life will serve to identify goals for timely interventions to improve patients' QoL at the end-of-life. However, there are few population-based studies reporting the self-reported QoL of large groups of cancer patients in their last months of life. Furthermore, previous studies reported on patients admitted to a single hospital who received medical care with palliative intent or on patients with limited expected survival who were included in a clinical trial regarding advanced palliative care, both potential subject

to selection bias. Moreover, clinical trial data does not reflect the daily practice. Population-based information about QoL during the final course of the disease for all cancer patients is lacking. Therefore, this study used data of a large population-based cohort to assess health related quality of life (HRQoL) in the last year of life of advanced cancer patients stratified by four different periods of time before death.

METHODS

Study design and setting

Data from the PROFILES ('Patient Reported Outcomes Following Initial treatment and Long term Evaluation of Survivorship') registry were used for secondary analyses. The PROFILES registry is an ongoing data collection of PROs within the sampling frame of the Netherlands Cancer Registry(NCR) and can be linked with clinical data of all individuals newly diagnosed with cancer in the Netherlands [10].

Study population

The current analysis included patients with cancer between May 2009 and October 2015 who received a questionnaire in their last year of life, using all study samples from the PROFILES registry. In all study samples, participants were included if they were older than 18 years at diagnosis and excluded if they were not able to complete a Dutch questionnaire according to their (ex-)attending specialist (i.e., cognitive impairment, non-native speaker, too ill to participate). Ethical approval was obtained for all study samples separately.

Data collection

A detailed description of the data-collection has been published previously [11]. In brief, in each study sample, cancer patients were informed about the study via a letter by their (ex-)attending specialist. This letter contained either an informed consent and a postal questionnaire, or a secured link to a web-based informed consent and online questionnaire. In study samples where the secured link was provided, the patient could return a postcard to request a paper-and-pencil questionnaire, if preferred. Data from the PROFILES registry are available for non-commercial scientific research, subject to study question, privacy and confidentiality restrictions, and registration[12].

Study measures

Sociodemographic and clinical data

Socio-demographic and clinical data were obtained from the NCR. Socio-demographic variables included date of birth, gender, and socio-economic status (SES). SES was based on postal code of the residence area of the patient, combining aggregated individual fiscal data on the economic value of the home and household incomes, and was categorized into low, medium or high [13]. Questions on educational level and partnership were added to all questionnaire packages. Clinical data include cancer type, stage, and date of diagnosis. Cancer type was classified according to the third International Classification of Diseases for Oncology (ICDO-3) [14] and disease stage was classified according to TNM [15] or Ann Arbor Code (Hodgkin lymphoma and Non-Hodgkin lymphoma). TNM5 was used for patients diagnosed from 2002 to 2003, TNM6 for patients diagnosed from 2003 to 2010, and TNM7 was used for patients diagnosed from 2010. Comorbidity was categorized according to the adapted Self-administered Comorbidity Questionnaire (SCQ). Patients were asked to report comorbid conditions present in the past 12 months. The total score was the sum of all positive responses (range 0-14) and categorized into no comorbid condition, one comorbid condition, and at least two comorbid conditions [16]. Vital status and date of death were obtained from the Dutch municipal personal records database and was last verified on February 1st 2017.

Quality of life

The European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire (EORTC QLQ-C30) (version 3.0) was used to assess HRQoL in cancer patients [17, 18]. It contains five functional scales on physical, role, cognitive, emotional and social functioning, an overall health status/global QoL scale, three symptom scales on fatigue, nausea and vomiting, and pain, and six single items. The scores were linearly transformed into a score between 0 and 100 [19]. A higher score for a functional scale indicates a higher level of functioning, but a higher score for a symptom scale represents a higher level of symptomatology. Furthermore, the recently developed QLQ-C30 summary score has been used, because the sensitivity of the global QoL (ql) in advanced cancer patients seems limited, i.e. it seems not particularly well suited for detecting changes between patient groups [20]. This QLQ-C30 summary score was developed by Giesinger et al [21], with a higher score indicating a better HRQoL.

Normative population

A reference cohort of 2,194 individuals from the general Dutch population (Center panel) was used to obtain the normative population. This reference cohort is representative for the Dutch speaking adult population in the Netherlands [22]. The normative population completed a questionnaire in November 2013 that included the EORTC QLQ-C30, and items on socio-demographics. From this normative population, an age- and gender-matched selection ($N=288$) was made to compare HRQoL with the patient group.

Statistical analysis

Descriptive statistics were used to assess sociodemographic and clinical data and to determine the HRQoL in the last year of life of cancer patients. Respondents and non-respondents were compared using t-test (age) and Chi-square test (gender, partnership, SES). Four groups were created: patients who completed a questionnaire in their last 3 months of life ($N=61$), last 3-6 months ($N=110$), last 6-9 months ($N=138$), and last 9-12 months ($N=129$). Differences between the four groups in HRQoL were analysed with a one-way ANOVA, followed by a Bonferroni post-hoc test. A Bonferroni correction was applied ($P<0.008$) to account for multiple testing. Additionally, clinical relevance of the differences was assessed using the meta-analysis of Cocks et al. [23], who published a guideline to aid interpretation of differences in EORTC QLQ-C30 scores defining thresholds for trivial, small, medium and large differences per subscale. A multivariable regression model was used to determine the associations between the EORTC QoL summary score and moment of completing the questionnaire (time before death in months as continuous variable), adjusted for gender, age, cancer type, and initial treatment.

RESULTS

A total of 892 patients received a questionnaire in their last year of life and 458 patients (51%) completed the questionnaire (mean age 72, standard deviation [SD] 9) (Table 1). Most common diagnoses were colorectal cancer (58%), lymphoma (22%), and gynaecological cancer (12%). Non-respondents ($n=434$) were more often female and were more often in the last 3 months of life compared to respondents ($p<0.05$).

[Table 1]

HRQoL in patients at the end of life

Overall, the mean QLQ-C30 summary score of all patients in their last year of life was 73(SD 19), while in the last 3 months of life the mean QLQ-C30 summary score was 62(SD 22) (Table 2). Most severe symptom burden in the last year of life was found for fatigue (44(SD 30)) and dyspnoea (30(SD 34)), while in the last 3 months fatigue 57(SD 29) and pain 39(SD 35) were most burdensome.

HRQoL trajectory towards the end of life

Significant differences of the QLQ-C30 summary score between different time periods before death were found ($F(3,234) = 9.57$, $p = .000$). HRQoL was statistically significantly lower in the last 3 months of life compared to the last 3-6 months ($p=.0001$), the last 6-9 months ($p= .000$), and the last 9-12 months of life ($p =.000$) (Table 2 and Figure 1). All functioning subscales were significantly lower in the last 3 months of life, compared to patients in the previous time periods ($p<0.008$). Subscales global QoL, cognitive and social functioning showed a large clinically relevant mean lower score in patients in their last 3 months compared to patients in their last 9-12 months, respectively 17, 14 and 18 points. Physical and role functioning showed a medium clinically relevant lower score, respectively 19 and 23 points. Differences of the symptoms fatigue and appetite loss between the last 3 months and the last 9-12 months were also statistically significant. Fatigue, appetite loss, pain, insomnia, dyspnoea and nausea and vomiting showed medium clinically relevant differences between the last 3 months and the last 9-12 months of life (range 11-19 points).

[Table 2]

[Figure 1]

HRQoL steeply declined in the last six months towards death; the QLQ-C30 summary score and the moment of completing the questionnaire (time before death in months) were statistically significant associated ($\beta=2.3$, 95% CI 0.23-4.33, $p=.029$).

Normative population

The QLQ-C30 summary score of advanced cancer patients was significantly lower in patients in their last year compared to the normative population, respectively 73 (SD 19) vs. 87 (SD 13), $p <0.000$

(Table 2). Overall, patients in their last year of life reported a lower functioning and a higher symptom burden on all subscales compared to the normative population.

Discussion

Advanced cancer patients experience a significantly impaired HRQoL in their last year of life, especially in the last three months of life. Patients also experience a high symptom burden, in particular regarding fatigue, dyspnoea and pain. HRQoL of cancer patients in their last year of life is worse compared to the normative population, particularly in the final phase of life.

The marked lower QoL in the last 3 months of life is in line with the theoretical disease trajectory as described by Lynn and Adamson [6] and is in accordance with previous smaller studies [7-9]. A short period of evident decline is typical for cancer, as most patients with malignancies maintain a high level of functioning for a substantial period. However, once the cancer advances, the patient's QoL sharply declines in the final weeks preceding death.

Our analyses demonstrate that QoL measurement using patient-reported outcomes (PROs) is feasible in cancer patients in their last year of life. A response rate in the last year of life of 51% can be considered reasonable compared to the overall response rate of cancer patients in the Profiles registry (73-75%) [24-26]. As expected, we see a lower response rate among patients who participated in their last three months of life (36%). Completing a questionnaire in the final phase of life is obviously more difficult, possibly due to deterioration and higher symptom burden (as shown in this study). Using PROs in (early and late) palliative oncological care is important, as it provides valuable information about the QoL that would support end-of-life decision-making about subsequent treatment and supportive care. Furthermore, monitoring QoL and symptoms increases awareness among health care professionals to better anticipate on patients' changing needs [27, 28] and improves clinical outcomes (i.e. fewer ER visits, fewer hospitalizations, and better survival) [28].

Strengths and limitations

An important strength of the current analysis is the large population-based sample of cancer patients in their last year of life, including different primary cancer sites. Another strength is that, through linkage with cancer registry data and the Dutch municipal personal records database, we had access to complete and comprehensive data on socio-demographic and clinical characteristics, for the full

population of respondents and non-respondents. Furthermore, in our analysis we have used the EORTC QLQ-C30, a widely used instrument to measure HRQoL within oncology. However, many instruments to assess HRQoL of patients with advanced cancer are available [29]. The EORTC QLQ-C30 seems suitable for patients with advanced cancer, although for patients in their final weeks of life the shortened version of this questionnaire, the EORTC QLQ-C15-PAL might be more appropriate [30]. A limitation of our study is its design, a cross-sectional analyses, based on a collection of separate study samples, with different inclusion criteria.

Practical implications

Our results clearly show a progressive deterioration in QoL towards the end of life. This marked decline of QoL may be an indicator of approaching death and therefore should be an important trigger for end-of-life communication and decision-making about subsequent treatment and care. Ideally, this should start earlier. However, timing of these end-of-life discussions remains challenging, “it always seems too early, until it’s too late”. Therefore, in current practice, a change in QoL or symptom burden can serve as a welcome starting point for these discussions to help professional caregivers to overcome the experienced barriers [31, 32]. The routine assessment of patient-reported outcomes (PROs) in advanced cancer patients helps to provide information on QoL and symptom burden and are widely recommended for clinical oncology practice [33] and for palliative care [34] .

Conclusion

Cancer patients experience a significantly impaired QoL and high symptom burden in their last year of life, especially in the last three months of life. This considerable decline of function and increase of symptom burden in the final months of life might serve as an indicator for end-of-life communication and supportive care.

Acknowledgements

The PROFILES registry was funded by an Investment Grant (#480-08-009) of the Netherlands Organization for Scientific Research (The Hague, The Netherlands). Dr. Olga Husson is supported by a Social Psychology Fellowship from the Dutch Cancer Society (#KUN2015-7527). These funding agencies had no further role in conducting the study.

Disclosure

The authors have declared no conflicts of interest.

References

1. Torre LA, Siegel RL, Ward EM, Jemal A. Global Cancer Incidence and Mortality Rates and Trends--An Update. *Cancer Epidemiol Biomarkers Prev* 2016; 25: 16-27.
2. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin* 2015; 65: 5-29.
3. <http://www.who.int/mediacentre/factsheets/fs297/en/> Last accessed on November 2017.
4. Temel JS, Greer JA, Muzikansky A et al. Early palliative care for patients with metastatic non-small-cell lung cancer. *N Engl J Med* 2010; 363: 733-742.
5. Kavalieratos D, Corbelli J, Zhang D et al. Association Between Palliative Care and Patient and Caregiver Outcomes: A Systematic Review and Meta-analysis. *JAMA* 2016; 316: 2104-2114.
6. Lynn J and Adamson DM. Living well at the end of life. Adapting health care to serious chronic illness in old age. RAND Health: Santa Monica. 2003.
7. Giesinger JM, Wintner LM, Oberguggenberger AS et al. Quality of life trajectory in patients with advanced cancer during the last year of life. 2011; 14: 904-912.
8. Hwang SS, Chang VT, Fairclough DL et al. Longitudinal quality of life in advanced cancer patients: pilot study results from a VA medical cancer center. 2003; 25: 225-235.
9. Elmqvist MA, Jordhoy MS, Bjordal K et al. Health-related quality of life during the last three months of life in patients with advanced cancer. *Support Care Cancer* 2009; 17: 191-198.
10. www.cijfersoverkanker.nl NCR: Netherlands Cancer Registration. Last accessed on November 2017.
11. van de Poll-Franse LV, Horevoorts N, van Eenbergen M et al. The Patient Reported Outcomes Following Initial treatment and Long term Evaluation of Survivorship registry: scope, rationale and design of an infrastructure for the study of physical and psychosocial outcomes in cancer survivorship cohorts. *Eur J Cancer* 2011; 47: 2188-2194.
12. www.profilesregistry.nl The PROFILES Registry. Last accessed on November 2017.
13. van Duijn C and Keij I. Sociaal-economische status indicator op postcode niveau. *Maandstatistiek van de bevolking* 2002; 50: 32-35. <https://www.cbs.nl/-/media/imported/documents/2002/08/b-15-02-02.pdf>
14. Fritz A PC, Jack A, Shanmugaratnam K, Sobin L, Parkin DM, Whelan S. International Classification of Diseases for Oncology First Revision. WHO Library Cataloguing-in-Publication Data 2013.
15. Sobin LH, Fleming ID. TNM Classification of Malignant Tumors, fifth edition (1997). Union Internationale Contre le Cancer and the American Joint Committee on Cancer. *Cancer* 1997; 80: 1803-1804.
16. Sangha O, Stucki G, Liang MH et al. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum* 2003; 49: 156-163.
17. Niezgoda HE, Pater JL. A validation study of the domains of the core EORTC quality of life questionnaire. *Qual Life Res* 1993; 2: 319-325.
18. Aaronson NK, Ahmedzai S, Bergman B et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993; 85: 365-376.
19. Fayers PM AN, Bjordal K, Groenvold M, Curran D, Bottomley A, on behalf of the EORTC Quality of Life Group. The EORTC QLQ-C30 Scoring Manual (3rd Edition). In. Brussels: 2001.
20. Nordin K, Steel J, Hoffman K, Glimelius B. Alternative methods of interpreting quality of life data in advanced gastrointestinal cancer patients. *Br J Cancer* 2001; 85: 1265-1272.
21. Giesinger JM, Kieffer JM, Fayers PM et al. Replication and validation of higher order models demonstrated that a summary score for the EORTC QLQ-C30 is robust. 2016; 69: 79-88.
22. www.centerdata.nl/en Centerpanel. Last accessed on November 2017
23. Cocks K, King MT, Velikova G et al. Evidence-based guidelines for determination of sample size and interpretation of the European Organisation for the Research and Treatment of Cancer Quality of Life Questionnaire Core 30. *J Clin Oncol* 2011; 29: 89-96.
24. Husson O, Thong MS, Mols F et al. Information provision and patient reported outcomes in patients with metastasized colorectal cancer: results from the PROFILES registry. *J Palliat Med* 2013; 16: 281-288.
25. Horevoorts NJ, Vissers PA, Mols F et al. Response rates for patient-reported outcomes using web-based versus paper questionnaires: comparison of two invitational methods in older colorectal cancer patients. *J Med Internet Res* 2015; 17: e111.
26. Nicolaije KA, Ezendam NP, Pijnenborg JM et al. Paper-Based Survivorship Care Plans May be Less Helpful for Cancer Patients Who Search for Disease-Related Information on the Internet: Results of the Registrationsystem Oncological Gynecology (ROGY) Care Randomized Trial. *J Med Internet Res* 2016; 18: e162.
27. Etkind SN, Daveson BA, Kwok W et al. Capture, transfer, and feedback of patient-centered outcomes data in palliative care populations: does it make a difference? A systematic review. *J Pain Symptom Manage* 2015; 49: 611-624.

28. Basch E, Deal AM, Kris MG et al. Symptom Monitoring With Patient-Reported Outcomes During Routine Cancer Treatment: A Randomized Controlled Trial. *J Clin Oncol* 2016; 34: 557-565.
29. van Roij J FH, Van de Poll-Franse L, Zijlstra M, Raijmakers NJH. Measuring Quality of life in patient with advanced cancer: A systematic review of self-administered instruments. *under review*
30. Groenvold M, Petersen MA, Aaronson NK et al. The development of the EORTC QLQ-C15-PAL: a shortened questionnaire for cancer patients in palliative care. *Eur J Cancer* 2006; 42: 55-64.
31. Graneek L, Krzyzanowska MK, Tozer R, Mazzotta P. Oncologists' strategies and barriers to effective communication about the end of life. *J Oncol Pract* 2013; 9: e129-135.
32. Baile WF, Lenzi R, Parker PA et al. Oncologists' attitudes toward and practices in giving bad news: an exploratory study. *J Clin Oncol* 2002; 20: 2189-2196.
33. Wintner LM, Sztankay M, Aaronson N et al. The use of EORTC measures in daily clinical practice-A synopsis of a newly developed manual. *Eur J Cancer* 2016; 68: 73-81.
34. Stover AM, Basch EM. The Potential Role of Symptom Questionnaires in Palliative and Supportive Cancer Care Delivery. *Curr Oncol Rep* 2017; 19: 12.

Table captions

Table 1 Sociodemographic and clinical characteristics

Table 2 Health related quality of life of cancer patients in their last months of life (n=458)

Figure 1 Differences in HRQoL and its subscales in cancer patients in their last year of life (n=458)

Table 1 Sociodemographic and clinical characteristics

	Respondents (N=458)	Non respondents (N=434)	p-value	Normative population ^a (N=288)
Age (mean(SD); range min-max)	72(9); 40-96	73(11); 21-99	0.1985	71(11); 24-90
Gender (% male)	59%	49%	0.005*	54%
Cancer type			0.000*	n.a.
Colon/rectum	58% (n=264)	48% (n=208)		
Lymphoma	22% (n=100)	15% (n=66)		
Gynaecological	12% (n=54)	20% (n=88)		
Prostate	6.1%(n=28)	9.5% (n=41)		
Other	2.6% (n=12)	7.1% (n=31)		
Metastasis at diagnosis			0.254	n.a.
Yes	25% (n=114)	22% (n=94)		
No	75% (n=344)	78% (n=340)		
Time since diagnosis (yrs.)				
(mean(SD))	3.6(2.6)	3.6(2.8)	0.7915	n.a.
Up to two years	27% (n=125)	33% (n=141)		
2 – 4 years	40% (n=185)	34% (n=147)		
More than 4 years	32% (n=148)	34% (n=146)		
Moment of receiving questionnaire			0.000*	n.a.
Last 3 months of life	14% (n=65)	27% (n=118)		
3-12 months before death	86% (n=393)	73% (n=316)		
Comorbidity				
No comorbid condition	30% (n=126)	-		23% (n=66)
One comorbid condition	24% (n=110)	-		26% (n=76)
More than one comorbid conditions	48% (n=222)	-		51% (n=146)
Most frequent conditions				
Hypertension	34% (n=138)	-		37% (n=107)
Back pain	30% (n=116)	-		35% (n=101)
Arthritis	29% (n=113)	-		33% (n=96)
Heart disease	27% (n=109)	-		24% (n=69)
Diabetes mellitus	20% (n=80)	-		11% (n=32)
Pulmonary disease	17% (n=66)	-		12% (n=35)
Partnership^b				
Yes	74% (n=338)	-		69%
No	24% (n=110)	-		31%

Social economic status^b

High	23.6% (n=102)	23% (n=100)	-
Intermediate	41.6% (n=180)	41.5% (n=180)	-
Low	34.9% (n=151)	27.4% (n=119)	-

Due to rounding off, percentages can exceed 100% and an asterisk indicated statistically significance at $p < 0.01$

^a Matched normative population on age and gender

^b Due to missings it does not add up to 458 (missings did not exceed 5%)

						Clinical	
	I: Last 3	II: Last 3-6	III: Last 6-9	IV: Last 9-12		relevance	Normative
	months of	months of	months of	months of		of mean	population
	life	life	life	life		difference	
						between I	
						and IV ²	
	N =65	N =118	N =142	N =133			N =288
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	P-value ¹		Mean (SD)
Summary score QoL	62 (22)	73 (19)	75 (18)	77 (17)	0.0000*		87 (13)
Quality of life							
Physical functioning	48 (28)	63 (24)	62 (25)	67 (25)	0.0000*	medium	83 (19)
Role functioning	39 (37)	59 (35)	57 (34)	62 (32)	0.0001*	medium	83 (25)
Emotional functioning	64 (26)	75 (25)	78 (23)	83 (20)	0.0000*	-	87 (18)
Cognitive functioning	68 (28)	80 (29)	76 (28)	82 (20)	0.0004*	large	88 (17)
Social functioning	60 (29)	74 (31)	72 (30)	78 (24)	0.0007*	large	92 (18)
Global quality of life	50 (27)	61 (24)	60 (24)	67 (20)	0.0000*	large	75 (19)
Symptoms							
Fatigue	57 (29)	44 (30)	42 (29)	38 (28)	0.0006*	medium	23 (24)
Nausea/vomiting	18 (24)	15 (25)	11 (22)	9.8 (20)	0.0536	medium	3 (11)
Pain	39 (35)	28 (29)	27 (32)	25 (29)	0.0322	medium	22 (27)
Dyspnoea	37 (38)	33 (34)	28 (34)	26 (32)	0.0758	medium	12 (22)
Insomnia	37 (36)	30 (32)	26 (31)	22 (31)	0.0214	medium	19 (26)
Appetite loss	36 (34)	25 (35)	18 (27)	17 (30)	0.0002*	medium	5 (16)
Constipation	21 (31)	12 (25)	14 (24)	12 (24)	0.0893	small	9 (18)
Diarrhoea	22 (33)	11 (20)	13 (24)	19 (28)	0.0081	small	6 (14)
¹ A one-way ANOVA was conducted to determine if QoL differed for patients in different number of months before death, followed by post-hoc Bonferroni test (not shown).							
² Indication of clinical relevance of mean differences, as reported by Cocks et al, 2010							
*statistically significant; a Bonferroni correction was applied (P<0.008)) to account for multiple testing.							

Figure 1 Differences in HRQoL and its subscales in cancer patients in their last year of life (n=458)