

Tilburg University

Interlacing eigenvalues and graphs

Haemers, W.H.

Publication date:
1994

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Haemers, W. H. (1994). *Interlacing eigenvalues and graphs*. (Research memorandum / Tilburg University, Department of Economics; Vol. FEW 675). Unknown Publisher.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Interlacing Eigenvalues and Graphs

by

Willem H. Haemers

Tilburg University, Department of Econometrics,
P.O.Box 90153, 5000 LE Tilburg, The Netherlands.
e-mail: haemers@kub.nl

Dedicated to J.J. Seidel⁶⁵²⁸¹

Abstract

We give several old and some new applications of eigenvalue interlacing to matrices associated to graphs. Bounds are obtained for characteristic numbers of graphs, such as the size of a maximal (co)clique, the chromatic number, the diameter and the bandwidth in terms of the eigenvalues of the standard adjacency matrix or the Laplacian matrix. We also deal with inequalities and regularity results concerning the structure of graphs and block designs.

1 Preface

Between 1975 and 1979, under the inspiring supervision of J.J. Seidel, I did my PhD research on applications of eigenvalue techniques to combinatorial structures. It turned out that eigenvalue interlacing provides a handy tool for obtaining inequalities and regularity results concerning the structure of graphs in terms of eigenvalues of the adjacency matrix. After fifteen years my thesis [14] became an obscure reference (I myself have no spare copies left) and in addition I came across some new applications. This made me decide to write the present paper, which is an attempt to survey the various kinds of applications of eigenvalue interlacing and I am very glad to have the opportunity to present it in this issue of Linear Algebra and Applications, dedicated to the person who educated me in combinatorial matrix theory and made me a believer in the power of eigenvalue techniques.

⁶⁵²⁸¹This paper is meant to be published in a special issue of Linear Algebra and its Applications in the honour of J.J. Seidel

2 Interlacing

Consider two sequences of real numbers: $\lambda_1 \geq \dots \geq \lambda_n$, and $\mu_1 \geq \dots \geq \mu_m$ with $m < n$. The second sequence is said to *interlace* the first one whenever

$$\lambda_i \geq \mu_i \geq \lambda_{n-m+i}, \quad \text{for } i = 1, \dots, m .$$

The interlacing is called *tight* if there exist an integer $k \in [0, m]$ such that

$$\lambda_i = \mu_i \quad \text{for } 0 \leq i \leq k \quad \text{and} \quad \lambda_{n-m+i} = \mu_i \quad \text{for } k+1 \leq i \leq m .$$

If $m = n - 1$ the interlacing inequalities become $\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \dots \geq \mu_m \geq \lambda_n$, which clarifies the name. Godsil [13] reserves the name interlacing for this particular case and calls it generalized interlacing otherwise. Throughout, the λ_i 's and μ_i 's will be eigenvalues of matrices A and B , respectively. Basic to eigenvalue interlacing is Rayleigh's principle, a standard (and easy to prove) result from linear algebra, which can be stated as follows. Let u_1, \dots, u_n be an orthonormal set of eigenvectors of the symmetric matrix A , such that u_i is a λ_i -eigenvector (we use this abbreviation for an eigenvector corresponding to the eigenvalue λ_i). Then

$$\frac{u^\top A u}{u^\top u} \geq \lambda_i \quad \text{if } u \in \langle u_1, \dots, u_i \rangle \quad \text{and} \quad \frac{u^\top A u}{u^\top u} \leq \lambda_i \quad \text{if } u \in \langle u_1, \dots, u_{i-1} \rangle^\perp .$$

In both cases equality implies that u is a λ_i -eigenvector of A .

Theorem 2.1 *Let S be a real $n \times m$ matrix such that $S^\top S = I$ and let A be a symmetric $n \times n$ matrix with eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$. Define $B = S^\top A S$ and let B have eigenvalues $\mu_1 \geq \dots \geq \mu_m$ and respective eigenvectors v_1, \dots, v_m .*

- i. *The eigenvalues of B interlace those of A .*
- ii. *If $\mu_i = \lambda_i$ or $\mu_i = \lambda_{n-m+i}$ for some $i \in [1, m]$, then B has a μ_i -eigenvector v such that Sv is a μ_i -eigenvector of A .*
- iii. *If for some integer ℓ , $\mu_i = \lambda_i$ for $i = 1, \dots, \ell$ (or $\mu_i = \lambda_{n-m+i}$ for $i = \ell, \dots, m$), then Sv_i is a μ_i -eigenvector of A for $i = 1, \dots, \ell$ (respectively $i = \ell, \dots, m$).*
- iv. *If the interlacing is tight then $SB = AS$.*

Proof. With u_1, \dots, u_n as above, for each $i \in [1, m]$, take a non-zero vector s_i in

$$\langle v_1, \dots, v_i \rangle \cap \langle S^\top u_1, \dots, S^\top u_{i-1} \rangle^\perp . \tag{1}$$

Then $Ss_i \in \langle u_1, \dots, u_{n-1} \rangle^\perp$, hence by Rayleigh's principle

$$\lambda_i \geq \frac{(Ss_i)^\top A (Ss_i)}{(Ss_i)^\top (Ss_i)} = \frac{s_i^\top B s_i}{s_i^\top s_i} \geq \mu_i ,$$

and similarly (or by applying the above inequality to $-A$ and $-B$) we get $\lambda_{n-m+i} \leq \mu_i$, proving (i).

If $\lambda_i = \mu_i$, then s_i and Ss_i are λ_i -eigenvectors of B and A respectively, proving (ii).

We prove (iii) by induction on ℓ . Assume $Sv_i = u_i$ for $i = 1, \dots, \ell - 1$, then we may take $s_\ell = v_\ell$ in (1), but in proving (ii) we saw that Ss_ℓ is a λ_ℓ -eigenvector of A . (The statement between parenthesis follows by considering $-A$ and $-B$.) Thus we have (iii). Let the interlacing be tight, then by (iii) Sv_1, \dots, Sv_m is an orthonormal set of eigenvectors of A for the eigenvalues μ_1, \dots, μ_m . So we have $SBv_i = \mu Sv_i = ASv_i$ for $i = 1, \dots, m$. Since the vectors v_i form a basis, it follows that $SB = AS$. \square

If we take $S = [I \ O]^\top$, then B is just a principal submatrix of A and we have:

Corollary 2.2 *If B is a principal submatrix of a symmetric matrix A , then the eigenvalues of B interlace the eigenvalues of A .*

Suppose rows and columns of

$$A = \begin{bmatrix} A_{1,1} & \cdots & A_{1,m} \\ \vdots & & \vdots \\ A_{m,1} & \cdots & A_{m,m} \end{bmatrix}$$

are partitioned according to a partitioning X_1, \dots, X_m of $\{1, \dots, n\}$ with characteristic matrix \tilde{S} (that is, $(\tilde{S})_{i,j} = 1$ if $i \in X_j$ and 0 otherwise). The *quotient matrix* is the matrix \tilde{B} whose entries are the average row sums of the blocks of A , more precisely:

$$(\tilde{B})_{i,j} = \frac{1}{|X_i|} \mathbf{1}^\top A_{i,j} \mathbf{1} = \frac{1}{|X_i|} (\tilde{S}^\top A \tilde{S})_{i,j}$$

($\mathbf{1}$ denotes the all-one vector). The partition is called *regular* (or *equitable*) if each block $A_{i,j}$ of A has constant row (and column) sum, that is $A\tilde{S} = \tilde{S}\tilde{B}$.

Corollary 2.3 *Suppose \tilde{B} is the quotient matrix of a symmetric partitioned matrix A .*

- i.** *The eigenvalues of \tilde{B} interlace the eigenvalues of A .*
- ii.** *If the interlacing is tight then the partition is regular.*

Proof. Put $D = \text{diag}(|X_1|, \dots, |X_m|)$, $S = \tilde{S}D^{-\frac{1}{2}}$. Then the eigenvalues of $B = S^\top AS$ interlace those of A . This proves (i), because B and $\tilde{B} = D^{-\frac{1}{2}}BD^{\frac{1}{2}}$ have the same spectrum. If the interlacing is tight then $SB = AS$, hence $A\tilde{S} = \tilde{S}\tilde{B}$. \square

Theorem 2.1(i) is a classical result, see Courant and Hilbert [6] (Vol.1, Ch.I). For the special case of a principal submatrix (Corollary 2.2) the result even goes back to Cauchy and is therefore often referred to as Cauchy interlacing. Interlacing for the quotient

matrix (Corollary 2.3) is especially applicable to combinatorial structures (as we shall see). Payne (see for instance [28]) has applied the extremal inequalities $\lambda_1 \geq \mu_i \geq \lambda_n$ to finite geometric structures several times. He contributes the method to Higman and Sims and therefore calls it the "Higman-Sims technique".

3 Graphs and subgraphs

Throughout the paper G is a graph on n vertices (undirected, simple and loopless) having an adjacency matrix A with eigenvalues $\lambda_1 \geq \dots \geq \lambda_n$. The size of the largest coclique (independent set of vertices) of G is denoted by $\alpha(G)$. Both Corollaries 2.2 and 2.3 lead to a bound for $\alpha(G)$.

Theorem 3.1 $\alpha(G) \leq |\{i | \lambda_i \geq 0\}|$ and $\alpha(G) \leq |\{i | \lambda_i \leq 0\}|$.

Proof. A has a principal submatrix $B = O$ of size $\alpha(G)$. Corollary 2.2 gives $\lambda_\alpha \geq \mu_\alpha = 0$ and $\lambda_{n-\alpha+1} \leq \mu_1 = 0$. \square

Theorem 3.2 If G is regular then $\alpha(G) \leq n \frac{-\lambda_n}{\lambda_1 - \lambda_n}$ and if a coclique C meets this bound then every vertex not in C is adjacent to precisely $-\lambda_n$ vertices of C .

Proof. We apply corollary 2.3. Let $k = \lambda_1$ be the degree of G and put $\alpha = \alpha(G)$. The coclique gives rise to a partition of A with quotient matrix

$$B = \begin{bmatrix} 0 & k \\ \frac{k\alpha}{n-\alpha} & k - \frac{k\alpha}{n-\alpha} \end{bmatrix}.$$

B has eigenvalues $\mu_1 = k$ (row sum) and $\mu_2 = -\frac{k\alpha}{n-\alpha}$ ($\text{trace}(B) - k$) and so $\lambda_n \leq \mu_2$ gives the required inequality. If equality holds then $\mu_2 = \lambda_n$ and since $\mu_1 = \lambda_1$ the interlacing is tight and hence the partition is regular. \square

The first bound is due to Cvetković [7]. The second bound is an unpublished result of Hoffman. There are many examples where equality holds. For instance, a 4-coclique in the Petersen graph is tight for both bounds. The second bound can be generalized to arbitrary graphs in the following way:

Theorem 3.3 If G has smallest degree δ then $\alpha(G) \leq n \frac{-\lambda_1 \lambda_n}{\delta^2 - \lambda_1 \lambda_n}$.

Proof. Now we let k denote the average degree of the vertices of the coclique. Then the quotient matrix B is the same as above, except maybe for the entry $(B)_{2,2}$. Interlacing gives

$$-\lambda_1 \lambda_n \geq -\mu_1 \mu_2 = -\det(B) = \frac{k^2 \alpha}{n - \alpha} \geq \frac{\delta^2 \alpha}{n - \alpha},$$

which yields the required inequality. \square

If G is regular of degree k then $\delta = \lambda_1 = k$ and Theorem 3.3 reduces to Hoffman's bound (3.2). Lovász [23] proved that Hoffman's bound is also an upper bound for the Shannon capacity of G . This a concept from information theory defined as follows. Denote by G^ℓ the product of ℓ copies of G (that is the graph with vertex set $\{1, \dots, n\}^\ell$, where two vertices are adjacent if all of the coordinate places correspond to adjacent or coinciding vertices of G ; if we denote the Kronecker product of ℓ copies of a matrix M by $M^{\otimes \ell}$, then the adjacency matrix of G^ℓ is given by $(A + I)^{\otimes \ell} - I$). The number

$$\Theta(G) = \sup_{\ell} \sqrt[\ell]{\alpha(G^\ell)}$$

is called the Shannon capacity of G . Clearly $\Theta(G) \geq \alpha(G)$, so Lovász' bound implies Hoffman's bound. But conversely, Lovász' bound can be proved using 3.2.

Theorem 3.4 *Let G be regular of degree k , then $\Theta(G) \leq n \frac{-\lambda_n}{k - \lambda_n}$.*

Proof. First note that the above proof of 3.2 remains valid if the ones in A are replaced by arbitrary real numbers, as long as A remains symmetric with constant row sum. So we may apply Hoffman's bound to $A_\ell = (A - \lambda_n I)^{\otimes \ell} - (-\lambda_n)^\ell I$ to get a bound for $\alpha(G^\ell)$. It easily follows that A_ℓ has row sum $(k - \lambda_n)^\ell - (-\lambda_n)^\ell$ and smallest eigenvalue $-(-\lambda_n)^\ell$. So we find $\alpha(G^\ell) \leq (n \frac{-\lambda_n}{k - \lambda_n})^\ell$. \square

For the pentagon C_5 we get $\Theta(C_5) \leq \sqrt{5}$. This is sharp since C_5^2 has a coclique of size 5. More generally, one can obtain results on the size of induced subgraphs, analogues to Hoffman's bound.

Theorem 3.5 *Let G be regular of degree k and suppose G has an induced subgraph G' with n' vertices and m' edges. Then*

$$\lambda_2 \geq \frac{2m' \frac{n}{n'} - n'k}{n - n'} \geq \lambda_n .$$

If equality holds on either side then G' is regular and so is the subgraph induced by the vertices not in G' .

Proof. We now have a quotient matrix

$$B = \begin{bmatrix} \frac{2m'}{n'} & k - \frac{2m'}{n'} \\ \frac{n'k - 2m'}{n - n'} & k - \frac{n'k - 2m'}{n - n'} \end{bmatrix} ,$$

with eigenvalues k and $\frac{2m'}{n'} - \frac{n'k-2m'}{n-n'}$ and Theorem 2.3 gives the result. \square

If $m' = 0$ we get Hoffman's bound back. If $m' = \frac{1}{2}n'(n' - 1)$ Theorem 3.5 gives that the size of a clique is bounded above by

$$n \frac{1 + \lambda_2}{n - k + \lambda_2} ,$$

which is again Hoffman's bound applied to the complement of G . Like in Theorem 3.3, also the above result can be generalized to non-regular graphs. Better bounds can sometimes be obtained if more is known about the structure of G' by considering a refinement of the partition. This is for example the case if G' is a bipartite. See [14] for details.

4 Chromatic number

A colouring of a graph G is a partition of its vertices into cocliques (colour classes). Therefore the number of colour classes, and hence the chromatic number $\chi(G)$ of G is bounded below by $\frac{n}{\alpha(G)}$. Thus upper bounds for $\alpha(G)$ give lower bounds for $\chi(G)$. For instance if G is regular Theorem 3.2 implies that $\chi(G) \geq 1 - \frac{\lambda_1}{\lambda_n}$. This bound remains however valid for non-regular graphs. (But note that it does not follow from 3.3.)

Theorem 4.1

i. If G is not the empty graph then $\chi(G) \geq 1 - \frac{\lambda_1}{\lambda_n}$.

ii. If $\lambda_2 > 0$ then $\chi(G) \geq 1 - \frac{\lambda_{n-\chi(G)+1}}{\lambda_2}$.

Proof. Let X_1, \dots, X_χ ($\chi = \chi(G)$) denote the colour classes of G and let u_1, \dots, u_n be an orthonormal set of eigenvectors of A (where u_i corresponds to λ_i). For $i = 1, \dots, \chi$ let s_i denote the restriction of u_1 to X_i , that is

$$(s_i)_j = \begin{cases} (u_1)_j & \text{if } j \in X_i , \\ 0 & \text{otherwise ,} \end{cases}$$

and put $\tilde{S} = [s_1 \ \dots \ s_\chi]$ (if some $s_i = 0$ we delete it from \tilde{S} and proceed similarly) and $D = \tilde{S}^\top \tilde{S}$, $S = \tilde{S} D^{-\frac{1}{2}}$ and $B = S^\top A S$. Then B has zero diagonal (since each colour class corresponds to a zero submatrix of A) and an eigenvalue λ_1 ($d = D^{\frac{1}{2}} \mathbf{1}$ is a λ_1 -eigenvector of B). Moreover interlacing theorem 2.1 gives that the remaining eigenvalues of B are at least λ_n . Hence

$$0 = \text{trace}(B) = \mu_1 + \dots + \mu_\chi \geq \lambda_1 + (\chi - 1)\lambda_n ,$$

which proves (i), since $\lambda_n < 0$. The proof of (ii) is similar, but a bit more complicated. With s_1, \dots, s_χ as above, choose a non-zero vector s in

$$\langle u_{n-\chi+1}, \dots, u_n \rangle \cap \langle s_1, \dots, s_\chi \rangle^\perp .$$

The two spaces have non-trivial intersection since the dimensions add up to n and u_1 is orthogonal to both. Redefine s_i to be the restriction of s to X_i , and let \tilde{S}, D, S , and d be analogous to above. Put $A' = A - (\lambda_1 - \lambda_2)u_1u_1^\top$, then the largest eigenvalue of A' equals λ_2 , but all other eigenvalues of A are also eigenvalues of A' with the same eigenvectors. Define $B = S^\top A' S$. Now B has again zero diagonal (since $u_1^\top S = 0$). Moreover, B has smallest eigenvalue $\mu_\chi \leq \lambda_{n-\chi+1}$, because

$$\mu_\chi \leq \frac{d^\top B d}{d^\top d} = \frac{s^\top A' s}{s^\top s} \leq \lambda_{n-\chi+1} .$$

So interlacing gives

$$0 = \text{trace}(B) = \mu_1 + \dots + \mu_\chi \leq \lambda_{n-\chi+1} + (\chi - 1)\lambda_2 .$$

Since $\lambda_2 > 0$, (ii) follows. □

The first inequality is due to Hoffman [19]. The proof given here seems to be due to the author [15] and is a customary illustration of interlacing, see for example Lovász [24] (problem 11.21) or Godsil [13] (p.84). In [14] more inequalities of the above kind are given. But only the two treated here turned out to be useful. The condition $\lambda_2 > 0$ is not strong; only the complete multipartite graphs, possibly extended with some isolated vertices have $\lambda_2 \leq 0$. The second inequality looks a bit awkward, but can be made more explicit if the smallest eigenvalue λ_n has large multiplicity m_n , say. Then (ii) yields $\chi \geq \min\{1 + m_n, 1 - \frac{\lambda_n}{\lambda_2}\}$ (indeed, if $\chi \leq m_n$, then $\lambda_n = \lambda_{n-\chi+1}$, hence $\chi \geq 1 - \frac{\lambda_n}{\lambda_2}$). For strongly regular graphs with $\lambda_2 > 0$ it is shown in [14], by use of Seidel's absolute bound (see Delsarte Goethals and Seidel [11]), that the minimum is always taken by $1 - \frac{\lambda_n}{\lambda_2}$, except for the pentagon (see Section 7 for more about strongly regular graphs). So we have

Corollary 4.2 *If G is a strongly regular graph, not the pentagon or a complete multipartite graph, then*

$$\chi(G) \geq 1 - \frac{\lambda_n}{\lambda_2} .$$

For example if G is the Kneser graph $K(m, 2)$ (i.e. the complement of the line graph of K_m) then G is strongly regular with eigenvalues $\lambda_1 = \frac{1}{2}(m-2)(m-3)$, $\lambda_2 = 1$ and $\lambda_n = 3 - m$ (for $m \geq 4$). The above bound gives $\alpha(G) \geq m - 2$, which is tight, whilst Hoffman's lower bound (4.1(i)) equals $\frac{1}{2}m$. On the other hand, if m is even, Hoffman's bound is tight for the complement of G whilst the above bound is much less.

5 Designs

In case we have a non-symmetric matrix N (say) or different partitions for rows and columns, we can still use interlacing by considering the matrix

$$A = \begin{bmatrix} O & N \\ N^\top & O \end{bmatrix}.$$

Then we find results in terms of the eigenvalues of A , which now satisfy $\lambda_i = \lambda_{n-i+1}$ for $i = 1, \dots, n$. The positive eigenvalues of A are the singular values of N , they are also the square roots of the non-zero eigenvalues of NN^\top (and of $N^\top N$). In particular if N is the $(0,1)$ incidence matrix of some design or incidence structure D , we consider the bipartite incidence graph G . An edge of G corresponds to a flag (an incident point-block pair) of D and D is a 1 - (v, k, r) design precisely when G is biregular with degrees k and r .

Theorem 5.1 *Let D be a 1 - (v, k, r) design with b blocks and let D' be a substructure with v' points, b' blocks and m' flags, then*

$$(m' \frac{v}{v'} - b'k)(m' \frac{b}{b'} - v'r) \leq \lambda_2^2(v - v')(b - b').$$

Equality implies that all four substructures induced by the point set of D' or its complement, and the block set of D' or its complement form a 1 -design (possibly degenerate).

Proof. We apply Corollary 2.3. The substructure D' gives rise to a partition of A with the following quotient matrix

$$B = \begin{bmatrix} 0 & 0 & \frac{m'}{v'} & r - \frac{m'}{v'} \\ 0 & 0 & \frac{b'k - m'}{v - v'} & r - \frac{b'k - m'}{v - v'} \\ \frac{m'}{b'} & k - \frac{m'}{b'} & 0 & 0 \\ \frac{v'r - m'}{b - b'} & k - \frac{v'r - m'}{b - b'} & 0 & 0 \end{bmatrix}.$$

We easily have $\lambda_1 = -\lambda_n = \mu_1 = -\mu_4 = \sqrt{rk}$ and $\det(B) = rk \left(\frac{m' \frac{v}{v'} - b'k}{v - v'} \right) \left(\frac{m' \frac{b}{b'} - v'r}{b - b'} \right)$.

Interlacing gives

$$\frac{\det(B)}{rk} = -\mu_2\mu_3 \leq -\lambda_2\lambda_{n-1} = \lambda_2^2,$$

which proves the first statement. If equality holds then $\lambda_1 = \mu_1$, $\lambda_2 = \mu_2$, $\lambda_{n-1} = \mu_3$ and $\lambda_n = \mu_4$, so we have tight interlacing, which implies the second statement. \square

The above result becomes especially useful if we can express λ_2 in terms of the design parameters. For instance if D is a 2 - (v, k, λ) design, then $\lambda_2^2 = r - \lambda = \lambda \frac{v-k}{k-1}$ (see for

example Hughes and Piper [20]) and if D is a generalized quadrangle of order (s, t) , then $\lambda_2^2 = s + t$ (see for instance Payne and Thas [29]). Let us consider two special cases.

Corollary 5.2 *If a symmetric 2- (v, k, λ) design has a symmetric 2- (v', k', λ') subdesign (possibly degenerate) then $(k'v - kv')^2 \leq (k - \lambda)(v - v')^2$.*

Proof. Take $b = v, r = k, b' = v', m' = v'k'$ and $\lambda_2^2 = k - \lambda$ and apply 5.1. \square

Corollary 5.3 *Let X be a subset of the points and let Y be a subset of the blocks of a 2- (v, k, λ) design D , such that no point of X is incident with a block of Y , then $kr|X||Y| \leq (r - \lambda)(v - |X|)(b - |Y|)$. If equality holds then the incidence structure D' formed by the points in X and the blocks not in Y is a 2-design.*

Proof. Take $m' = 0, v' = |X|, b' = |Y|$ and $\lambda_2^2 = r - \lambda$. Now 5.1 gives the inequality and that D' is a 1-design. But then D' is a 2-design, because D is. \square

If equality holds in 5.2, the subdesign is called *tight*. There are many examples of tight subdesigns of a symmetric designs, see Haemers and Shrikhande [17] or Jungnickel [21]. Wilbrink used Theorem 5.1 to shorten the proof of Feit's result on the number of points and blocks fixed by an automorphism group of a symmetric design, see Lander [22]. The inequality of the second corollary is for example tight for hyperovals and (more generally) maximal arcs in finite projective planes. If we take $|X| = v - k$ we obtain $|Y| \leq \frac{b}{m}$, which is Mann's inequality for the number of repeated blocks in a 2-design. This is an unusual approach to Mann's inequality. For bounds concerning the intersection numbers of D (these are the possible intersection sizes of two blocks of D) it is mostly better to consider the matrix $N^T N$, whose entries are precisely the intersection numbers. We give one illustration.

Theorem 5.4 *Suppose ρ is an intersection number of a 2- (v, k, λ) design D with b blocks and r blocks through a point.*

i $\rho \geq k - r + \lambda$.

ii *Calling blocks equivalent if they are the same or meet in $k - r + \lambda$ points, defines an equivalence relation.*

iii *The number of blocks in an equivalence class is at most $\frac{b}{b - v + 1}$.*

iv *Equality in (iii) for all classes implies that the intersection size of two distinct blocks only depends on whether these blocks are in the same class or in different classes (that is, D is strongly resolvable).*

Proof. Put $A = N^T N$, then $\lambda_1 = rk, \lambda_2 = r - \lambda$ and A has a principal submatrix

$$B = \begin{bmatrix} k & \rho \\ \rho & k \end{bmatrix}.$$

So $\mu_1 = k + \rho$ and $\mu_2 = k - \rho$ and (i) follows from Cauchy interlacing.

Assume the first two blocks b_1 and b_2 meet in $k - r + \lambda$ points, then $\mu_2 = \lambda_2$ and by 2.1(ii), A has a λ_2 -eigenvector $[1, -1, 0, \dots, 0]^\top$ (since B has μ_2 -eigenvector $[1, -1]^\top$), which implies that $(A)_{i,1} = (A)_{i,2}$ for $i = 3, \dots, b$. Hence every block $\notin \{b_1, b_2\}$ meets b_1 and b_2 in the same number of points. Therefore having intersection $k - r + \lambda$ is a transitive relation which proves (ii).

Suppose the first b' blocks of D are equivalent. This gives a partitioning of A with quotient matrix

$$B = \begin{bmatrix} x & rk - x \\ (rk - x)\frac{b'}{b-b'} & rk - (rk - x)\frac{b'}{b-b'} \end{bmatrix},$$

wherein $x = b'(k - r + \lambda) + r - \lambda$. Then $\mu_1 = kr = \lambda_1$ and $\mu_2 = x - (rk - x)\frac{b'}{b-b'}$. Now $\mu_2 \geq \lambda_b \geq 0$ leads to the inequality (iii) (using the 2-design identities $bk = vr$ and $rk - v\lambda = r - \lambda$).

From the proof of (ii) we know that $(A)_{i,1} = (A)_{i,2} = \dots = (A)_{i,b'}$ for $i = b' + 1, \dots, b$. Equality in (iii) implies tight interlacing, therefore the row sums of block $A_{2,1}$ are constant hence all entries of $A_{2,1}$ are equal. This proves (iv). \square

See [14] for more general and other results on intersection numbers. The above theorem appeared in Beker and Haemers [1], where the intersection number $k - r + \lambda$ is treated in detail. Properties (i) and (ii), however, are much older and due to Majumdar [25].

6 Laplace matrix

The Laplace matrix L of a graph G is defined by

$$(L)_{i,j} = \begin{cases} \text{the degree of } i \text{ if } i = j, \\ -1 \text{ if } i \text{ and } j \text{ are adjacent,} \\ 0 \text{ otherwise.} \end{cases}$$

This matrix is singular and positive semi-definite with eigenvalues

$$0 = \theta_1 \leq \dots \leq \theta_n,$$

say (Laplace eigenvalues are usually ordered increasingly). If G is regular of degree k with (standard) adjacency matrix A then $L = kI - A$, so $\theta_i = k - \lambda_i$ and we have an easy one to one correspondence between eigenvalues of L and A . For non-regular graphs there is a different behaviour and the Laplace spectrum seems to be the more natural one. For instance, the number of components equals the nullity of L (i.e. the multiplicity of the eigenvalue 0), whilst this number is not deducible from the spectrum of A (indeed, $K_{1,4}$ and C_4 plus an isolated vertex have the same standard spectrum). Notice that the Laplace matrix of a subgraph G' of G is not a submatrix of L , unless G' is a component. So the interlacing techniques of Section 3 do not work in such a straightforward manner

here. But we can obtain results if we consider off-diagonal submatrices of L in a way similar to the previous section.

Lemma 6.1 *Let X and Y be disjoint sets of vertices of G , such that there is no edge between X and Y , then*

$$\frac{|X||Y|}{(n-|X|)(n-|Y|)} \leq \left(\frac{\theta_n - \theta_2}{\theta_n + \theta_2} \right)^2.$$

Proof. Put $\theta = -\frac{1}{2}(\theta_n + \theta_2)$ and

$$A = \begin{bmatrix} O & L + \theta I \\ L + \theta I & O \end{bmatrix}.$$

Then $\lambda_1 = -\lambda_{2n} = \theta$ and $\lambda_2 = -\lambda_{2n-1} = \frac{1}{2}(\theta_n - \theta_2)$. The sets X and Y give rise to a partitioning of A with quotient matrix

$$B = \begin{bmatrix} 0 & 0 & \theta & 0 \\ 0 & 0 & \theta - \theta \frac{|X|}{n-|Y|} & \theta \frac{|X|}{n-|Y|} \\ \theta \frac{|Y|}{n-|X|} & \theta - \theta \frac{|Y|}{n-|X|} & 0 & 0 \\ 0 & \theta & 0 & 0 \end{bmatrix}.$$

Clearly $\mu_1 = \lambda_1 = \theta$ and $\mu_4 = \lambda_{2n} = -\theta$. Using interlacing we find $\theta^2 \frac{|X||Y|}{(n-|X|)(n-|Y|)} = -\mu_2\mu_3 \leq -\lambda_2\lambda_{2n-1} = (\frac{1}{2}(\theta_n - \theta_2))^2$, which gives the required inequality. \square

A direct consequence of this Lemma is an inequality of Helmberg, Mohar, Poljak and Rendl [18], concerning the bandwidth of G . A symmetric matrix M is said to have bandwidth w if $(M)_{i,j} = 0$ for all i, j satisfying $|i - j| \geq w$. The bandwidth $w(G)$ of a graph G is the smallest possible bandwidth for its adjacency matrix (or Laplace matrix). This number (or rather, the vertex order realizing it) is of interest for some combinatorial optimization problems.

Theorem 6.2 *Suppose G is not the empty graph and define $b = \lceil n \frac{\theta_2}{\theta_n} \rceil$, then*

$$w(G) \geq \begin{cases} b & \text{if } n - b \text{ is even,} \\ b - 1 & \text{if } n - b \text{ is odd.} \end{cases}$$

Proof. Order the vertices of G such that L has bandwidth $w = w(G)$. If $n - w$ is even, let X be the first $\frac{1}{2}(n - w)$ vertices and let Y be the last $\frac{1}{2}(n - w)$ vertices. Then 6.1 applies and thus we find the first inequality. If $n - w$ is odd, take for X and Y the first and last $\frac{1}{2}(n - w - 1)$ vertices and the second inequality follows. If b and w have different

parity, then $w - b \geq 1$ and so the better inequality holds. \square

In case $n - w$ is odd, the bound can be improved a little by applying Theorem 6.1 with $|X| = \frac{1}{2}(n - w + 1)$ and $|Y| = \frac{1}{2}(n - w - 1)$. It is clear that the result remains valid if we consider graphs with weighted edges.

Next we consider an application of interlacing by Van Dam and Haemers [9], which gives a bound for the diameter of G .

Lemma 6.3 *Suppose G has diameter d and $n \geq 2$ vertices, and let P be a polynomial of degree less than d , such that $P(0) = 1$, then*

$$\max_{i \neq 1} |P(\theta_i)| \geq \frac{1}{n-1} .$$

Proof. Assume vertex 1 and n have distance d , then $(L^\ell)_{1,n} = 0$ for $0 \leq \ell \leq d - 1$, hence $(P(L))_{1,n} = 0$. Define

$$A = \begin{bmatrix} O & P(L) \\ P(L) & O \end{bmatrix} .$$

Then $(A)_{1,2n} = (A)_{2n,1} = 0$ and A has row sum $P(\theta_1) = P(0) = 1$. This leads to a partition of A with quotient matrix

$$B = \begin{bmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 1 - \frac{1}{n-1} & \frac{1}{n-1} \\ \frac{1}{n-1} & 1 - \frac{1}{n-1} & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} .$$

The eigenvalues of B are $\mu_1 = -\mu_4 = 1$ and $\mu_2 = -\mu_3 = \frac{1}{n-1}$ and those of A are $\pm P(\theta_i)$. Interlacing gives

$$\frac{1}{n-1} = \mu_2 \leq \lambda_2 \leq \max_{i \neq 1} |P(\theta_i)| ,$$

which proves the inequality. \square

The problem is to find a good choice for the polynomial P . If $\theta_2 > 0$ and L has $\ell < d$ distinct non-zero eigenvalues, we can take P such that $P(\theta_i) = 0$ for all $i \in [2, n]$. This leads to a contradiction in 6.3, proving the well-known result that $d \leq \ell - 1$ if G is connected. In general, it turned out that the Chebyshev polynomials are a good choice for P . The Chebyshev polynomial T_ℓ of degree ℓ can be defined by

$$T_\ell(x) = \frac{1}{2}(x + \sqrt{x^2 - 1})^\ell + \frac{1}{2}(x - \sqrt{x^2 - 1})^\ell .$$

We need the following properties (see Rivlin [31])

$$|T_\ell(x)| \leq 1 \text{ if } |x| \leq 1 , \quad |T_\ell(x)| \geq 1 \text{ if } |x| \geq 1 , \quad T_\ell\left(\frac{x+y}{x-y}\right) > \frac{1}{2} \left(\frac{\sqrt{x+\sqrt{y}}}{\sqrt{x-\sqrt{y}}}\right)^\ell \text{ if } x > y \geq 0 .$$

Theorem 6.4 *If G is a connected graph with diameter $d > 1$, then*

$$d < 1 + \frac{\log 2(n-1)}{\log(\sqrt{\theta_n} + \sqrt{\theta_2}) - \log(\sqrt{\theta_n} - \sqrt{\theta_2})}.$$

Proof. Define $Q(x) = T_{d-1}\left(\frac{\theta_n + \theta_2 - 2x}{\theta_n - \theta_2}\right)$ ($\theta_n \neq \theta_2$, since G is not complete or empty) and put $P(x) = \frac{Q(x)}{Q(0)}$ ($Q(0) \neq 0$, since $\frac{\theta_n + \theta_2}{\theta_n - \theta_2} \geq 1$). Then $|Q(\theta_i)| \leq 1$ for $i = 2, \dots, n$ and $P(0) = 1$. By use of 6.3 we have

$$n-1 \geq \min_{i \neq 1} \frac{1}{|P(\theta_i)|} \geq Q(0) = T_{d-1}\left(\frac{\theta_n + \theta_2}{\theta_n - \theta_2}\right) > \frac{1}{2} \left(\frac{\sqrt{\theta_n} + \sqrt{\theta_2}}{\sqrt{\theta_n} - \sqrt{\theta_2}}\right)^{d-1}.$$

and the bound follows by taking logarithms (the connectivity of G guarantees a non-zero denominator) \square

Computing the diameter of a given graph probably goes faster than computing the above bound. Yet the bound can be of use if the graph is not explicitly given, but the involved eigenvalues are. This is for instance the case with coset graphs of linear codes with given weights. Then eigenvalues correspond to weights in the dual code and diameter bounds lead to bounds for the covering radius of the code (see Delorme and Solé [10]).

7 Regularity

Corollary 2.3(ii) gives a sufficient condition for a partition of a matrix A to be regular. This has turned out to be handy for proving various kinds of regularity. In Sections 3 and 5 we mentioned some examples. Here we give a few more. If we apply Theorem 2.3 to the trivial one-class partition of the adjacency matrix of a graph G with n vertices and m edges we obtain

$$\frac{2m}{n} \leq \lambda_1,$$

and equality implies that G is regular. This is a well-known result, see Cvetković, Doob and Sachs [8]. In fact (since $2m = \text{trace}(A^2) = \sum_{i=1}^n \lambda_i^2$), it implies that G is regular if and only if $\sum_{i=1}^n \lambda_i^2 = n\lambda_1$.

Next we consider less trivial partitions. For a vertex v of G , we denote by $X_i(v)$ the set of vertices at distance i from v . The *neighbour partition* of G with respect to v is the partition into $X_0(v)$, $X_1(v)$ and the remaining vertices. If G is connected, the partition into the $X_i(v)$'s is called the *distance partition* with respect to v . We give examples of tight interlacing involving strongly regular and distance-regular graphs. A graph is *distance-regular around v* if the distance partition with respect to v is regular. If G is distance-regular around each vertex with the same quotient matrix, then G is called *distance-regular*. A *strongly regular graph* is a distance-regular graph of diameter 2. A

distance-regular graph of diameter d has precisely $d + 1$ distinct eigenvalues, being the eigenvalues of the quotient matrix of the distance partition. See Brouwer, Cohen and Neumaier [3] for more about distance-regular graphs. For strongly regular graphs there is a nice survey by Seidel [32].

Theorem 7.1 *Suppose G is regular of degree k ($0 < k < n - 1$) and let t_v be the number of triangles through the vertex v . Then*

$$nk - 2k^2 + 2t_v \leq -\lambda_2 \lambda_n (n - k - 1) .$$

If equality holds for every vertex, then G is strongly regular.

Proof. The neighbour partition has the following quotient matrix

$$B = \begin{bmatrix} 0 & k & 0 \\ 1 & \frac{2t_v}{k} & \frac{k^2 - k - 2t_v}{k} \\ 0 & \frac{k^2 - k - 2t_v}{n - k - 1} & \frac{nk - 2k^2 + 2t_v}{n - k - 1} \end{bmatrix} .$$

Interlacing gives:

$$k \frac{nk - 2k^2 + 2t_v}{n - k - 1} = -\det(B) = -k\mu_2\mu_3 \leq -k\lambda_2\lambda_n .$$

This proves the inequality. If equality holds then $\lambda_2 = \mu_2$ and $\lambda_n = \mu_3$ so (since $k = \lambda_1 = \mu_1$) the interlacing is tight and the neighbour partition is regular with quotient matrix B . By definition, equality for all vertices implies that G is strongly regular. \square

The average number of triangles through a vertex is $\frac{1}{2n} \text{trace}(A^3) = \frac{1}{2n} \sum_{i=1}^n \lambda_i^3$. So if we replace t_v by this expression the above inequality remains valid. Equality then means automatically equality for all vertices so strong regularity. In [16] we looked for similar results for distance-regular graphs of diameter $d > 2$, in order to find sufficient conditions for distance-regularity in terms of the eigenvalues. Therefor one needs to prove regularity of the distance partition. The problem is, however, that in general all eigenvalues $\neq \lambda_1$ of a distance-regular graph have a multiplicity greater than 1, whilst the quotient matrix has all multiplicities equal to 1. So for $d \geq 3$ there is not much chance for tight interlacing. But because of the special nature of the partition we still can conclude regularity.

Lemma 7.2 *Let A be a symmetric partitioned matrix such that $A_{i,j} = 0$ if $|i - j| > 1$ and let B be the quotient matrix. For $i = 1, \dots, m$, let $v_i = [v_{i,1}, \dots, v_{i,m}]^T$ denote a μ_i -eigenvector of B . If $\lambda_0 = \mu_0$, $\lambda_1 = \mu_1$ and $\lambda_n = \mu_m$ and if any three consecutive rows of $[v_1 \ v_2 \ v_m]$ are independent, then the partition is regular.*

Proof. By (iii) of Theorem 2.1 $A\tilde{S}v_i = \mu_i\tilde{S}v_i$ for $i = 1, 2, m$. By considering the ℓ^{th} block row of A we get

$$v_{i,\ell-1}A_{\ell,\ell-1}\mathbb{1} + v_{i,\ell}A_{\ell,\ell}\mathbb{1} + v_{i,\ell+1}A_{\ell,\ell+1}\mathbb{1} = \mu_i v_{i,\ell}\mathbb{1} \quad \text{for } i = 1, 2, m,$$

(wherein the undefined terms have to be taken equal to zero). Since, for $i = 1, 2, m$ and $j = \ell - 1, \ell, \ell + 1$, the matrix $(v_{i,j})$ is non-singular, we find $A_{\ell,j}\mathbb{1} \in \langle \mathbb{1} \rangle$ for $j = \ell - 1, \ell, \ell + 1$ (and hence for $j = 1, \dots, m$). Thus the partition is regular. \square

In [16] it was proved that the independence condition in the above lemma is always fulfilled if we consider the distance partition of a graph. So we have

Theorem 7.3 *Let G be a connected graph and let B be the quotient matrix of the distance partition with respect to a vertex v . If $\lambda_0 = \mu_0$, $\lambda_1 = \mu_1$ and $\lambda_n = \mu_m$ then G is distance-regular around v .*

Using this result it was proved (among others) that if G has the same spectrum and the same number of vertices at maximal distance from each vertex as a distance regular graph G' of diameter 3, then G is distance-regular (with the same parameters as G').

As a last illustration of tight interlacing we prove a result from [14] used by Peeters [30] in his contribution to the special Seidel-volume. Suppose G is a strongly regular graph, pseudo-geometric to a generalized quadrangle of order (s, t) . This means that G has $n = (s + 1)(st + 1)$ vertices and the following quotient matrix for its neighbour partitions

$$\begin{bmatrix} 0 & st + s & 0 \\ 1 & s - 1 & st \\ 0 & t + 1 & st + s - t - 1 \end{bmatrix},$$

for integers s and t . So the eigenvalues of G are $\lambda_1 = st + s$, $\lambda_2 = s - 1$ and $\lambda_n = -t - 1$.

Proposition 7.4 *Let C be a component of the graph induced by the neighbours of any vertex v of G . Let C have c vertices. Then c is a multiple of s and every vertex not adjacent to v is adjacent to precisely $\frac{c}{s}$ vertices of C .*

Proof. Assume $c < st + s$ (otherwise the result is immediate), then C gives rise to a refinement of the neighbour partition with quotient matrix

$$B = \begin{bmatrix} 0 & c & st + s - c & 0 \\ 1 & s - 1 & 0 & st \\ 1 & 0 & s - 1 & st \\ 0 & \frac{c}{s} & t + 1 - \frac{c}{s} & st + s - t - 1 \end{bmatrix}.$$

It easily follows that B has eigenvalues $\mu_1 = st + s$, $\mu_2 = \mu_3 = s - 1$ and $\mu_4 = -t - 1$. Thus we have tight interlacing, so the partition is regular and the result follows. \square

8 Miscellanea

Mohar [27] obtained necessary conditions for the existence of a long cycle in a graph G using Cauchy interlacing. In particular he finds an eigenvalue condition for Hamiltonicity.

Theorem 8.1 *Let G be regular of degree k . If G has a cycle of length ℓ , then*

$$2 \cos \frac{i\pi}{\ell} \leq \sqrt{k + \lambda_{2i+1}} \quad \text{for } i = 1, \dots, \lfloor \frac{n-1}{2} \rfloor .$$

Proof. Let N be the vertex-edge incidence matrix of G . Consider the (bipartite) graph G' with adjacency matrix

$$A' = \begin{bmatrix} O & N \\ N^\top & O \end{bmatrix} .$$

(G' is called the subdivision of G ; roughly, G' is obtained from G by putting a vertex of degree 2 in the middle of each edge.) If G has a cycle of length ℓ , then G' has a cycle of length 2ℓ as an induced subgraph, so the eigenvalues of the 2ℓ -cycle interlace the eigenvalues of A' . Since G has adjacency matrix $A = NN^\top - kI$ (see also the beginning of Section 5), the eigenvalues of G' are $\pm\sqrt{k + \lambda_i}$ and (possibly) zero. The eigenvalues of the 2ℓ -cycle are $2 \cos(\frac{i\pi}{\ell})$ ($i = 0, \dots, 2\ell-1$) and Cauchy interlacing finishes the proof. \square

For example, if G is the Petersen graph and $\ell = 10$, then $i = 3$ gives $2 \cos \frac{3\pi}{10} \leq 1$, which is false. This proves that the Petersen graph is not Hamiltonian.

Eigenvalues of graphs have application in chemistry via the so called Hückel theory, see [8]. For instance, a (carbon) molecule is chemically stable if its underlying graph has half of its eigenvalues positive and half its eigenvalues negative. Manolopoulos, Woodall and Fowler [26] proved that certain graphs, feasible for a molecule structure, satisfy the desired eigenvalue property and hence provide chemically stable molecules. Their method essentially uses interlacing, though they call it Rayleigh's inequalities. We illustrate their approach by considering the more general question how to make graphs (on an even number of vertices) with $\lambda_{\frac{1}{2}n} \geq 0$ and $\lambda_{\frac{1}{2}n+1} \leq 0$. By Theorem 3.1, graphs with a coclique of size $\frac{1}{2}n$ satisfy this property. This includes the bipartite graphs. But there are many more such graphs. We call a graph G an *expanded line graph* of a graph G' if G can be obtained from G' in the following way. The vertices of G are all the ordered pairs (i, j) for which $\{i, j\}$ is an edge in G' . Vertices corresponding to the same edge are adjacent and vertices corresponding to disjoint edges are not adjacent. Of the vertices of G that correspond to intersecting edges $\{i, j\}$ and $\{i, k\}$ (say) of G' , either (i, j) is adjacent to (k, i) , or (j, i) is adjacent to (i, k) , but (i, j) is never adjacent to (i, k) and (j, i) is never adjacent to (k, i) . For example the triangle has the following two expanded line graphs.



Theorem 8.2 *If G is an expanded line graph of a graph G' with adjacency matrix A , then*

- i. $\lambda_{\frac{1}{2}n} \geq 0$ and equality implies that A has a 0-eigenvector $u = [u_1, \dots, u_n]^\top$, where $u_i = u_j$ if i and j correspond to the same edge of G' .
- ii. $\lambda_{\frac{1}{2}n-1} \leq 0$ and equality implies that A has a 0-eigenvector $u = [u_1, \dots, u_n]^\top$, where $u_i = -u_j$ if i and j correspond to the same edge of G' .

Proof. We start with a matrix description of the above construction. Let N be the vertex-edge incidence matrix of G' and let \tilde{N} be a matrix obtained from N by replacing in each column one of the two 1's by a -1 (arbitrarily). Consider the matrices $B = N^\top N$ and $\tilde{B} = \tilde{N}^\top \tilde{N}$ (so that $B - 2I$ is the adjacency matrix of the line graph of G' and $\tilde{B} - 2I$ is a signed version of it). Next we substitute a 2×2 matrix for every entry of \tilde{B} as follows. Replace

$$1 \text{ by } \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix} \text{ or } \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \quad -1 \text{ by } \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \text{ or } \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix},$$

$$2 \text{ by } \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \text{ and } 0 \text{ by } \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}.$$

Then the matrix A , thus obtained, is the adjacency matrix of an expanded line graph of G' and every expanded line graph of G' can be obtained like this. By construction we have a partition of A into $m = \frac{1}{2}n$ classes of size 2 with quotient matrix $\frac{1}{2}B$. Obviously B has smallest eigenvalue $\mu_m \geq 0$. So interlacing gives $\lambda_m \geq 0$ and by (ii) of Theorem 2.1 we have the required 0-eigenvector in case $\lambda_m = \mu_m = 0$. To prove (ii), we multiply every odd row and column of A by -1 . Then the eigenvalues of A remain the same, but now the partition has quotient matrix $-\frac{1}{2}\tilde{B}$ which has largest eigenvalue at most 0 and (ii) follows by interlacing. \square

In [26] the authors considered so-called leapfrog fullerenes, which are special cases of expanded line graphs. By use of the second parts of the above statements they were able to show that leapfrog fullerenes have no eigenvalue 0, and so give rise to stable molecules.

Eigenvalue interlacing has been applied to graphs in many more cases than mentioned in this paper. For instance, Brouwer and Mesner [5] used it to prove that the connectivity of a strongly regular graph equals its degree and in Brouwer and Haemers [4] eigenvalue interlacing is a basic tool for their proof of the uniqueness of the Gewirtz graph. There are also some papers using eigenvalue interlacing that are submitted to the same Seidel-volume as the present paper. Brouwer [2] uses it to find bounds for the toughness of a graph and Ghinelly and Löwe [12] use interlacing to reconstruct generalized quadrangles.

References

- [1] H.J. Beker and W.H. Haemers, *2-designs with an intersection number $k - n$* , J. Comb. Theory A-28 (1980) 64-81.
- [2] A.E. Brouwer, *Toughness and spectrum of a graph*, preprint.
- [3] A.E. Brouwer, A.M. Cohen and A. Neumaier, *Distance-regular graphs*, Springer-Verlag, 1989.
- [4] A.E. Brouwer and W.H. Haemers, *The Gewirtz graph - An exercise in the theory of graph spectra*, Europ. J. Comb. 14 (1993) 397-407.
- [5] A.E. Brouwer and D.M. Mesner, *The connectivity of strongly regular graphs*, Europ. J. Comb. 6 (1985) 215-216.
- [6] R. Courant and D. Hilbert, *Methoden der Mathematischen Physik*, Springer verlag, Berlin, 1924.
- [7] D.M. Cvetković, *Graphs and their spectra*, (Thesis, Univ. Beograd), Publ. Elektrotehn. Fak. Ser. Mat. Fiz. 354-356, (1971) 1-50.
- [8] D.M. Cvetković, M. Doob and H. Sachs, *Spectra of graphs - theory and applications*, V.E.B. Deutscher Verlag der Wissenschaften, Berlin, 1979.
- [9] E.R. van Dam and W.H. Haemers, *Eigenvalues and the diameter of graphs*, Discussion paper 9343, Center for economic research, Tilburg University, 1993; also Lin. Multilin. Alg. (to appear).
- [10] C. Delorme and P. Solé, *Covering index, covering radius and eigenvalues*, Europ. J. Comb. 12 (1991), 95-108.
- [11] P. Delsarte, J.M. Goethals and J.J. Seidel, *Spherical codes and designs*, Geometriae Dedicata 6 (1977), 363-388; also in: *Geometry and combinatorics* (Selected works of J.J. Seidel edited by D.G. Corneil and R. Mathon), Academic Press, 1991, pp. 68-93.
- [12] D. Ghinelli and S. Löwe, *Generalized quadrangles with a regular point and association schemes*, preprint.
- [13] C.D. Godsil, *Algebraic combinatorics*, Chapman and Hall, New York, 1993.
- [14] W.H. Haemers, *Eigenvalue techniques in design and graph theory* (thesis Technical University Eindhoven 1979), Math. Centre Tract 121, Mathematical Centre, Amsterdam, 1980.
- [15] W.H. Haemers, *Eigenvalue methods*, in: *Packing and covering in combinatorics* (A. Schrijver ed.), Math. Centre Tract 106, Mathematical Centre, Amsterdam, 1979, pp. 15-38.
- [16] W.H. Haemers, *Distance-regularity and the spectrum of graphs*, Research Memorandum FEW 582, Tilburg University, 1992; also Linear Alg. Appl. (to appear).

- [17] W.H. Haemers and M.S. Shrikhande, *Some remarks on subdesigns of symmetric designs*, J. Stat. Plann. Inference 3 (1979) 361-366.
- [18] C. Helmborg, B. Mohar, S. Poljak and F. Rendl, *A spectral approach to bandwidth and separator problems in graphs*, Proceedings third IPCO conf. (G. Rinaldi and L. Wolsey eds.), (1993) 183-194.
- [19] A.J. Hoffman, *On eigenvalues and colourings of graphs*, in: *Graph theory and its applications*, (B. Harris, ed.), Acad. Press, New York, 1970, pp. 79-91.
- [20] D.R. Hughes and F.C. Piper, *Design theory*, Cambridge Univ. Press, Cambridge, 1985.
- [21] D. Jungnickel, *On subdesigns of symmetric designs*, Math. Z. 181 (1982), 383-393.
- [22] E.S. Lander, *Symmetric designs: an algebraic approach*, London Math. Soc. Lect. Note Ser. 74, Cambridge Univ. Press, Cambridge, 1983.
- [23] L. Lovász, *On the Shannon capacity of a graph*, IEEE trans. Inf. Theory IT-25 (1979) 1-7.
- [24] L. Lovász, *Combinatorial problems and exercises*, North Holland, Amsterdam, 1979.
- [25] K.N. Majumdar, *On some theorems in combinatorics relating to incomplete block designs*, Ann. Math. Statist. 24 (1953), 379-389.
- [26] D.E. Manolopoulos, D.R. Woodall and P.W. Fowler, *Electronic stability of fullerenes: eigenvalue theorems for leapfrog carbon clusters*, J. Chem. Soc. Faraday Trans. 88 (1992), 2427-2435.
- [27] B. Mohar, *A domain monotonicity theorem for graphs and Hamiltonicity*, Discrete Appl. Math. 36 (1992), 169-177.
- [28] S.E. Payne, *An inequality for generalized quadrangles*, Proc. Amer. Math. Soc. 71 (1978) 147-152.
- [29] S.E. Payne and J.A. Thas, *Finite generalized quadrangles*, Research Notes in Math. 110, Pitman Adv. Publ. Program, London, 1984.
- [30] M.J.P. Peeters, *Uniqueness of strongly regular graphs having minimal p-rank*, Research Memorandum FEW 626, Tilburg University, 1993; also Lin. Alg. Appl. (to appear).
- [31] T.J. Rivlin, *Chebyshev polynomials* (second edition), John Wiley and Sons, New York, 1990.
- [32] J.J. Seidel, *Strongly regular graphs*, in: *Surveys in combinatorics* (B. Bollobás ed.), London Math. Soc. Lect. Note Ser. 38, Cambridge Univ. Press, Cambridge, 1979, pp. 157-180.