Tilburg University

Long-term quality-of-life assessment after laparoscopic and classic cholecystectomy

Lachinski, A.J.; Markuszewska-Proczko, M.; Stefaniak, T.J.; Vingerhoets, A.J.J.M.

Published in:
Surgical Endoscopy

Publication date:
2004

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Lachinski, A. J., Markuszewska-Proczko, M., Stefaniak, T. J., & Vingerhoets, A. J. J. M. (2004). Long-term quality-of-life assessment after laparoscopic and classic cholecystectomy. *Surgical Endoscopy*, 18(7), 1152-1153.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Long-term quality-of-life assessment after laparoscopic and classic cholecystectomy

Positive results of studies comparing the laparoscopic and classic treatment of surgical problems are always encouraging. They reinforce what surgeons believe in and they provide us with reliable scientific material that supports our point of view when we have to defend ourselves against the criticism of more conservative colleagues. Such findings also provide evidence to buttress what we observe in everyday practice—that after laparoscopic treatment patients experience more satisfaction and a better quality of life.

Therefore, the recent article by Topcu et al. should give us much reason for satisfaction [9]. However, the article fails to supply a psychological rationale for the comparison of those two methods of surgical intervention for cholelithiasis. In addition, it has some methodological weaknesses that undermine their optimistic conclusions.

First, quality of life, although very popular and in fashion nowadays, does not seem to be the most appropriate psychological parameter for a differentiation between two groups of patients suffering from a medical problem that is not likely to have a profound impact on quality of life. Although quality of life is especially important for chronic medical conditions, its relevance for acute medical conditions is moderate at best. It has indeed been demonstrated that a decrease in health-related quality of life can be observed in progressive or chronic diseases, such as chronic pancreatitis [6], chronic pain syndromes [2], and cancers [1]. Although there is no doubt that cholelithiasis may decrease the quality of life during its acute symptomatic phase, it is highly improbable that it will be significantly altered in the postoperative course after cholecystectomy, independent of the operative technique that is applied.

Second, the protracted period of ~3 years that elapsed between the operation and the data collection casts more doubt on the reliability of the findings. Health-related quality of life is a dynamic psychological variable that may be influenced by many factors—such as additional illnesses that developed later or personal problems—that were not controlled by the researchers. In addition, there is reason to assume that sociocultural factors—in particular, income and insurance—also have a strong association with quality of life [3–5]. It is stated in

the methods section that the patients operated laparoscopically had to pay for the operation, whereas the costs for the open procedures were covered by insurance. Given the ample body of literature supporting the wealth–health connection [3–5], it makes no sense to measure quality of life such a long time after cholecystectomy. The only way to obtain reliable and equivocal quality-of-life data in these conditions would be to measure it repeatedly—for example, every month after the operation. In addition, because there was no random assignment, the preoperative quality-of-life scores should have been determined, so as to control for possible differences in pretreatment levels. It is very likely that the reported posttreatment differences in quality of life simply reflect preexisting pretreatment differences, due to the connection of this measure with income and insurance.

The fact that significant differences were found for every aspect of quality of life also lends support to this hypothesis. We could speculate that the classic treatment might be worse in terms of cosmesis. Thus, it would be understandable if the social aspect of quality of life were impacted, but it is difficult to believe that other aspects of quality of life still show significant differences as long as 3 years after the operation.

To summarize, the study by Topcu et al. although conceptually interesting, is characterized by some methodological drawbacks that compromise its scientific value. It lacks initial measurement of quality of life, and the period between the operation and the measurement is too long. Because randomization was not possible, we would at least have expected that the authors would have controlled statistically for differences in the relevant pretreatment variables—in particular, income levels and type of insurance. In addition, more repeated measures, starting shortly after the treatment, would have provided greater insight into the dynamics of this most important outcome variable.

It should be emphasized that the study represents an important approach to the evaluation of the results of surgical treatment. This approach uses psychological parameters, such as quality of life, body image, or subjective perception of the severity of the illness, as measures to evaluate the effects of a medical procedure [8]. In addition to mortality, morbidity, and recurrence (e.g., cancer), these subjective psychological parameters

should be considered essential in determining the effect of treatment [7].

References

1. Bottomley A, Efficace F, Thomas R, Vanvoorden V, Ahmedzai SH (2003) Health-related quality of life in non-small-cell lung cancer: methodologic issues in randomized controlled trials. *J Clin Oncol* 21: 2982–2992
2. Kalso E, Allan L, Dellemijn PL, Faura CC, Ilias WK, Jensen TS, Perrot S, et al. (2003) Recommendations for using opioids in chronic non-cancer pain. *Eur J Pain* 7: 381–386
3. King LA, Napa CK (1998) What makes a life good? *J Pers Soc Psychol* 75: 156–65
4. Klose T (2003) A utility-theoretic model for QALYs and willingness to pay. *Health Econ* 12: 17–31
5. Lee PP, Smith JP, Kington RS (1999) The associations between self-rated vision and hearing and functional status in middle age. *Ophthalmology* 106: 401–405
6. Makarewicz W, Stefaniak T, Kossakowska M, Basinski A, Suchorzewski M, Stanek A, Gruca ZB (2003) Quality of life improvement after videothoroscopic splanchnicectomy in chronic pancreatitis patients: case control study. *World J Surg* 27: 986–911
7. Stefaniak T, Kaska L, Makarewicz W, Gruca Z (2003) Psychological and analytic perspective to reliable outcome validation in minimally invasive surgery. Abstracts of 11th EAES Congress, 17
8. Stefaniak T, Trus M, Babinska D, Vingerhoets AJJM (2003) Laparoscopic versus classic cholecystectomy—recovery mediated via wound-related subjective perspective of illness. Abstracts of 11th EAES Congress, 49
9. Topcu O, Karakayali F, Kuzu MA, Ozdemir S, Erverdi N, Elhan A, Aras N (2003) Comparison of long-term quality of life after laparoscopic and open cholecystectomy. *Surg Endosc* 17: 291–295

A. Lachinski¹
A. Vingerhoets¹
M. Markuszewska-Proczko²
T. Stefaniak^{1,3}

¹ Department of General Gastroenterological, and Endocrinological Surgery
 Medical University of Gdansk
 1 Kieturakis Street, PL-80-742
 Gdansk, Poland

² Department of Psychology and Health, and Research Institute for Psychology and Health
 Tilburg University, Tilburg
 The Netherlands

³ Laboratory of Psychosomatics and Psychology of Surgery
 Department of General, Gastroenterological, and Endocrinological Surgery
 Medical University of Gdansk
 Poland

Online publication: 12 May 2004