Tilburg University

Extension to higher dimensions of the Jaeschke-Eicker result on the standardized empirical process

Einmahl, J.H.J.

Published in:
Communications in statistics: Part A: Theory and methods

Publication date:
1996

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Einmahl, J. H. J. (1996). Extension to higher dimensions of the Jaeschke-Eicker result on the standardized empirical process. *Communications in statistics: Part A: Theory and methods*, 25(4), 813-822.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Eindhoven University of Technology

Department of Mathematics and Computing Science

Probability Theory, Statistics, Operations Research and Systems Theory

Reprint Series

**Extension to higher dimensions of the
Jaeschke-Eicker result on the standardized
empirical process**

by

J.H.J. Einmahl

Reprint COSOR 96/387

**Extension to higher dimensions of the
Jaeschke-Eicker result on the standardized
empirical process**

by

J.H.J. Einmahl

Reprint COSOR 96/387

This article appeared in: Commun.Statist.-Theory Meth.
25, No. 4, 1996, p. 813-822.

Reprinted for private circulation by the author.

Eindhoven University of Technology

Department of Mathematics and Computing Science
Probability Theory, Statistics, Operations Research and
Systems Theory Group
Secretary, Main Building, Room 9.10, Tel. (040)-2473130
P.O. Box 513, 5600 MB Eindhoven, Netherlands

Volume 25, Number 4, 1996, p. 813-822

communications in statistics

THEORY AND METHODS

Editor-in-Chief

WILLIAM B. SMITH
Department of Statistics
Texas A&M University
College Station, TX 77843-3143

Senior Editors

WILLIAM R. SCHUCANY
Southern Methodist University, Texas

ANANT M. KSHIRSAGAR
University of Michigan, Ann Arbor

Algorithms Editor

JAMES E. GENTLE
George Mason University, Virginia

Founding Editor

DONALD B. OWEN
(Volumes 1-20)

EXTENSION TO HIGHER DIMENSIONS OF THE
JAESCHKE-EICKER RESULT
ON THE STANDARDIZED EMPIRICAL PROCESS

John H.J. Einmahl

Dept. of Mathematics and Computing Science, Eindhoven University of
Technology, P.O. Box 513, 5600 MB Eindhoven, The Netherlands

Key Words and Phrases: Asymptotic distribution; empirical process; heavy weights.

ABSTRACT

The asymptotic distribution of the sup-norm of the heavily weighted empirical process is established in the multidimensional case. This theorem extends in particular the famous result in Jaeschke (1975, 1979) to higher dimensions. There is a striking difference between the behaviour for higher dimensions and that for dimension one, especially the limiting distribution is now a simple transformation of a standard exponential random variable.

1. INTRODUCTION AND MAIN RESULTS

Let $X_1, \dots, X_n$ be independent random vectors, each uniformly distributed on $[0, 1]^d$, $d \in N$, and define the multivariate empirical distribution function by

$$F_n(t) = \frac{1}{n} \# \{1 \leq i \leq n : X_i \leq t\}, \quad t \in [0, 1]^d,$$

where ' $\leq$ ' denotes the componentwise ordering. Note that $F(t) = P(X_i \leq t) = \prod_{j=1}^d t_j$, $t = (t_1, \dots, t_d) \in [0, 1]^d$. The multivariate empirical process is written as

$$\alpha_n(t) = n^{1/2}(F_n(t) - F(t)), \quad t \in [0, 1]^d.$$

Observe that $\text{Var}(\alpha_n(t)) = F(t)(1 - F(t))$ and hence it is natural to consider the standardized empirical process

$$\frac{\alpha_n(t)}{(F(t)(1 - F(t)))^{1/2}}, \quad t \in [0, 1]^d, \quad (1.1)$$

a process having constant variance equal to 1, or, more generally,

$$\alpha_{n,\nu}(t) = \frac{\alpha_n(t)}{(F(t)(1 - F(t)))^\nu}, \quad t \in [0, 1]^d, \quad \nu \in [0, 1]. \quad (1.2)$$

(We will not consider the case $\nu > 1$, since then $\sup_{t \in [0, 1]^d} |\alpha_{n,\nu}(t)| = \infty$ a.s.) It is well-known that for $\nu \in [0, \frac{1}{2})$, and not for $\nu \in [\frac{1}{2}, 1]$, $\alpha_{n,\nu}$ converges weakly to $B/(F(1 - F))^\nu$, where B is a continuous mean zero Gaussian process with $EB(s)B(t) = F(s \wedge t) - F(s)F(t)$, where $s \wedge t$ has to be understood component-wise. Here, however, we are interested in the values of $\nu \in [\frac{1}{2}, 1]$, leading to so-called heavily weighted empirical processes, with special interest in the case $\nu = \frac{1}{2}$. To be more precise, we will consider the weak limiting behaviour of

$$\sup_{t \in [0, 1]^d} |\alpha_{n,\nu}(t)|, \quad \nu \in [\tfrac{1}{2}, 1]. \quad (1.3)$$

Of course, for $\nu \in [0, \frac{1}{2})$, we immediately have from the weak convergence of $\alpha_{n,\nu}$ as a process that

$$\sup_{t \in [0, 1]^d} |\alpha_{n,\nu}(t)| \rightarrow_d \sup_{t \in [0, 1]^d} \frac{|B(t)|}{(F(t)(1 - F(t)))^\nu}, \quad \text{as } n \rightarrow \infty.$$

For $d = 1$, the weak limiting behaviour of the properly normalized statistic in (1.3) is well-known. It is the purpose of this paper to extend the one dimensional

results to higher dimensions; surprisingly the behaviour turns out to be completely different then. Let us briefly describe the one dimensional results now. When $d = 1$ the behaviour of the cases $\nu = \frac{1}{2}$ and $\nu \in (\frac{1}{2}, 1]$ is substantially different. First we consider the standardized empirical process, i.e. the case $\nu = \frac{1}{2}$. Set $a_n = (2 \log \log n)^{-\frac{1}{2}}$ and $b_n = (2 \log \log n)^{\frac{1}{2}} + \{\frac{1}{2} \log \log \log n - \frac{1}{2} \log \frac{\pi}{4}\} / (2 \log \log n)^{\frac{1}{2}}$. Then, as $n \rightarrow \infty$,

$$(\sup_{t \in [0,1]} |\alpha_{n,\frac{1}{2}}(t)| - b_n) / a_n \rightarrow_d \Gamma, \quad (1.4)$$

where Γ is a standard Gumbel random variable, i.e. $P(\Gamma \leq x) = \exp(-\exp(-x))$, $x \in \mathbb{R}$. This famous result in empirical process theory is established in Jaeschke (1975, 1979); closely related results (with F in the denominator in (1.1) replaced by F_n) can be found in Eicker (1979). Now we consider $\nu \in (\frac{1}{2}, 1]$. Let N and $\tilde{N}$ be two independent homogeneous unit intensity Poisson processes on $[0, \infty)$, then it is shown in Mason (1983) that, as $n \rightarrow \infty$,

$$n^{\frac{1}{2}-\nu} \sup_{t \in [0,1]} |\alpha_{n,\nu}(t)| \rightarrow_d \sup_{t>0} \frac{|N(t)-t| \vee |\tilde{N}(t)-t|}{t^\nu}. \quad (1.5)$$

Without making this more precise, we mention that the behaviour of the left hand side of (1.5) is determined by the extreme tails of the (uniform) distribution, whereas in (1.4) the moderate tails are responsible for the behaviour of its left hand side.

Now we turn to our main result, i.e. the proper extensions of (1.4) and (1.5) to dimension $d \geq 2$. For this purpose let Y be a mean one exponential random variable ($P(Y \leq x) = 1 - e^{-x}$, $x \geq 0$) and set $L = ((d-1)!Y)^{-\nu}$.

THEOREM 1. Let $d \geq 2$ and $\nu \in [\frac{1}{2}, 1]$, then, as $n \rightarrow \infty$,

$$\frac{\sup_{t \in [0,1]^d} |\alpha_{n,\nu}(t)|}{n^{\nu-\frac{1}{2}} (\log n)^{\nu(d-1)}} \rightarrow_d L. \quad (1.6)$$

Remark 1. It is striking that in the multivariate case the behaviour differs substantially from that in the one dimensional situation. Observe also that in (1.6) the

cases $\nu = \frac{1}{2}$ and $\nu \in (\frac{1}{2}, 1]$ are treated 'simultaneously', in contrast to the behaviour in dimension one. It will become clear from the proof that the behaviour of the left hand side of (1.6) is determined by the left tail ($\{t: F(t) \text{ 'small'}\}$) of the distribution, to be more precise, we will show that the 'multivariate minimum' (the X_i with $F(X_i) = \min_{1 \leq j \leq n} F(X_j)$) plays the dominant role.

Remark 2. There is already a related result for $d = 2$ and $\nu \in (\frac{1}{2}, 1)$ in the literature (Csörgö and Horváth (1990)). This result yields

$$\frac{n^{\frac{1}{2}-\nu} \sup_{t \in [0,1]^d} |\alpha_n(t)|}{(F(t))^\nu} \rightarrow_d \sup_{t_1, t_2 > 0} \frac{|N(t_1, t_2) - t_1 t_2|}{(t_1 t_2)^\nu} \quad (n \rightarrow \infty), \quad (1.7)$$

where N is a homogeneous unit intensity Poisson process on $[0, \infty)^2$. However, it can be readily verified that the right hand side of (1.7) is equal to infinity with probability one. This means that the normalization $(n^{\frac{1}{2}-\nu})$ is not appropriate, which is also confirmed by Theorem 1.

Remark 3. It is standard to generalize the one dimensional results (1.4) and (1.5) to X_i 's having a continuous distribution function, instead of being uniformly $[0, 1]$ distributed. Similarly for $d \geq 2$, it readily follows from the fact that the uniform $[0, 1]^d$ distribution has independent marginals, that Theorem 1 remains true when this distribution is replaced by a probability distribution on $\mathbb{R}^d$ with distribution function $F(t) = \prod_{j=1}^d F_j(t_j)$, $t = (t_1, \dots, t_d) \in \mathbb{R}^d$, where the F_j , $1 \leq j \leq d$, are continuous one dimensional distribution functions on $\mathbb{R}$ (of course, in (1.6), $[0, 1]^d$ has to be replaced by $\mathbb{R}^d$ then). The class of these distributions is still restricted, since the marginals remain independent. On the other hand, when $d \geq 2$, some serious restriction on the distribution of the X_i is needed, since it easily follows that when, e.g., $d = 2$ and the X_i are uniformly distributed on the line segment from $(1, 0)$ to $(0, 1)$, then $\sup_{t \in [0,1]^2} |\alpha_{n,\nu}(t)| = \infty$ a.s. for all $\nu \in (0, 1]$, i.e. trying to extend Theorem 1 to this distribution is senseless.

Now we present an essential generalization of Theorem 1, namely to the case where the marginals may be dependent.

THEOREM 2. Let the X_i have a distribution on $[0, 1]^d$, with a continuous density f with respect to Lebesgue measure on $[0, 1]^d$, $d \geq 2$, satisfying for some m and $M : 0 < m \leq f(t) \leq M < \infty$, $t \in [0, 1]^d$. Then, as $n \rightarrow \infty$, (1.6) remains true for $\nu \in [\frac{1}{2}, 1]$.

Remark 4. Theorem 2 shows that for a wide class of distributions the rate of convergence of $\sup_{t \in [0, 1]^d} |\alpha_{n, \nu}(t)|$ and the limiting law are the same. Observe that this class, in turn, can be extended to distributions on $\mathbb{R}^d$ as indicated in Remark 3. It is also shown in Remark 3, however, that (1.6) is definitely not satisfied for all distributions. Finally, note that the rate in (1.6) is also not the 'slowest' possible one, even not within the class of continuous distribution functions, since, e.g., again when $d = 2$, the uniform distribution on the line segment from $(0, 0)$ to $(1, 1)$ leads to an essentially one dimensional situation, i.e. the rates in (1.4) and (1.5) apply.

2. PROOFS

For the proof of Theorem 1 the following three results on probabilities, concerning the empirical distribution function of independent uniform- $[0, 1]^d$ random vectors, which all can be found in Einmahl (1987, pages 19, 38 and 26 respectively), will be required.

FACT 1. Let $d \in \mathbb{N}$ and $\nu \in [\frac{1}{2}, 1]$. Then for any $\delta \in (0, 1)$, $0 < \alpha \leq \beta \leq \frac{1}{2}(1 - \delta)$ and $\lambda \geq 0$

$$P\left(\sup_{\alpha \leq F(t) \leq \beta} n^{\frac{1}{2}-\nu} |\alpha_n(t)| / (F(t))^\nu \geq \lambda\right) \\ \leq C \int_{(1-\delta)\alpha}^{\beta/(1-\delta)} \frac{(\log(1/x))^{d-1}}{x} \exp\left(-\frac{1}{2}(1-\delta)\lambda^2 (nx)^{2\nu-1} \psi\left(\frac{\lambda}{(n\alpha)^{1-\nu}}\right)\right) dx,$$

where $C = C(d, \delta) \in (0, \infty)$ and $\psi(y) = 2y^{-2}\{(1+y)\log(1+y) - y\}$, $y > 0$.

FACT 2. Let $d \in \mathbb{N}$, $m \in \{1, \dots, n\}$ and $0 < \alpha \leq 1/e$. Then

$$P\left(\sup_{F(t) \leq \alpha} nF_n(t) \geq m\right) \leq c_1 \binom{n}{m} (c_2 \alpha)^m (\log(1/\alpha))^{d-1},$$

where $c_1 = c_1(d)$, $c_2 = c_2(d)$ and $c_1, c_2 \geq 1$. Of course $c_1(1)$ and $c_2(1)$ can both be taken equal to 1.

FACT 3. Let $d \in \mathbb{N}$ and $\gamma \in (0, \infty)$. Then, as $n \rightarrow \infty$,

$$P\left(\sup_{F(t) \leq \frac{(d-1)!}{\gamma n (\log n)^{d-1}}} nF_n(t) = 0\right) \rightarrow e^{-1/\gamma}.$$

PROOF OF THEOREM 1. From the weak convergence of α_n it is immediate that for all $0 < a < b < 1$

$$\frac{1}{n^{\nu-\frac{1}{2}}(\log n)^{\nu(d-1)}} \sup_{a \leq F(t) \leq b} |\alpha_{n,\nu}(t)| \rightarrow_P 0 \quad (n \rightarrow \infty). \quad (2.1)$$

This shows that in the 'middle' no contribution to the limit is made. Hence it suffices to consider the left tail and the right tail separately. However, it can be shown as in Theorem 3.2 in Einmahl (1987), that 'large d-dimensional points' behave (modulo multiplicative constants) as 'small (or large) 1-dimensional points'. By using (1.4) and (1.5), this statement leads to the fact that (2.1) remains true with $b = 1$. So, taking $a = \frac{1}{4}$, it suffices to show, that as $n \rightarrow \infty$,

$$\frac{1}{n^{\nu-\frac{1}{2}}(\log n)^{\nu(d-1)}} \sup_{F(t) \leq \frac{1}{4}} |\alpha_{n,\nu}(t)| \rightarrow_d L. \quad (2.2)$$

Write $a_n = 1/(n(\log n)^{\frac{1}{2}(d-1)})$ and note that

$$\sup_{a_n \leq F(t) \leq \frac{1}{4}} |\alpha_{n,\nu}(t)| \leq \frac{4}{3} \sup_{a_n \leq F(t) \leq \frac{1}{4}} |\alpha_n(t)| / (F(t))^\nu.$$

We hence have for arbitrary $\varepsilon > 0$ by Fact 1, with $\delta = \frac{1}{2}$,

$$P\left(\sup_{a_n \leq F(t) \leq \frac{1}{4}} n^{\frac{1}{2}-\nu} |\alpha_{n,\nu}(t)| \geq \varepsilon (\log n)^{\nu(d-1)}\right)$$

$$\leq C \int_{\frac{1}{2}a_n}^{\frac{1}{2}} \frac{(\log(1/x))^{d-1}}{x} dx \quad (2.3)$$

$$\cdot \exp\left(-\frac{1}{4}\left(\frac{3}{4}\right)^2 \varepsilon^2 (\log n)^{2\nu(d-1)} \left(\frac{1}{2}na_n\right)^{2\nu-1} \psi\left(\frac{3}{4}\varepsilon (\log n)^{\nu(d-1)} (na_n)^{\nu-1}\right)\right).$$

Using $x\psi(x) \rightarrow \infty$ ($x \rightarrow \infty$), the right hand side of (2.3) is for large n bounded from above by

$$C(\log n)^d \exp(-(\log n)^{\frac{1}{2}\nu(d-1)}),$$

which tends to 0, as $n \rightarrow \infty$. Since $\varepsilon > 0$ is arbitrary, it is now sufficient to prove (2.2) with $\frac{1}{4}$ replaced by a_n . But since on $\{t \in [0, 1]^d : F(t) \leq a_n\}$ we have $(1 - a_n)^\nu \leq (1 - F(t))^\nu \leq 1$ and $\lim_{n \rightarrow \infty} (1 - a_n)^\nu = 1$, it finally remains to show that

$$L_n = \frac{1}{n^{\nu-\frac{1}{2}}(\log n)^{\nu(d-1)}} \sup_{F(t) \leq a_n} \frac{|\alpha_n(t)|}{(F(t))^\nu} \rightarrow_d L \quad (n \rightarrow \infty). \quad (2.4)$$

Now Fact 2 yields

$$\begin{aligned} P\left(\sup_{F(t) \leq a_n} nF_n(t) \geq 2\right) &\leq c_1 \binom{n}{2} (c_2 a_n)^2 (\log(1/a_n))^{d-1} \\ &= O((\log n)^{-\frac{1}{2}\nu(d-1)}) \rightarrow 0 \quad (n \rightarrow \infty), \end{aligned} \quad (2.5)$$

and Fact 3 yields

$$P\left(\sup_{F(t) \leq a_n} nF_n(t) = 0\right) \rightarrow 0 \quad (n \rightarrow \infty). \quad (2.6)$$

Statements (2.5) and (2.6) play the crucial role in the proof of (2.4). They imply, that, with arbitrary high probability (n large) there are observations in the region $\{t \in [0, 1]^d : F(t) \leq a_n\}$, but in that region $nF_n(t)$ is at most one. Set $M_n = \sup_{F(t) \leq a_n} nF_n(t)$. We have by (2.5) and (2.6)

$$\mathbb{I}_{\{1\}}(M_n) \rightarrow_P 1 \quad (n \rightarrow \infty). \quad (2.7)$$

Write

$$L'_n = \frac{n^{1-\nu}}{(\log n)^{\nu(d-1)}} \frac{|\frac{1}{n} - F(X)_{1:n}|}{(F(X)_{1:n})^\nu},$$

where $F(X)_{1:n} = \min_{1 \leq i \leq n} F(X_i)$. Then it follows that for $n \geq 3$

$$L_n = L'_n \mathbb{I}_{\{1\}}(M_n) + L_n \mathbb{I}_{\{0,2,3,\dots,n\}}(M_n). \quad (2.8)$$

Also, using (2.7),

$$L'_n \mathbb{I}_{\{1\}}(M_n) / \frac{1}{(\log n)^{\nu(d-1)} n^\nu (F(X)_{1:n})^\nu} \rightarrow_P 1 \quad (n \rightarrow \infty). \quad (2.9)$$

From (2.7)-(2.9) and

$$P\left(\frac{1}{(\log n)^{\nu(d-1)} n^\nu (F(X)_{1:n})^\nu} \leq x\right) = \{P(F(X_1) \geq \frac{1}{x^{1/\nu} n (\log n)^{d-1}})\}^n,$$

it follows that for a proof of (2.4), it remains to show that for $x > 0$, as $n \rightarrow \infty$,

$$\{P(F(X_1) \geq \frac{1}{x^{1/\nu} n (\log n)^{d-1}})\}^n \rightarrow P(L \leq x). \quad (2.10)$$

Using, e.g., that $-\log F(X_1)$ has a gamma distribution with density $((d-1)!)^{-1} x^{d-1} e^{-x} 1_{(0,\infty)}(x)$, a straightforward but tedious calculation shows that the left hand side of (2.10) converges to

$$\exp\left(-\frac{1}{(d-1)! x^{1/\nu}}\right), \text{ as } n \rightarrow \infty. \quad (2.11)$$

Observing that

$$\begin{aligned} P(L \leq x) &= P\left(\frac{1}{((d-1)! Y)^\nu} \leq x\right) = P\left(Y \geq \frac{1}{(d-1)! x^{1/\nu}}\right) \\ &= \exp\left(-\frac{1}{(d-1)! x^{1/\nu}}\right), \end{aligned}$$

now completes the proof of (2.4) and hence that of Theorem 1. $\square$

For the proof of Theorem 2 we will need the following 'local' generalization of the one dimensional probability integral transform, which may be of independent interest.

PROPOSITION 1. Under the condition on f in Theorem 2 we have

$$\lim_{s \downarrow 0} \frac{(d-1)! P(F(X_1) \leq s)}{s(\log(1/s))^{d-1}} = 1. \quad (2.12)$$

PROOF. The proof is essentially easy, but the details are somewhat technical. Therefore we will only give a sketch of the proof. When $F(t) = \prod_{j=1}^d t_j$, i.e. the uniform distribution on $[0, 1]^d$, then (2.12) easily follows, e.g., by using that $-\log F(X_1)$ has a gamma density (see the end of the proof of Theorem 1). If $F(t) = c \prod_{j=1}^d t_j$, $c \in (0, \infty)$, for small $F(t)$, then (2.12) also easily follows by using the fact that it holds for $c = 1$, which we just observed.

Now we turn to the general case. Let $\delta > 0$ be arbitrary. Then, because of continuity of f , we can find an $\eta \in (0, 1)$ such that on $t \in [0, \eta]^d$, $|f(t) - f(\bar{0})| \leq \varepsilon f(\bar{0})$ ($\bar{0} = (0, \dots, 0)$).

Hence we have for small $\eta > 0$

$$\begin{aligned} P(F(X_1) \leq s) \\ &= P(F(X_1) \leq s, X_1 \in [0, \eta]^d) + P(F(X_1) \leq s, X_1 \notin [0, \eta]^d) \\ &\leq P(F'(X'') \leq s) + P(F(X_1) \leq s, X_1 \notin [0, \eta]^d), \end{aligned} \quad (2.13)$$

where F' is a distribution function on $[0, 1]^d$, such that for F' small, $F'(t) = (1 - \varepsilon)f(\bar{0})\prod_{j=1}^d t_j$ and X'' is for small values of its distribution function, distributed according to $F''(t) = (1 + \varepsilon)f(\bar{0})\prod_{j=1}^d t_j$. Write $h(s) = ((d-1)!)^{-1}s(\log(1/s))^{d-1}$, then it easily follows by applying the just treated case with $c = (1 + \varepsilon)f(\bar{0})$, that, as $s \downarrow 0$,

$$P(F'(X'') \leq s) \leq (1 + o(1)) \frac{1 + \varepsilon}{1 - \varepsilon} h(s). \quad (2.14)$$

Also, by using that $m \leq f \leq M$, as $s \downarrow 0$,

$$P(F(X_1 \leq s, X_1 \notin [0, \eta]^d) = o(h(s)). \quad (2.15)$$

Hence by combining (2.13)-(2.15) we see that since $\varepsilon > 0$ is arbitrary that

$$\limsup_{s \downarrow 0} P(F(X_1) \leq s)/h(s) \leq 1.$$

It similarly follows that the 'liminf' is not smaller than 1. $\square$

PROOF OF THEOREM 2. The proof follows along the same lines as that of Theorem 1, using obvious modifications of Facts 1-3 (see section 6.2 in Einmahl (1987) for details). The only problem arises in the very last part of the proof, i.e. proving that the left hand side of (2.10) converges to (2.11). This, however, follows immediately from Proposition 1. $\square$

BIBLIOGRAPHY

- Csörgő, M. and Horváth, L. (1990). Asymptotic tail behaviour of uniform multivariate empirical processes, *Ann. Probab.*, 18, 1723-1738.
- Eicker, F. (1979). The asymptotic distribution of the suprema of the standardized empirical processes, *Ann. Statist.*, 7, 116-138.
- Einmahl, J.H.J. (1987). *Multivariate Empirical Processes*, CWI Tract 32, Mathematisch Centrum, Amsterdam.
- Jaeschke, D. (1975). Über die Grenzverteilung des Maximums der normierten empirischen Verteilungsfunktion, Dissertation, Dortmund University.
- Jaeschke, D. (1979). The asymptotic distribution of the supremum of the standardized empirical distribution function on subintervals, *Ann. Statist.*, 7, 108-115.
- Mason, D.M. (1983). The asymptotic distribution of weighted empirical distribution functions, *Stochastic Process. Appl.*, 15, 99-109.

Nr.	Author(s)	Title
90/290	<i>Vija K. Rohatgi</i> <i>F.W. Steutel</i> <i>Gábor J. Székely</i>	Infinite divisibility of products and quotients of I.I.D. random variables
90/291	<i>J. van der Wal</i>	Monotonicity of the throughput of a closed exponential queueing network in the number of jobs
90/292	<i>M.W.I. van Kraaij</i> <i>W.Z. Venema</i> <i>J. Wessels</i>	Automatic modelling and solving of strategic planning problems
90/293	<i>A.A. Stoorvogel</i> feedback <i>H.L. Trentelman</i>	The quadratic matrix inequality in singular H Control with state
90/294	<i>I.J.B.F. Adan</i> <i>J. Wessels</i> <i>W.H.M. Zijm</i>	Analysis of the symmetric shortest queue problem
90/295	<i>A. van den Boom</i> <i>R. Kool</i>	The Benelux Working Group on Software: Activities and software developments
90/296	<i>F.W. Steutel</i> <i>J.G.F. Thiemann</i>	The Gamma process and the Poisson distribution
90/297	<i>C. Withagen</i>	A differential game between government and firms: comments
90/298	<i>B. Veltman</i> <i>B.J. Lageweg</i> <i>J.K. Lenstra</i>	Multiprocessor scheduling with communication delays
90/299	<i>L.J.G. Langenhoff</i> <i>W.H.M. Zijm</i>	An analytical theory of multi-echelon production/distribution systems
90/300	<i>I.J.B.F. Adan</i> <i>J. Wessels</i> <i>W.H.M. Zijm</i>	Analysis of the asymmetric shortest queue problem
90/301	<i>Jan van Geldrop</i> <i>Shou Jilin</i> <i>Cees Withagen</i>	Existence of general equilibria in economies with natural exhaustible resources and an infinite horizon
91/303	<i>Henk Zijm</i>	Operational control of automated PCB assembly lines
91/304	<i>Ivo Adan</i> <i>Jaap Wessels</i> <i>Henk Zijm</i>	Flexible assembly and shortest queue problems
91/305	<i>D. Furth</i> <i>C. Withagen</i>	Mathematical economics, an introduction
91/306	<i>J. van Geldrop</i> <i>C. Withagen</i>	On the Negishi-approach to dynamic economic systems
91/307	<i>F.W. Steutel</i>	Counterexamples to Robertson's conjecture
91/308	<i>H.N. Linssen</i>	A table for solving the Behrens - Fisher problem
91/309	<i>Patrick Zwietering</i> <i>Emile Aarts</i>	Parallel Boltzmann machines: a mathematical model
91/310	<i>P.J. Zwietering</i> <i>M.J.A.L. van Kraaij</i> <i>E.H.L. Aarts</i> <i>J. Wessels</i>	Neural networks and production planning
91/311	<i>K. van Harn</i> <i>F.W. Steutel</i>	On a characterization of the exponential distribution
91/312	<i>P.J. Zwietering</i> <i>E.H.L. Aarts</i> <i>J. Wessels</i>	The design and complexity of exact multilayered perceptrons
91/313	<i>H.J.C. Huijberts</i> <i>A.J. van der Schaft</i>	Input-output decoupling with stability for Hamiltonian systems
92/314	<i>Charles van Marrewijk</i> <i>Casper G. de Vries</i> <i>Cees Withagen</i>	Optimal localized production experience and schooling
92/315	<i>I.J.B.F. Adan</i> <i>J. Wessels</i> <i>W.H.M. Zijm</i>	Analysis of the asymmetric shortest queue problem with threshold jockeying
92/316	<i>G.J. van Houtum</i> <i>W.H.M. Zijm</i>	Computational procedures for stochastic multi-echelon production systems
92/317	<i>H.J.C. Huijberts</i> <i>H. Nijmeijer</i> <i>L.L.M. van der Wegen</i>	Dynamic disturbance decoupling for nonlinear systems

92/318	<i>H.J.C. Huijberts</i>	A nonregular solution of the nonlinear dynamic disturbance decoupling problem with an application to a complete solution of the nonlinear model matching problem
92/319	<i>H.J.C. Huijberts H. Nijmeijer L.L.M. van der Wegen</i>	Minimality of dynamic input-output decoupling for nonlinear systems
92/320	<i>P.J.M. van Laarhoven E.H.L. Aarts J.K. Lenstra</i>	Job shop scheduling by simulated annealing
92/321	<i>M.C. van der Heijden A.G. de Kok</i>	Customer waiting times in an (R,S) inventory system with compound Poisson demand
92/322	<i>M.W.I. van Kraaij W.Z. Venema J. Wessels</i>	Support for problem solving in manpower planning problems
92/323	<i>H.J.C. Huijberts H. Nijmeijer</i>	Local nonlinear model matching: from linearity to nonlinearity
92/324	<i>A. Wagelmans S. van Hoesel A. Kolen</i>	Economic lot sizing: an $O(n \log n)$ algorithm that runs in linear time in the Wagner-Whitin case
92/325	<i>D. Levhari C. Withagen</i>	Optimal management of the growth potential of renewable resources
92/326	<i>F. Groot</i>	Note on the open-loop Von Stackelberg equilibrium in the cartel versus fringe
	<i>C. Withagen A. de Zeeuw</i>	model
92/327	<i>P.J. Zwitering E.H.L. Aarts J. Wessels</i>	Exact classification with two-layered perceptrons
92/328	<i>F.W. Steutel P. van der Laan J. Wessels R. Doornbos C.P.M. van Hoesel C.A.J. Hurkens J.K. Lenstra M.W.P. Savelsbergh A.G. de Kok J. Wessels</i>	On the number of first-rate statisticians Beslissen met statistische selectiemethoden De drie gratiën of: de schone Helena en haar lelijke zusters? Een voorbeeld van een weinig aannemelijk M-L schatter Productieplanning in praktijk en theorie
93/329		Lof der schoonheid Decision systems: The relation between problem specification and mathematical analysis
93/330	<i>H. Huijberts C. Withagen</i>	Local asymptotic stability of optimal steady states
93/331	<i>K. van Ham F.W. Steutel</i>	Stability equations for processes with stationary independent increments using branching processes and Poisson mixtures
93/332	<i>I.J.B.F. Adan J. Wessels W.H.M. Zijm</i>	The effect of varying routing probability in two parallel queues with dynamic routing under a threshold-type scheduling
93/333	<i>I.J.B.F. Adan J. Wessels W.H.M. Zijm</i>	Matrix-geometric analysis of the shortest queue problem with threshold jockeying
93/334	<i>J.H. van Geldrop C.A.A.M. Withagen</i>	General equilibrium and international trade with exhaustible resources
93/335	<i>G.J. van Houtum I.J.B.F. Adan J. Wessels W.H.M. Zijm</i>	The compensation approach for three or more dimensional random walks
93/336	<i>I.J.B.F. Adan J. Wessels</i>	Product forms as a solution base for queueing systems
93/337	<i>G. Kindervater J.K. Lenstra M. Savelsbergh</i>	Sequential and parallel local search for the time-constrained traveling salesman problem
93/338	<i>E.L. Lawler J.K. Lenstra A.H.G. Rinnooy Kan D.B. Shmoys</i>	Sequencing and scheduling: algorithms and complexity

93/339	I.J.B.F. Adan G.J. van Houtum J. Wessels W.H.M. Zijm	A compensation procedure for multiprogramming queues
93/340	I.J.B.F. Adan J. Wessels W.H.M. Zijm	Analysing multiprogramming queues by generating functions
93/341	P.J. Zwietering E.H.L. Aarts J. Wessels	The construction of minimal multilayered perceptrons: A case study for sorting
93/342	E.H.L. Aarts J. Wessels P.J. Zwietering	The applicability of neural nets for decision support
93/343	L.J.J. van der Bruggen J.K. Lenstra P.C. Schuur	Variable-depth search for the single-vehicle pickup and delivery problem with time windows
93/344	I.J.B.F. Adan J. Wessels W.H.M. Zijm	A compensation approach for two-dimensional Markov processes
93/345	O.J. Boxma G.J. van Houtum	The compensation approach applied to a 2 x 2 switch
94/346	P.J. Zwietering E.H.L. Aarts J. Wessels	The minimal number of layers of a perceptron that sorts
94/347	I. Adan G.J. van Houtum J. van der Wal	Upper and lower bounds for the waiting time in the symmetric shortest queue system
94/348	C.P.M. van Hoesel A.P.M. Wagelmans L.A. Wolsey	Polyhedral characterization of the economic lot-sizing problem with start-up costs
94/349	E.H.L. Aarts P.J.M. van Laarhoven J.K. Lenstra N.L.J. Ulder	A computational study of local search algorithms for job shop scheduling
94/350	J. Korst E. Aarts J.K. Lenstra J. Wessels	Periodic assignment and graph colouring
94/351	H. Zijm G.J. van Houtum	On multi-stage production/inventory systems under stochastic demand
94/352	Jan H. van Geldrop Cees A.A.M. Withagen	General equilibrium in an economy with exhaustible resources and an unbounded horizon
94/353	C. Withagen	Pollution and exhaustibility of fossil fuels
94/354	J.J.A.M. Brands F.W. Steutel R.J.G. Wilms	On the number of maxima in a discrete sample
94/355	J.A. Hoogeveen J.K. Lenstra B. Veltman	Tree, four, five, six, or the complexity of scheduling with communication delays
94/356	M.J.A. van Eenige J. van der Wal	A non-homogeneous MIPh1 queueing system subject to breakdowns
95/357	J.L. van den Berg E.B. Diks J.A.C. Resing J. van der Wal	The change of traffic characteristics in ATM networks
95/358	E. Balas J.K. Lenstra A. Vazacopoulos	The one-machine problem with delayed precedence constraints and its use in job shop scheduling
95/359	L. Fortuin T. de Kok I. Adan	Voorraadbeheer volgens model
95/360	C. Withagen	Pollution, abatement and balanced growth

95/361	E.E.M. van Berkum H.N. Linssen D.A. Overdijk	The reduction and ancillary classification of observations
95/362	J.K. Lenstra D.B. Shmoys	Computing near-optimal schedules
95/363	A.W.J. Kolen J.K. Lenstra	Combinatorics in operations research
95/364	H.P. Stehouwer E.H.L. Aarts J. Wessels	Multi-layered perceptrons for on-line lot sizing (extended abstract)
95/365	F.W. Steutel	Probabilistic methods in problems of applied analysis
95/366	K. van Harn F.W. Steutel	Infinite divisibility and the waiting-time paradox
95/367	R.F.M. Schapers R.P.E. Pauwels J.Th.M. Wijnen A.W.G.B. Smeets F.T. Bosman	Morphometric grading of transitional cell carcinoma of the urinary bladder
95/368	J.H.J. Einmahl	A Bahadur-Kiefer theorem beyond the largest observation
95/369	J. Beirlant J.H.J. Einmahl	Asymptotic confidence intervals for the length of the short under random censoring
95/370	J.A. Hoogeveen S.L. van de Velde	Minimizing total completion time and maximum cost simultaneously is solvable in polynomial time
95/371	H.L. Trentelman A.A. Stoorvogel	Sampled-data and discrete-time H_2 optimal control
95/372	A.A. Stoorvogel	Nonlinear L_1 optimal controllers for linear systems
95/373	A. Saberi Z. Lin A.A. Stoorvogel	H_2 almost disturbance decoupling problem with internal stability
95/374	A.A. Stoorvogel	The H_∞ control problem with zeros on the boundary of the stability domain
95/375	A. Saberi P. Sannuti A.A. Stoorvogel	H_2 optimal controllers with measurement feedback for continuous-time system - flexibility in closed-loop pole placement
95/376	R. Mantri A. Saberi Z. Lin A.A. Stoorvogel	Output regulation for linear discrete-time systems subject to input saturation
95/377	A.A. Stoorvogel A. Saberi	Continuity properties of solutions to H_2 and H_∞ Riccati equations
95/378	S. Weiland A.A. Stoorvogel	Rational representations of $e_2[0, \infty)$ systems
95/379	A. Di Bucchianico	Representations of stochastic processes
95/380	P. van der Laan	Subset selection for exponential populations: determination of the selection constant
95/381	J.L. van den Berg C.H.T.M. van der Heijden J.A.C. Resing J. Wessels	Adaptive traffic control in ATM networks
95/382	J.A. Hoogeveen S.L. van de Velde	Stronger Lagrangian bounds by use of slack variables: applications to machine scheduling problems
95/383	J.A. Hoogeveen S.L. van de Velde	Formulating a scheduling problem with almost identical jobs by using positional completion times
95/384	J.H.J. Einmahl F.H. Ruymgaart	Tail processes under heavy random censorship with applications
96/385	Z. Lin A.A. Stoorvogel A. Saberi	Output regulation for linear systems subject to input saturation
96/386	I. Adan M. van Eenige J. Resing	Fitting discrete distributions on the first two moments
96/387	J.H.J. Einmahl	Extension to higher dimensions of the Jaeschke-Eicker result on the standardized empirical process