

On the spectral characterization of mixed extensions of P3

Haemers, Willem H.; Sorgun, Sezer; Topcu, Hatice

Document version:

Publisher's PDF, also known as Version of record

Publication date:

2018

[Link to publication](#)

Citation for published version (APA):

Haemers, W. H., Sorgun, S., & Topcu, H. (2018). On the spectral characterization of mixed extensions of P3. (arXiv; Vol. 1810.12615). Ithaca: Cornell University Library.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright, please contact us providing details, and we will remove access to the work immediately and investigate your claim.

On the spectral characterization of mixed extensions of P_3

Willem H. Haemers*

haemers@uvt.nl

Sezer Sorgun†

srgnrzs@gmail.com

Hatice Topcu †

haticekamittopcu@gmail.com

Abstract

A mixed extension of a graph G is a graph H obtained from G by replacing each vertex of G by a clique or a coclique, whilst two vertices in H corresponding to distinct vertices x and y of G are adjacent whenever x and y are adjacent in G . If G is the path P_3 , then H has at most three adjacency eigenvalues unequal to 0 and -1 . Recently, the first author classified the graphs with the mentioned eigenvalue property. Using this classification we investigate mixed extension of P_3 on being determined by the adjacency spectrum. We present several cospectral families, and with the help of a computer we find all graphs on at most 25 vertices that are cospectral with a mixed extension of P_3 .

1 Introduction

Characterizations of graphs by means of the spectrum of the adjacency matrix is a well-studied subject. Although it is conjectured that almost all graphs are determined by the spectrum of the adjacency matrix, there are still relatively few graphs known to be determined by its spectrum. The reason is that in general this property is hard to prove. Some

*Dept. of Econometrics and O.R., Tilburg University, The Netherlands

†Dept. of Mathematics, Nevşehir Hacı Bektaş Veli University, Turkey

of these proofs are based on the classification of graphs with certain spectral properties, such as classifications in terms of the smallest eigenvalue. Recently the first author classified the graphs with all but at most three eigenvalues equal to 0 or -1 ; see [4]. Here this classification is applied to mixed extensions of the path P_3 (see next section), which have the mentioned spectral property. We give the characteristic polynomial of all graphs in the classification, and completely determine the mixed extensions of P_3 which are determined by the spectrum on at most 25 vertices. Also we present several infinite families of cospectral graphs with all but three eigenvalues equal to 0 or -1 .

2 Mixed extensions of P_3

Let G be a graph with vertex set $V(G) = \{1, \dots, m\}$ and let V_1, \dots, V_m be mutually disjoint nonempty finite sets. A graph H with vertex set $V(H) = V_1 \cup \dots \cup V_m$ is defined as follows. For each $i \in \{1, \dots, m\}$, all vertices of V_i are either mutually adjacent (form a clique), or mutually nonadjacent (form a coclique). For any $u \in V_i$ and $v \in V_j$ ($i \neq j$) $\{u, v\}$ is an edge in H if and only if $\{i, j\}$ is an edge in G . The graph H is called a *mixed extension* of G . A mixed extension is represented by an m -tuple (t_1, \dots, t_m) of nonzero integers, where $t_i > 0$ indicates that V_i is a clique of order t_i and $t_i < 0$ means that V_i is a coclique of order $-t_i$. We refer to [4] for basic results on mixed extensions, and to [1] or [3] for graph spectra.

Suppose H is a mixed extension of the path P_3 of type (t_1, t_2, t_3) . Then the adjacency matrix of H admits the following structure. (As usual, J is an all-ones matrix, J_n is the $n \times n$ all-ones matrix, and I_n is the identity matrix of order n .)

$$A = \begin{bmatrix} \varepsilon_1(J_{|t_1|} - I_{|t_1|}) & J & O \\ J & \varepsilon_2(J_{|t_2|} - I_{|t_2|}) & J \\ O & J & \varepsilon_3(J_{|t_3|} - I_{|t_3|}) \end{bmatrix},$$

where $\varepsilon_i = 1$ if $t_i > 1$ and $\varepsilon_i = 0$ otherwise ($i = 1, 2, 3$).

The given partition of A is equitable, therefore A has two kinds of eigenvalues: the ones that have eigenvectors in the span V of the characteristic vectors of the partition, and those whose eigenvectors are orthogonal to V . The first kind coincide with the eigenvalues of the quotient matrix

$$Q = \begin{bmatrix} \varepsilon_1(|t_1| - 1) & |t_2| & 0 \\ |t_1| & \varepsilon_2(|t_2| - 1) & |t_3| \\ 0 & |t_2| & \varepsilon_3(|t_3| - 1) \end{bmatrix}.$$

The second kind of eigenvalues of H remain eigenvalues if we subtract an all-one block from each nonzero block of A . So these eigenvalues are also eigenvalues of

$$B = \begin{bmatrix} -\varepsilon_1 I_{|t_1|} & O & O \\ O & -\varepsilon_2 I_{|t_2|} & O \\ O & O & -\varepsilon_3 I_{|t_3|} \end{bmatrix},$$

which are clearly all equal to 0 or -1 . So, if $n = |t_1| + |t_2| + |t_3|$ is the order of H , and $q(x) = x^3 - bx^2 - cx + d$ is the characteristic polynomial of Q , then the characteristic polynomial of A equals $p(x) = q(x)(x+1)^b x^{n-b-3}$. Indeed, the exponent of $(x+1)$ equals b , because the coefficient of x^{n-1} in $p(x)$ equals $\text{trace}(A) = 0$. Note that $d = -\det(Q) \geq 0$, and $\det(Q) = 0$ if and only if $\varepsilon_1 = \varepsilon_3 = 0$ that is, both end vertices of P_3 are replaced by a coclique. If this is the case then we call it an *improper* mixed extension of P_3 , because it is in fact a mixed extension of P_2 . If $\det(Q) \neq 0$, then the mixed extension is *proper*.

Proposition 2.1. *A proper mixed extension of the path P_3 has exactly two positive eigenvalues and one eigenvalue smaller than -1 .*

Proof. The quotient matrix Q has at least one positive eigenvalue, and $\det(Q) < 0$ gives that Q (and A) has exactly two positive eigenvalues. It is well-known that a graph with smallest adjacency eigenvalue at least -1 is the disjoint union of complete graphs. Therefore the smallest eigenvalue of A (and Q) is less than -1 . \square

3 Spectral characterizations

Some known results on spectral characterizations of graphs deal with special cases of mixed extensions of P_3 . This includes the pineapple graphs (type $(p, 1, -q)$, $p, q > 0$), and the complete graphs from which the edges of a complete bipartite subgraph are deleted (type (p, q, r) , $p, q, r > 0$). The latter graphs are determined by their spectrum for all $p, q, r > 0$; see [2]. For the pineapple graphs it is known for which p and q the graphs are determined by its spectrum. For the precise conditions on p and q we refer to [6]. It follows that among the connected graphs the pineapple graph is determined by its spectrum.

An improper mixed extension of P_3 is either a complete bipartite graph $K_{p,q}$ ($p \geq 2, q \geq 1$), or a complete split graph $CS_{p,q}$ ($p, q \geq 2$), which is a complete graph K_{p+q} from which the edges of a complete subgraph K_q are deleted. For these classes of graphs the spectral characterization is straightforward and known. The complete split graph $CS_{p,q}$ is determined by its spectrum for all $p, q \geq 2$. The complete bipartite graph $K_{p,q}$ is not determined by its adjacency spectrum if and only if pq has a divisor r strictly between p

	b	c	d
$(-p, -q, r)$	$r - 1$	$pq + qr$	$pq(r - 1)$
$(-p, q, r)$	$q + r - 2$	$q + r + pq - 1$	$pq(r - 1)$
$(p, -q, r)$	$p + r - 2$	$q(p + r) + (p - 1)(1 - r)$	$qr(p - 1) + pq(r - 1)$
(p, q, r)	$p + q + r - 3$	$2q + 2r + 2p - pr - 3$	$qpr + pr - p - q - r + 1$
$(-2, q, r, -2)$	$q + r - 1$	$2q + 2r$	$4qr$
$(-3, q, -2, s)$	$q + s - 1$	$2s + 5q - qs$	$6qs$
$(-2, -2, -3, s)$	s	$2s + 10$	$12s$
$(5, 2, -r, 4)$	8	$6r - 9$	$34r + 18$
$(4, 2, -r, 6)$	9	$8r - 15$	$40r + 25$
$(7, 2, -r, 3)$	9	$5r - 6$	$37r + 16$
$(3, 3, -r, 6)$	9	$9r - 15$	$45r + 25$
$(4, 3, -r, 3)^*$	7	$6r - 4$	$30r + 12$
$(7, 3, -r, 2)$	9	$5r + 1$	$37r + 9$
$(3, 4, -r, 4)$	8	$8r - 9$	$40r + 18$
$(3, 6, -r, 3)$	9	$9r - 6$	$45r + 16$
$(4, 6, -r, 2)^*$	9	$8r + 1$	$40r + 9$
$(5, 4, -r, 2)^*$	9	$6r + 1$	$34r + 8$
$(2, 2, 2, 7)$	9	1	65
$(2, 2, 6, 3)$	9	9	73
$(2, 2, 3, 4)$	7	5	51
$(2, 3, 2, 5)$	8	1	68
$(2, 3, 4, 3)$	8	7	74
$(2, 5, 2, 4)$	9	1	89
$(3, 2, 2, 3)$	6	3	40
$(2, 5, 3, 3)$	9	6	94
$(1, p, -q, r, 1)$	$p + r - 1$	$(q + 1)(p + r) - pr$	$pr(2q + 1)$
$K_p + K_{q,r}$	$p - 1$	qr	$qr(p - 1)$
$K_p + CS_{q,r}$	$p + q - 2$	$qr - (p - 1)(q - 1)$	$qr(p - 1)$

Table 1: Coefficients b, c, d of the characteristic polynomials of graphs in \mathcal{G}''

and q . Then $K_{r,pq/r}$ extended with $p+q-r-pq/r$ isolated vertices is cospectral with $K_{p,q}$. From now on we restrict to proper mixed extensions of P_3 .

We define \mathcal{G} to be the set of graphs with all but at most three adjacency eigenvalues equal to -1 or 0 . Note that a graph G in \mathcal{G} remains in \mathcal{G} if isolated vertices are added or deleted. Therefore, for the classification, we may restrict to the set of graphs in \mathcal{G} with no isolated vertices. By \mathcal{G}'' we denote the class of graphs in \mathcal{G} with no isolated vertices, with two positive eigenvalues and with one eigenvalue less than -1 . By Proposition 2.1 we know that a proper mixed extension of P_3 is in \mathcal{G}'' . The characterization of graphs in \mathcal{G} , mentioned in the introduction, when restricted to \mathcal{G}'' , is given by the following two theorems.

Theorem 3.1. *A disconnected graph H belongs to \mathcal{G}'' , if and only if H is one of the following.*

- (i) $K_p + K_{q,r}$ with $p, q, r \geq 2$,
- (ii) $K_p + CS_{q,r}$ with $p, q \geq 2, q \geq 1$.

Theorem 3.2. *A connected graph H belongs to \mathcal{G}'' if and only if H is one of the following.*

- (i) A mixed extension of P_3 of type $(-p, -q, r)$; $(-p, q, r)$; $(p, -q, r)$, or (p, q, r) with $p, q \geq 1$ and $r \geq 2$,
- (ii) a mixed extension of P_4 of type $(-2, q, r, -2)$; $(-3, q, -2, s)$, or $(-2, -2, -2, s)$ with $q, r, s \geq 1$,
- (iii) a mixed extension of P_4 of type $(p, q, -r, s)$ with $r \geq 1$ and $(p, q, s) \in \{(5, 2, 4), (4, 2, 6), (7, 2, 3), (3, 3, 6), (4, 3, 3), (7, 3, 2), (3, 4, 4), (3, 6, 3), (4, 6, 2), (5, 4, 2)\}$,
- (iv) a mixed extension of P_4 of type (p, q, r, s) , with $(p, q, r, s) \in \{(2, 2, 2, 7), (2, 2, 6, 3), (2, 2, 3, 4), (2, 3, 2, 5), (2, 3, 4, 3), (2, 5, 2, 4), (2, 5, 3, 3), (3, 2, 2, 3)\}$,
- (v) a mixed extension of P_5 of type $(1, p, -q, r, 1)$ with $p, q, r \geq 1$.

The characteristic polynomial $p(x)$ of a graph in \mathcal{G}'' with n vertices can be written as:

$$p(x) = (x^3 - bx^2 - cx + d)(x + 1)^b x^{n-b-3}, \text{ with } b \geq 0 \text{ and } d > 0$$

(note that the multiplicity of the eigenvalue -1 equals b , because $\text{trace}(A) = 0$). So the nonzero part of the spectrum of a graph in \mathcal{G}'' is determined by the coefficients b, c and d . Table 1 gives these coefficients for each type of the above classification. If two graphs in the classification have the same b, c and d , then the nonzero part of the spectrum is the same. We will call such a pair *pseudo-cospectral*. If two pseudo-cospectral graphs have different order, we can extend the smaller graph (or both graphs) with some isolated vertices, so that the two graphs become cospectral. Therefore Table 1 gives all information needed to decide which proper mixed extensions of P_3 are determined by their spectrum. Nevertheless, it is far too complicated to give a general result. Therefore we present some special cases.

Theorem 3.3. [2] *Suppose H is a mixed extension of P_3 of order n and type (p, q, r) with $p, q, r \geq 1$. Then H is determined by the spectrum of its adjacency matrix.*

Proof. From Table 1 we see that for this case the coefficient b equals $p + q + r - 3 = n - 3$. For every graph H' of order n' pseudo-cospectral with H , which belongs to one of the other types, we have $b < n' - 3$. So H' has more vertices than H , and we cannot obtain a graph cospectral with H by adding isolated vertices to H' . If H' is a mixed extension of P_3 of type (p', q', r') with $p', q', r' \geq 1$, and H' is cospectral with H , then H and H' have the coefficients b, c and d , which implies $p = p', q = q'$, and $r = r'$. \square

The following results follow straightforwardly from Table 1.

Proposition 3.4. *The following types of mixed extensions of P_3 are pseudo-cospectral with a non-isomorphic graph in \mathcal{G}'' .*

- (i) *Type $(p, -q, p)$ with $K_p + CS_{p,2q}$ where $p \geq 2, q \geq 1$,*
- (ii) *type $(p, (p-1)(p-2), p)$ with $K_{p(p-1)} + K_{p-1,p-1}$, where $p \geq 3$,*
- (iii) *type (p, q, p) with $K_p + CS_{p+q-1,r}$ where $r = 1 + pq/(p+q-1)$ is an integer, and $p \geq 2, q \geq 1$.*

Notice that this proposition does not give any graph which is cospectral and non-isomorphic with a mixed extension of P_3 , because in all three cases the second graph has more vertices than the first one. For the pineapple graph K_p^q , which is a mixed extension of P_3 of type $(p-1, 1, -q)$, we find (see [5]).

Proposition 3.5. *The pineapple graph $K_{2p}^{p^2}$ is cospectral with the mixed extension of P_3 of type $(p, -p, p)$ extended with $p(p-1)$ isolated vertices.*

See [6] for more examples of graphs non-isomorphic but cospectral with K_p^q . As remarked before, there exists no connected example. However, there do exist connected non-isomorphic cospectral mixed extensions of P_3 .

Proposition 3.6. *The mixed extensions of P_3 of types $(p, -q, q(2q-p-1)/(q-p))$ and $(-q, 2q-1, p(2q-p-1)/(q-p))$ are cospectral whenever $q(2q-p-1)/(q-p)$ is a positive integer, and $p, q \geq 1$.*

Note that there are infinitely many values of p and q for which the above fraction is integral. For example if $p = 1$ or $p = q - 1$.

4 Enumeration

Using Table 1, we generated by computer (using Maple) a list of all non-isomorphic graphs in \mathcal{G}'' on at most 25 vertices. The list contains almost 10000 graphs. We ordered the graphs alphabetically with respect to the coefficients b, c, d . Then the pseudo-cospectral graphs in \mathcal{G}'' become consecutive items in the table with the same b, c, d . Since the list is very long we only consider the mixed extensions of P_3 that have at least one non-isomorphic pseudo-cospectral mate in the list. By use of the shortened list we found the pseudo-cospectral examples of Propositions 3.4 and 3.6. Next we deleted the cases given in Proposition 3.4, 3.5 and 3.4 from the list. The final list is given in Table 2. Thus this table together with Propositions 3.4 to 3.6 give all mixed extensions of P_3 of order $n \leq 25$ for which there exist at least one pseudo-cospectral graph in \mathcal{G}'' of order at most 25. Since Proposition 3.4 gives no graphs cospectral with a mixed extension of P_3 , we can conclude the following:

Theorem 4.1. *Suppose H is a proper mixed extension of P_3 of order $n \leq 25$. Then H is determined by the spectrum of the adjacency matrix if and only if H is not one of the graphs given in Propositions 3.5 or 3.6, and every graph in Table 2 pseudo-cospectral with H has more vertices than H .*

References

- [1] A.E. Brouwer and W.H. Haemers, Spectra of Graphs, Springer, 2012.
- [2] M. Cámara and W.H. Haemers, Spectral characterizations of almost complete graphs, Discrete Appl. Math. 176 (2014), 19–23.
- [3] Dragoš Cvetković, Peter Rowlinson, Slobodan Simić, An Introduction to the Theory of Graph Spectra, London Mathematical Society Student Texts, 2009.
- [4] W.H. Haemers, Spectral characterization of mixed extensions of small graphs, Discrete Mathematics (to appear); also: <https://arxiv.org/abs/1712.01749>.
- [5] H. Topcu, S. Sorgun, W.H. Haemers, On the spectral characterization of pineapple graphs, Linear Algebra and its Applications 507 (2016), 267-273.
- [6] H. Topcu, S. Sorgun, W. H. Haemers, The graphs cospectral with the pineapple graph, Discrete Applied Mathematics (2018), <https://doi.org/10.1016/j.dam.2018.10.002>.

Table 2: Graphs in \mathcal{G}'' of order $n \leq 25$ pseudo-cospectral with a mixed extension of P_3

b	c	d	Type	p	q	r	s	n
2	10	14	$(-p, q, r)$	7	1	3		11
2	10	14	$(1, p, -q, r, 1)$	1	3	2		8
3	8	12	$(-p, q, r)$	4	1	4		9
3	8	12	$(-p, q, r, -s)$	2	1	3	2	8
3	9	15	$(-p, q, r)$	5	1	4		10
3	9	15	$(1, p, -q, r, 1)$	1	2	3		8
3	10	12	$(-p, q, r)$	3	2	3		8
3	10	12	$K_p + CS_{q,r}$	2	3	4		9
3	14	18	$(-p, -q, r)$	3	2	4		9
3	14	18	$(-p, q, -r, s)$	3	3	2	1	9
3	16	24	$(-p, -q, r)$	4	2	4		10
3	16	24	$(-p, q, r)$	6	2	3		11
3	16	36	$(-p, q, r)$	12	1	4		17
3	16	36	$(-p, -q, -r, s)$	2	2	3	3	10
3	16	36	$(1, p, -q, r, 1)$	2	4	2		10
3	16	36	$K_p + CS_{q,r}$	3	2	9		14
3	18	28	$(-p, q, r)$	7	2	3		12
3	18	28	$(p, -q, r)$	2	4	3		9
3	28	60	$(-p, -q, r)$	10	2	4		16
3	28	60	$(1, p, -q, r, 1)$	2	7	2		13
3	28	60	$K_p + CS_{q,r}$	3	2	15		20
3	40	72	$(-p, -q, r)$	6	4	4		14
3	40	72	$(-p, q, r)$	18	2	3		23
3	52	108	$(-p, -q, r)$	9	4	4		17
3	52	108	$(1, p, -q, r, 1)$	2	13	2		19
4	9	12	$(-p, q, r)$	2	2	4		8
4	9	12	$K_p + CS_{q,r}$	2	4	3		9
4	15	30	$(-p, q, r)$	5	2	4		11
4	15	30	$(p, -q, r)$	2	3	4		9
4	21	24	$(-p, -q, r)$	2	3	5		10
4	21	24	$K_p + CS_{q,r}$	2	4	6		12
4	23	54	$(-p, q, r)$	9	2	4		15
4	23	54	$K_p + CS_{q,r}$	3	3	9		15
4	39	102	$(-p, q, r)$	17	2	4		23
4	39	102	$(1, p, -q, r, 1)$	2	8	3		15
5	14	40	$(-p, q, r)$	8	1	6		15
5	14	40	$K_p + CS_{q,r}$	3	4	5		12
5	16	20	$(-p, -q, r)$	2	2	6		10
5	16	20	$K_p + CS_{q,r}$	2	5	4		11
5	18	48	$(-p, q, r)$	6	2	5		13
5	18	48	$K_p + CS_{q,r}$	3	4	6		13
5	24	90	$(-p, q, r)$	18	1	6		25
5	24	90	$K_p + CS_{q,r}$	4	3	10		17
5	28	88	$(-p, q, r)$	11	2	5		18
5	28	88	$(1, p, -q, r, 1)$	2	5	4		13
5	30	72	$(-p, q, r)$	8	3	4		15
5	30	72	$K_p + CS_{q,r}$	3	4	9		16
5	36	90	$(-p, -q, r)$	6	3	6		15
5	36	90	$(-p, q, r)$	10	3	4		17
5	36	120	$(-p, -q, r)$	12	2	6		20
5	36	120	$(-p, q, r)$	15	2	5		22

b	c	d	Type	p	q	r	s	n
5	42	144	$(-p, q, r)$	18	2	5		25
5	42	144	$K_p + CS_{q,r}$	4	3	16		23
5	57	153	$(-p, q, r)$	17	3	4		24
5	57	153	$(p, -q, r)$	3	9	4		16
5	70	200	$(-p, -q, r)$	8	5	6		19
5	70	200	$(1, p, -q, r, 1)$	2	12	4		20
6	17	50	$(-p, q, r)$	5	2	6		13
6	17	50	$K_p + CS_{q,r}$	3	5	5		13
6	19	48	$(-p, q, r)$	4	3	5		12
6	19	48	$(p, -q, r)$	2	3	6		11
6	19	60	$(-p, q, r)$	6	2	6		14
6	19	60	$(-p, q, -r, s)$	3	5	2	2	12
6	22	60	$(-p, q, r)$	5	3	5		13
6	22	60	$K_p + CS_{q,r}$	3	5	6		14
6	25	90	$(-p, q, r)$	9	2	6		17
6	25	90	$(1, p, -q, r, 1)$	2	4	5		13
6	28	84	$(-p, -q, r)$	7	2	7		16
6	28	84	$(-p, q, r)$	7	3	5		15
6	31	120	$(p, -q, r)$	4	5	4		13
6	31	120	$(-p, q, r)$	12	2	6		20
6	40	132	$(-p, q, r)$	11	3	5		19
6	40	132	$(p, -q, r)$	3	6	5		14
6	44	180	$(-p, -q, r)$	15	2	7		24
6	44	180	$(1, p, -q, r, 1)$	3	7	4		16
6	50	90	$(-p, -q, r)$	3	5	7		15
6	50	90	$(1, p, -q, r, 1)$	1	7	6		16
6	51	180	$(-p, -q, r)$	10	3	7		20
6	51	180	$K_p + CS_{q,r}$	4	4	15		23
6	55	192	$(-p, q, r)$	16	3	5		24
6	55	192	$(p, -q, r)$	4	8	4		16
7	16	48	$(-p, q, r)$	4	2	7		13
7	16	48	$(-p, q, r, -s)$	2	2	6	2	12
7	20	60	$(-p, q, r)$	4	3	6		13
7	20	60	$K_p + CS_{q,r}$	3	6	5		14
7	20	84	$(-p, q, r)$	12	1	8		21
7	20	84	$(1, p, -q, r, 1)$	2	3	6		13
7	23	105	$(-p, q, r)$	15	1	8		24
7	23	105	$K_p + CS_{q,r}$	4	5	7		16
7	26	108	$(-p, q, r)$	9	2	7		18
7	26	108	$(p, -q, r)$	3	4	6		13
7	28	120	$(-p, q, r)$	10	2	7		19
7	28	120	$K_p + CS_{q,r}$	4	5	8		17
7	30	42	$(-p, -q, r)$	2	3	8		13
7	30	42	$(-p, q, -r, s)$	3	7	2	1	13
7	33	63	$(-p, -q, r)$	3	3	8		14
7	33	63	$(1, p, -q, r, 1)$	1	4	7		14
7	35	135	$(-p, q, r)$	9	3	6		18
7	35	135	$(p, -q, r)$	3	5	6		14
7	38	150	$(-p, q, r)$	10	3	6		19
7	38	150	$K_p + CS_{q,r}$	4	5	10		19
7	64	224	$(-p, -q, r)$	8	4	8		20
7	64	224	$(-p, q, r)$	14	4	5		23

b	c	d	Type	p	q	r	s	n
7	68	240	$(-p, q, r)$	15	4	5		24
7	68	240	$K_p + CS_{q,r}$	4	5	16		25
7	78	210	$(-p, -q, r)$	5	6	8		19
7	78	210	$(-p, q, r)$	14	5	4		23
8	8	64	$(p, -q, r)$	3	2	7		12
8	8	64	$(Kp + Kq, r)$	9	2	4		15
8	13	32	$(-p, q, r)$	4	1	9		14
8	13	32	(p, q, r)	2	6	3		11
8	16	56	$(-p, q, r)$	7	1	9		17
8	16	56	$K_p + CS_{q,r}$	3	7	4		14
8	18	72	$(-p, q, r)$	9	1	9		19
8	18	72	$(-p, q, r, -s)$	2	3	6	2	13
8	19	40	$(-p, q, r)$	2	5	5		12
8	19	40	$(1, p, -q, r, 1)$	1	2	8		13
8	27	126	$(-p, q, r)$	9	2	8		19
8	27	126	$K_p + CS_{q,r}$	4	6	7		17
8	33	144	$(-p, q, r)$	8	3	7		18
8	33	144	$K_p + CS_{q,r}$	4	6	8		18
8	38	160	$(-p, -q, r)$	10	2	9		21
8	38	160	$(p, -q, r)$	3	5	7		15
8	45	180	$(-p, q, r)$	9	4	6		19
8	45	180	$K_p + CS_{q,r}$	4	6	10		20
8	51	126	$(-p, q, r)$	7	6	4		17
8	51	126	$K_p + CS_{q,r}$	3	7	9		19
8	54	270	$(-p, q, r)$	15	3	7		25
8	54	270	$(1, p, -q, r, 1)$	3	7	6		18
8	57	240	$(-p, -q, r)$	10	3	9		22
8	57	240	$(-p, q, r)$	12	4	6		22
8	68	256	$(-p, -q, r)$	8	4	9		21
8	68	256	$(p, -q, r)$	3	8	7		18
8	128	448	$(-p, -q, r)$	7	8	9		24
8	128	448	$(p, -q, r)$	3	14	7		24
9	3	61	(p, q, r)	3	3	6		12
9	3	61	$(p, q, -r, s)$	3	6	1	3	13
9	4	90	$(p, -q, r)$	4	2	7		13
9	4	90	$(p, q, -r, s)$	7	2	2	3	14
9	9	49	(p, q, r)	2	4	6		12
9	9	49	$(p, q, -r, s)$	4	6	1	2	13
9	15	135	$(p, -q, r)$	4	3	7		14
9	15	135	$(Kp + Kq, r)$	10	3	5		18
9	18	64	$(-p, q, r)$	4	2	9		15
9	18	64	$K_p + CS_{q,r}$	3	8	4		15
9	24	112	$(-p, q, r)$	7	2	9		18
9	24	112	$(1, p, -q, r, 1)$	2	3	8		15
9	24	126	$(-p, q, r)$	14	1	10		25
9	24	126	$K_p + CS_{q,r}$	4	7	6		17
9	28	72	$(-p, q, r)$	3	6	5		14
9	28	72	$(-p, -q, r)$	4	2	10		16
9	31	63	$(-p, q, r)$	3	7	4		14
9	31	63	$(1, p, -q, r, 1)$	1	3	9		15
9	31	147	$(-p, q, r)$	7	3	8		18
9	31	147	$K_p + CS_{q,r}$	4	7	7		18

b	c	d	Type	p	q	r	s	n
9	34	96	$(-p, q, r)$	4	6	5		15
9	34	96	$K_p + CS_{q,r}$	3	8	6		17
9	34	144	$(-p, q, r)$	6	4	7		17
9	34	144	$(1, p, -q, r, 1)$	2	4	8		16
9	38	168	$(-p, q, r)$	7	4	7		18
9	38	168	$K_p + CS_{q,r}$	4	7	8		19
9	49	273	$(-p, q, r)$	13	3	8		24
9	49	273	$(1, p, -q, r, 1)$	3	6	7		18
9	64	216	$(-p, -q, r)$	6	4	10		20
9	64	216	$(-p, q, r)$	9	6	5		20
9	94	336	$(-p, q, r)$	14	6	5		25
9	94	336	$(1, p, -q, r, 1)$	2	10	8		22
10	20	72	$(-p, q, r)$	3	3	9		15
10	20	72	$K_p + CS_{q,r}$	3	9	4		16
10	27	144	$(-p, q, r)$	8	2	10		20
10	27	144	$K_p + CS_{q,r}$	4	8	6		18
10	29	90	$(-p, q, r)$	3	6	6		15
10	29	90	$K_p + CS_{q,r}$	3	9	5		17
10	32	168	$(-p, q, r)$	7	3	9		19
10	32	168	$(p, -q, r)$	3	4	9		16
10	35	168	$(-p, q, r)$	6	4	8		18
10	35	168	$K_p + CS_{q,r}$	4	8	7		19
10	59	240	$(-p, q, r)$	8	6	6		20
10	59	240	$K_p + CS_{q,r}$	4	8	10		22
10	63	364	$(-p, q, r)$	13	4	8		25
10	63	364	$(p, -q, r)$	4	7	8		19
10	66	330	$(-p, -q, r)$	11	3	11		25
10	66	330	$(-p, q, r)$	11	5	7		23
11	-12	108	$(1, p, -q, r, 1)$	6	1	6		15
11	-12	108	$K_p + CS_{q,r}$	7	6	3		16
11	24	108	$(-p, q, r)$	4	3	10		17
11	24	108	$(-p, q, r, -s)$	2	3	9	2	16
11	30	162	$(-p, q, r)$	6	3	10		19
11	30	162	$K_p + CS_{q,r}$	4	9	6		19
11	45	297	$(-p, q, r)$	11	3	10		24
11	45	297	$(1, p, -q, r, 1)$	3	5	9		19
11	48	216	$(-p, q, r)$	6	6	7		19
11	48	216	$K_p + CS_{q,r}$	4	9	8		21
11	52	320	$(-p, q, r)$	10	4	9		23
11	52	320	$K_p + CS_{q,r}$	5	8	10		23
12	1	132	$(p, -q, r)$	4	2	10		16
12	1	132	$K_p + CS_{q,r}$	12	2	6		20
12	19	66	$(-p, q, r)$	3	2	12		17
12	19	66	(p, q, r)	2	9	4		15
12	33	180	$(-p, q, r)$	5	4	10		19
12	33	180	$K_p + CS_{q,r}$	4	10	6		20
12	43	210	$(-p, q, r)$	5	6	8		19
12	43	210	$K_p + CS_{q,r}$	4	10	7		21
12	49	324	$(-p, q, r)$	9	4	10		23
12	49	324	$K_p + CS_{q,r}$	5	9	9		23
12	57	396	$(-p, q, r)$	11	4	10		25
12	57	396	$(p, -q, r)$	4	6	10		20

b	c	d	Type	p	q	r	s	n
12	58	360	$(-p, q, r)$	9	5	9		23
12	58	360	$K_p + CS_{q,r}$	5	9	10		24
13	-7	165	(p, q, r)	6	4	6		16
13	-7	165	$K_p + CS_{q,r}$	12	3	5		20
13	26	96	$(-p, q, r)$	2	6	9		17
13	26	96	$K_p + CS_{q,r}$	3	12	4		19
13	54	360	$(-p, q, r)$	8	5	10		23
13	54	360	$K_p + CS_{q,r}$	5	10	9		24
14	-17	176	(p, q, r)	6	3	8		17
14	-17	176	$K_p + CS_{q,r}$	12	4	4		20
14	39	216	$(-p, q, r)$	4	6	10		20
14	39	216	$K_p + CS_{q,r}$	4	12	6		22
15	32	192	$(-p, q, r)$	4	4	13		21
15	32	192	$(-p, q, r, -s)$	2	4	12	2	20
15	52	252	$(-p, q, r)$	4	9	8		21
15	52	252	$(1, p, -q, r, 1)$	2	4	14		22
16	-37	198	(p, q, r)	8	2	9		19
16	-37	198	$K_p + CS_{q,r}$	12	6	3		21
16	-28	288	$(p, -q, r)$	9	2	9		20
16	-28	288	$K_p + CS_{q,r}$	9	9	4		22

b	c	d	Type	p	q	r	s	n
16	-19	252	$(p, -q, r)$	6	2	12		20
16	-19	252	(p, q, r)	6	4	9		19
16	-10	252	(p, q, r)	5	5	9		19
16	-10	252	$K_p + CS_{q,r}$	15	3	6		24
16	31	210	$(-p, q, r)$	7	2	16		25
16	31	210	$K_p + CS_{q,r}$	4	14	5		23
16	38	210	$(-p, q, r)$	3	7	11		21
16	38	210	$(1, p, -q, r, 1)$	2	3	15		22
16	72	330	$(-p, q, r)$	5	11	7		23
16	72	330	$(1, p, -q, r, 1)$	2	5	15		24
17	33	225	$(-p, q, r)$	5	3	16		24
17	33	225	$K_p + CS_{q,r}$	4	15	5		24
17	36	270	$(-p, q, r)$	6	3	16		25
17	36	270	$(-p, q, -r, s)$	3	15	2	3	23
18	-17	420	$(-p, q, -r, s)$	3	5	2	14	24
18	-17	420	$K_p + CS_{q,r}$	8	12	5		25
18	35	240	$(-p, q, r)$	4	4	16		24
18	35	240	$K_p + CS_{q,r}$	4	16	5		25
19	40	300	$(-p, q, r)$	4	5	16		25
19	40	300	$(-p, q, r, -s)$	2	5	15	2	24