

Disconnected Vertex Sets and Equidistant Code Pairs

Haemers, W.H.

Publication date:
1996

[Link to publication](#)

Citation for published version (APA):
Haemers, W. H. (1996). *Disconnected Vertex Sets and Equidistant Code Pairs*. (FEW Research Memorandum; Vol. 728). Operations research.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright, please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Disconnected vertex sets and equidistant code pairs

Willem H. Haemers

Abstract

Let A and B be two disjoint vertex sets of a graph such that there are no edges between A and B . We give an upperbound for $|A| \cdot |B|$ in terms of eigenvalues of matrix representations of the graph. Applied to graphs in the Hamming association scheme it leads to bounds for equidistant code pairs, which improve and reprove some known results.

1 Introduction

Throughout G is a graph with vertex set V . Two disjoint subsets A and B of V are *disconnected* if there is no edge between A and B . We define $\Phi(G)$ to be the maximum of $\sqrt{|A| \cdot |B|}$ for disconnected sets A and B in G . Suppose V is the set of words of length n over an alphabet $\{1, \dots, q\}$ and define two words adjacent if their Hamming distance (i.e. the number of coordinates in which they differ) is not equal to a fixed $\delta \in \{1, \dots, n\}$. Then a pair of disconnected sets becomes an *equidistant code pair*.

The quantity $\Phi(G)$ has an application in information theory and leads to a lower bound for the two-way communication complexity of functions defined on $V \times V$ that are constant over the non-edges of G . About ten years ago this application caused some activity in the study of equidistant code pairs. The best result is due to Ahlswede [1], who gives the exact value of $\Phi(G)$ for $q = 2, 4$ and 5 , for every δ and n .

In this paper we will give a bound for $\Phi(G)$ in terms of the eigenvalues of a matrix associated with G . In case the complement of G is given by a relation of an association scheme the bound takes an easy form, which applied to the Hamming scheme leads to a bound for equidistant code pairs. This bound is not as accurate as Ahlswede's result, but it is more general and it turns out to be sharp for some values of q , n and δ , and for $q \rightarrow \infty$ for any fixed n and δ .

2 Disconnected vertex sets

Let $V = \{1, \dots, v\}$. We define $\mathcal{M}(G)$ to be the collection of symmetric $v \times v$ matrices M with all row and column sums equal to 1, such that $(M)_{i,j} = 0$ if i and j are distinct non-adjacent vertices of G . Let $\lambda_1(M), \dots, \lambda_v(M)$ denote the eigenvalues of a matrix $M \in \mathcal{M}(G)$, such that $\lambda_1(M)$ has eigenvector $\mathbf{1}$ (the all-one vector), so $\lambda_1(M) = 1$. Put

$$\lambda(M) = \max_{i \neq 1} |\lambda_i(M)|.$$

Lemma 2.1 *If A and B are disconnected vertex sets of G and $M \in \mathcal{M}(G)$, then*

$$\frac{|A| \cdot |B|}{(v - |A|)(v - |B|)} \leq \lambda^2(M).$$

Proof. See [7] Theorem 2.1, or [12] Lemma 6.1. □

Theorem 2.2 *For any $M \in \mathcal{M}(G)$*

$$\Phi(G) \leq v \frac{\lambda(M)}{1 + \lambda(M)}.$$

Proof. Put $\Phi = \Phi(G)$ and take A and B such that $\Phi^2 = |A| \cdot |B|$. Then by Lemma 2.1

$$\frac{\Phi^2}{\lambda^2(M)} \leq (v^2 - v(|A| + |B|) + \Phi^2) \leq (v^2 - 2v\sqrt{|A| \cdot |B|} + \Phi^2) = (v - \Phi)^2.$$

Clearly $v \geq \Phi$, so $\Phi \leq (v - \Phi)\lambda(M)$, which yields the required bound. □

In order to investigate when the bound of Theorem 2.2 is best possible, we define

$$\phi(G) = \min_{M \in \mathcal{M}(G)} v \frac{\lambda(M)}{1 + \lambda(M)}$$

and we let $\mathcal{M}'(G)$ denote the set of matrices from $\mathcal{M}(G)$ for which the above minimum is attained. Thus Theorem 2.2 becomes $\Phi(G) \leq \phi(G)$. To determine $\phi(G)$ we need to find a matrix in $\mathcal{M}'(G)$. For that purpose the automorphisms of G can be helpful.

Lemma 2.3 *Let \mathcal{A} be an automorphism group of G . Then $\mathcal{M}'(G)$ contains a matrix which is constant over each orbit of the action of \mathcal{A} on $V \times V$.*

Proof. Let P_g denote the permutation matrix corresponding to $g \in \mathcal{A}$ and take $M' \in \mathcal{M}'(G)$. Then clearly $P_g M' P_g^\top \in \mathcal{M}'(G)$ and, by Rayleigh's principle, $|u^\top P_g M' P_g^\top u| \leq \lambda(M')$ for every unit vector u orthogonal to $\underline{1}$. Define

$$\overline{M} = \frac{1}{|\mathcal{A}|} \sum_{g \in \mathcal{A}} P_g M' P_g^\top,$$

then clearly $\overline{M} \in \mathcal{M}(G)$ and \overline{M} is constant over \mathcal{A} -orbits on $V \times V$. Let u ($u \perp \underline{1}$) be a unit eigenvector for the eigenvalue $\pm \lambda(\overline{M})$. Then $\lambda(\overline{M}) = |u^\top \overline{M} u| \leq \lambda(M')$, so $\lambda(\overline{M}) = \lambda(M')$ and hence $\overline{M} \in \mathcal{M}'(G)$. \square

In particular we may take the diagonal constant if G has a transitive automorphism group. Theorem 2.2 leads to a more explicit bound in terms of the Laplacian eigenvalues of G . (If A is the standard adjacency matrix of G and D is the diagonal matrix containing the vertex degrees, then $F = D - A$ is the Laplacian matrix of G . It easily follows that F is positive semi-definite and singular; see for example Brualdi and Ryser [6].)

Theorem 2.4 *Suppose F is the Laplacian matrix of G and let $0 = \mu_1 \leq \mu_2 \leq \dots \leq \mu_v$ be the eigenvalues of F , then*

$$\phi(G) \leq \frac{v}{2} \left(1 - \frac{\mu_2}{\mu_v} \right)$$

with equality if G has an automorphism group that acts transitively on the edges.

Proof. Define

$$M = \frac{-2}{\mu_2 + \mu_v} F + I.$$

Then $M \in \mathcal{M}(G)$ and $\lambda(M) = (\mu_v - \mu_2)/(\mu_v + \mu_2)$, which yields the inequality.

Suppose G has an automorphism group which acts transitively on the edges. Then, by Lemma 2.3 there exists a matrix $M' \in \mathcal{M}'(G)$ such that $M' = xF + D$ for some constant x and diagonal matrix D . Now $M' \underline{1} = \underline{1}$ gives $D = I$ and so

$$\lambda(M') = \max\{|x\mu_2 + 1|, |x\mu_v + 1|\}.$$

It follows that $\lambda(M')$ is minimal if $x\mu_2 + 1 = -x\mu_v - 1$, that is, if $x = -2/(\mu_2 + \mu_v)$. Thus $M \in \mathcal{M}'(G)$. \square

Example. Suppose G is the triangular graph $T(2m)$ (that is, the line graph of K_{2m}). Then $v = m(2m - 1)$, $\mu_2 = 2m$ and $\mu_v = 4m - 2$. Theorem 2.4 gives $\phi(G) = \binom{m}{2}$. We easily have that $\Phi(G) \geq \binom{m}{2}$, so $\Phi(G) = \binom{m}{2}$.

Next we consider association schemes. For theory and notation see [4], [5] or [10]. Let S be an n -class association scheme defined on the set V with relations R_0, \dots, R_n . For $\delta \in \{1, \dots, n\}$ we denote by G_δ the graph (V, R_δ) and by $G_{\bar{\delta}}$ the complement of G_δ .

Theorem 2.5 *If Q is the matrix of dual eigenvalues of S , then*

$$\phi(G_{\bar{\delta}}) \leq \frac{v}{\sum_{j=0}^n |Q_{\delta,j}|}.$$

Equality holds if the automorphism group of S acts transitively on each relation.

Proof. Let A_0, \dots, A_n (with $A_0 = I$) be the adjacency matrices of S and let E_0, \dots, E_n (with $vE_0 = J$, the all-one matrix) be the minimal idempotents. Then the matrix Q of dual eigenvalues is given by

$$vE_j = \sum_{i=0}^n Q_{i,j} A_i, \text{ for } j = 0, \dots, n.$$

So $v(E_j)_{k,l} = Q_{\delta,j}$ whenever $\{k, l\} \in R_\delta$. Define $\mathcal{P}_\delta = \{j | 1 \leq j \leq n, Q_{\delta,j} > 0\}$, $m = \sum_{j \in \mathcal{P}_\delta} Q_{\delta,j} + \frac{1}{2}$ and

$$M = \frac{1}{m} \left(\frac{1}{2} I - \sum_{j \in \mathcal{P}_\delta} (E_j - Q_{\delta,j} E_0) \right).$$

Then, since $E_j \mathbf{1} = \mathbf{0}$ for $j \neq 0$ one readily verifies that $M \in \mathcal{M}(G_{\bar{\delta}})$. Moreover $\sum_{j \in \mathcal{P}_\delta} E_j$ has only 0 and 1 as eigenvalues. This implies that $\lambda_i(M) = \pm \frac{1}{2m}$ for $i \neq 1$, so $\lambda(M) = \frac{1}{2m}$, and thus we find $\phi(G_{\bar{\delta}}) \leq \frac{v}{2m+1}$. By use of $\sum_{j=0}^n Q_{\delta,j} = 0$ and $Q_{\delta,0} = 1$, we obtain

$$m + \frac{1}{2} = 1 + \sum_{j \in \mathcal{P}_\delta} Q_{\delta,j} = \frac{1}{2} \sum_{j=0}^n |Q_{\delta,j}|,$$

and the required inequality follows.

Next assume that S admits an automorphism group which is transitive on each relation. Then by Lemma 2.3 there exists a matrix $M' \in \mathcal{M}'(G_{\bar{\delta}})$ which is a linear combination of A_0, \dots, A_n , that is, M' is in the Bose-Messner algebra of S . Let $\lambda_{j'}(M')$ denote the eigenvalue of M' whose eigenspace is given by E_j . We claim that we may assume that $\lambda_{j'}(M') = \lambda(M')$ if $Q_{\delta,j} \leq 0$. Indeed, suppose this is not the case, then define $d = \lambda(M') - \lambda_{j'}(M')$, $m = 1 - dQ_{\delta,j}$ and $M'' = \frac{1}{m}(M' + d(E_j - Q_{\delta,j}E_0))$. It follows that $M'' \in \mathcal{M}(G_{\bar{\delta}})$, $m \geq 1$ and $\lambda_{j'}(M'') = \lambda(M'') = \frac{1}{m}\lambda(M') \leq \lambda(M')$. So we can redefine $M' = M''$, which proves the claim. Similarly, we may assume that $\lambda_{j'}(M') = -\lambda(M')$ if $Q_{\delta,j} \geq 0$ and $j \neq 0$. It now follows that

$$E = \frac{1}{2\lambda(M')}(\lambda(M')I - M' + (\lambda(M') - 1)E_0)$$

has eigenvalue 0 and 1 only, and hence E is an idempotent of S . Therefore E is the sum of those E_j that correspond to the eigenvalue 1 of E , that is $E = \sum_{j \in \mathcal{P}_\delta} E_j$. In addition, since $(M')_{k,l} = 0$ for $\{k, l\} \in R_\delta$, we have $\sum_{j \in \mathcal{P}_\delta} Q_{\delta,j} = \frac{1}{2\lambda(M')} - \frac{1}{2}$. This implies that $M' = M$, so $M \in \mathcal{M}'(G_{\bar{\delta}})$. \square

A graph $G_{\bar{\delta}}$ in a 2-class association scheme is the same as a strongly regular graph. An example of such a graph is the triangular graph $T(m)$, described in the example above. It is not difficult to see that for strongly regular graphs the bounds of Theorem 2.4 and Theorem 2.5 coincide.

3 Equidistant code pairs

Suppose $V = \{1, \dots, q\}^n$, the set of words of length n over an alphabet of size q , and define two words to be in relation R_δ if their Hamming distance (the number of coordinate places in which they differ) equals δ . This defines the well known Hamming association scheme $H(n, q)$. For a graph $G_{\bar{\delta}}$ in $H(n, q)$ two disconnected sets in $G_{\bar{\delta}}$ are called equidistant code pairs (at distance δ) and we write Φ_δ and ϕ_δ in stead of $\Phi(G_{\bar{\delta}})$ and $\phi(G_{\bar{\delta}})$ respectively.

Lemma 3.1

$$\Phi_\delta^2 \geq \max_{0 \leq \delta' \leq \delta} \binom{n - \delta'}{\delta - \delta'} (q - 1)^{\delta - \delta'} \left(\left\lfloor \frac{q}{2} \right\rfloor \left\lceil \frac{q}{2} \right\rceil \right)^{\delta'}.$$

Proof. Take for A the set of words (x_1, \dots, x_n) with $1 \leq x_i \leq \frac{q}{2}$ if $i \leq \delta'$ and $x_i = 1$ if $i > \delta'$, and let B consist of the words (x_1, \dots, x_n) with $\frac{q}{2} < x_i \leq q$ if $i \leq \delta'$ and $x_i \neq 1$ for precisely $\delta - \delta'$ values of $i > \delta'$. Then A and B form an equidistant code pair at distance δ with sizes $\lfloor \frac{q}{2} \rfloor^{\delta'}$ and $\lceil \frac{q}{2} \rceil^{\delta'} \binom{n-\delta'}{\delta-\delta'} (q-1)^{\delta-\delta'}$, respectively. \square

The above construction was given by Ahlswede [1]. He proves that equality holds for $q = 4$ and $q = 5$ and conjectures equality for all $q \geq 4$. For $q = 2$ and $q = 3$ there exist better constructions of equidistant code pairs (see below).

The Hamming scheme is self-dual, which means that the dual eigenvalues coincide with the eigenvalues. They are given by (see [8]):

$$Q_{\delta,j} = \sum_{k=0}^n (-1)^k (q-1)^{j-k} \binom{\delta}{k} \binom{n-\delta}{j-k} \quad \text{for } \delta, j \in \{0, \dots, n\}.$$

The automorphism group of $H(n, q)$ is transitive on each relation, so Theorem 2.5 gives the exact value of ϕ_δ for all n, q and δ .

Example If $n = 6$ and $q = 6$, then for $j = 0, \dots, 6$ the respective values of $Q_{4,j}$ are 1, 6, -9, -44, 111, -90 and 25. Theorem 2.5 gives $\phi_4 = 46656/286 \approx 163.13$. With Lemma 3.1 (take $\delta' = 2$) we find $45\sqrt{6} \leq \Phi_4 \leq 23328/143$.

This example shows that our bound will not prove Ahlswede's conjecture. But it can give interesting results in some cases.

Theorem 3.2 *If $q > 2$ then*

$$\phi_\delta \leq \frac{q^n}{(q-2)^{n-\delta} 2^\delta}.$$

Equality holds if and only if $\delta = n$.

Proof. The inequality follows from Theorem 2.5 and

$$\sum_{j=0}^n |Q_{\delta,j}| \geq \left| \sum_{j=0}^n (-1)^j Q_{\delta,j} \right| = \left| \sum_{k=0}^n \binom{\delta}{k} \sum_{j=0}^n (-1)^{j-k} (q-1)^{j-k} \binom{n-\delta}{j-k} \right| = 2^\delta (q-2)^{n-\delta}.$$

If j runs from 0 to n , $Q_{n,j}$ alternates in sign, so we have equality if $\delta = n$. The dual eigenvalues of any association scheme satisfy $\sum_{j=0}^n \frac{1}{\mu_j} Q_{\delta,j} Q_{n,j} = 0$ if $\delta \neq n$ ($\mu_j = \text{rk } E_j$). Therefore $Q_{\delta,j}$ cannot alternate in sign if $\delta \neq n$, so then we have strict inequality. \square

Corollary 3.3 *If $q > 2$ then*

$$\Phi_\delta \leq \frac{q^n}{(q-2)^{n-\delta} 2^\delta}.$$

Equality holds if and only if $\delta = n$ and q is even.

Proof. If $\delta = n$ and q is even, Lemma 3.1 (with $\delta' = \delta$) gives $\Phi_n \geq (\frac{q}{2})^n$, which equals ϕ_n . If q is odd, $(\frac{q}{2})^{2n}$ is not an integer, so $\Phi_n \neq \phi_n$. \square

We see that the lower bound of Lemma 3.1 and the upper bound of Corollary 3.3 tend to the same value $(\frac{q}{2})^\delta$ if $q \rightarrow \infty$. More precisely:

Corollary 3.4

$$\Phi_\delta = \left(\frac{q}{2}\right)^\delta + O(q^{\delta-1}) \quad (q \rightarrow \infty).$$

For $q \geq 4$ Ahlswede and Mörs [3] showed that $\Phi_\delta < \Phi_n$ if $\delta < n$. This result now follows directly from Corollary 3.3 when q is even and, by Lemma 3.1, also when q is odd and n is not too big.

For $q = 3$ not much is known about Φ_δ . Ahlswede [1] has a construction for equidistant code pairs and conjectures that it is best possible. If this is true then Φ_δ attains its maximal value

$$\left(\frac{3}{2}\right)^{\lfloor \frac{n}{3} \rfloor} 2^{\frac{n}{2}}$$

if $\delta = \lceil \frac{2n}{3} \rceil$. Theorem 3.2 gives $\phi_n = (\frac{3}{2})^n$, thus we have that $\Phi_n < \Phi_\delta$ if $\delta = \lceil \frac{2n}{3} \rceil$ ($n > 2$). By use of Theorem 2.5, stronger results are possible, but it turns out that the bound ϕ_δ is not good enough to prove that Φ_δ is maximal if $\delta = \lceil \frac{2n}{3} \rceil$.

For $q = 2$ the value Φ_δ is known for all δ ; see [1]. It attains the maximal value $2^{\lfloor \frac{n}{2} \rfloor}$ if $\delta = \lfloor \frac{n}{2} \rfloor$ and if $\delta = \lceil \frac{n}{2} \rceil$. This result was first proved by Ahlswede, El Gamal and Pang [2] and has several different proofs now (see [1]). We shall see that our bound provides yet another proof. The construction is as follows. Take for A the set of words (x_1, \dots, x_n) with $x_{2i-1} = x_{2i}$ for $1 \leq i \leq \frac{n}{2}$ and $x_n = 1$ if n is odd. Take for B the set of words with $x_{2i-1} \neq x_{2i}$ for $1 \leq i \leq \frac{n}{2}$ and x_n fixed if n is odd. Then A and B are equidistant code pairs at distance $\lfloor \frac{n}{2} \rfloor$ or $\lceil \frac{n}{2} \rceil$ and $|A| = |B| = 2^{\lfloor \frac{n}{2} \rfloor}$.

Theorem 3.5 *If $q = 2$ then*

$$\phi_\delta \leq 2^{\lfloor \frac{n}{2} \rfloor} \text{ for } \delta \in \{1, \dots, n\}.$$

Equality holds if $\delta \in \{\lfloor \frac{n}{2} \rfloor, \lceil \frac{n}{2} \rceil\}$.

Proof. With $i = \sqrt{-1}$ we have

$$\sum_{j=0}^n i^j Q_{\delta,j} = \sum_{k=0}^n i^{-k} \binom{\delta}{k} \sum_{j=0}^n i^{j-k} \binom{n-\delta}{j-k} = (1-i)^\delta (1+i)^{n-\delta} = 2^{\frac{n}{2}} \omega,$$

wherein $\omega = e^{\frac{\pi i}{4}(n-2\delta)}$. Hence

$$\sum_{j=0}^n |Q_{\delta,j}| = \sum_{j \text{ even}} |Q_{\delta,j}| + \sum_{j \text{ odd}} |Q_{\delta,j}| \geq 2^{\frac{n}{2}} (|\operatorname{Re} \omega| + |\operatorname{Im} \omega|) = 2^{\lceil \frac{n}{2} \rceil},$$

and the inequality follows by use of Theorem 2.5. The construction above shows that $\Phi_\delta = \phi_\delta = 2^{\lfloor \frac{n}{2} \rfloor}$ if $|\frac{n}{2} - \delta| \leq \frac{1}{2}$. □

With a similar argument and a bit more work as in the proof of Theorem 3.2 it can be seen that if $|\frac{n}{2} - \delta| \geq 1$ the bound ϕ_δ is strictly less than $2^{\lfloor \frac{n}{2} \rfloor}$.

4 Concluding remarks

There is some similarity between our function $\phi(G)$ and Lovász's function $\theta(G)$ (see [13]). The latter function gives an upper bound for a coclique (independent set of vertices) in G , which is, in a sense, a vertex set disconnected to itself. Lovász's $\theta(G)$ is known to be computable in polynomial time (see [11]), but we do not know the complexity of the computation of $\phi(G)$. The computation of $\Phi(G)$ seems to be NP-hard; see Garey and Johnson [9], p.196.

If we apply Theorem 2.5 to the Johnson association scheme $J(m, n)$, we obtain bounds for equidistant code pairs in (binary) constant weight codes. It seems certainly worthwhile to work this out and we intend to do so. The problem is that the formulas for the dual eigenvalues are rather complicated. For the special case that $\delta = n$ we believed that $\sum_j |Q_{\delta,j}| = \binom{\frac{m}{2}}{n}$, but had no proof, until Volker Strehl (private communication) provided us with a computer-generated proof by use of Zeilberger's algorithm.

Acknowledgement. The research for this paper was done when the author was a guest of the Sonderforschungsbereich "Diskrete Strukturen in der Mathematik" at the University of Bielefeld. Special thanks go to Professor Rudi Ahlswede for the invitation, and several fruitful discussions concerning the subject.

References

- [1] R. Ahlswede, *On code pairs with specified Hamming distances*, Colloquia Mathematica Societas János Bolya **52**, Combinatorics, Eger (Hungary), 1987, pp. 9-47.
- [2] R. Ahlswede, A. El Gamal and K.F. Pang, *A two-family extremal problem in Hamming space*, Discrete Math. **49** (1984), 1-5.
- [3] R. Ahlswede and M. Mörs, *Inequalities for code pairs*, Europ. J. Combinatorics **9** (1988), 175-181.
- [4] A.E. Brouwer, A.M. Cohen and A. Neumaier, *Distance-regular graphs*, Springer-Verlag, Berlin, 1989.
- [5] A.E. Brouwer and W.H. Haemers, *Association schemes*, chapter 15 in: Handbook of Combinatorics (R. Graham, M. Grötschel and L. Lovász eds.), Elsevier Science, Amsterdam, 1995, pp. 747-771.
- [6] R.A. Brualdi and H.J. Ryser, *Combinatorial matrix theory*, Cambridge University Press, Cambridge, 1991.
- [7] E.R. van Dam and W.H. Haemers, *Eigenvalues and the diameter of graphs*, Linear Multilinear Algebra **39** (1995), 33-44.
- [8] Ph. Delsarte, *An algebraic approach to the association schemes of coding theory*, Philips Research Rep. Suppl. **10**, 1973.
- [9] M.R. Garey and D.S. Johnson, *Computers and intractability: A guide to the theory of NP-completeness*, Freeman, San Francisco, 1979.
- [10] C.D. Godsil, *Algebraic combinatorics*, Chapman and Hall, New York - London, 1993.

- [11] M. Grötschel, L. Lovász and A. Schrijver, *The ellipsoid method and its consequences in combinatorial optimization*, *Combinatorica* **1** (1981), 169-197.
- [12] W.H. Haemers, *Interlacing eigenvalues and graphs*, *Linear Algebra Appl.* **226-228** (1995), 593-616.
- [13] L. Lovász, *On the Shannon capacity of a graph*, *IEEE Trans. Inform. Theory* **25** (1979), 1-7.

Willem H. Haemers,
Tilburg University,
Dept. of Econometrics,
P.O.Box 90153,
5000 LE Tilburg,
The Netherlands.
e-mail: haemers@kub.nl