

Tilburg University

Cross domain self-monitoring in anosognosia for memory loss in Alzheimer's disease

Chapman, Silvia; Colvin, Leigh E.; Vuorre, Matti; Cocchini, Gianna; Metcalfe, Janet; Huey, Edward D.; Cosentino, Stephanie

Published in:
Cortex

DOI:
[10.1016/j.cortex.2018.01.019](https://doi.org/10.1016/j.cortex.2018.01.019)

Publication date:
2018

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

Chapman, S., Colvin, L. E., Vuorre, M., Cocchini, G., Metcalfe, J., Huey, E. D., & Cosentino, S. (2018). Cross domain self-monitoring in anosognosia for memory loss in Alzheimer's disease. *Cortex*, *101*, 221-233. <https://doi.org/10.1016/j.cortex.2018.01.019>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Published in final edited form as:

Cortex. 2018 April ; 101: 221–233. doi:10.1016/j.cortex.2018.01.019.

Cross Domain Self-Monitoring in Anosognosia for Memory Loss in Alzheimer's disease

Silvia Chapman^{1,5}, Leigh E. Colvin^{1,4}, Matti Vuorre⁴, Gianna Cocchini⁵, Janet Metcalfe⁶, Edward D. Huey^{1,2,3}, and Stephanie Cosentino^{1,2,3}

¹Cognitive Neuroscience Division, Columbia University Medical Center, New York, NY, United States

²Taub Institute for Research on Alzheimer's disease and the Aging Brain, Gertrude H. Sergievsky Center, Columbia University Medical Center, New York, NY, United States

³Department of Neurology, Columbia University Medical Center, New York, NY, United States

⁴Teachers College, Columbia University, New York, NY, United States

⁵Goldsmiths College, University of London, London, United Kingdom

⁶Department of Psychology, Columbia University, New York, NY, United States

Abstract

Anosognosia for memory loss is a common feature of Alzheimer's disease (AD). Recent theories have proposed that anosognosia, a disruption in awareness at a *global* level, may reflect specific deficits in self-monitoring, or *local* awareness. Though anosognosia for memory loss has been shown to relate to memory self-monitoring, it is not clear if it relates to self-monitoring deficits in other domains (i.e., motor). The current study examined this question by analyzing the relationship between anosognosia for memory loss, memory monitoring, and motor monitoring in 35 individuals with mild to moderate AD. Anosognosia was assessed via clinical interview before participants completed a metamemory task to measure memory monitoring, and a computerized agency task to measure motor monitoring. Cognitive and psychological measures included memory, executive functions, and mood. Memory monitoring was associated with motor monitoring; however, anosognosia was associated only with memory monitoring, and not motor monitoring. Cognition and mood related differently to each measure of self-awareness. Results are interpreted within a hierarchical model of awareness in which local self-monitoring processes are associated across domain, but appear to only contribute to global levels awareness in a domain-specific fashion.

Data statement

Due to restrictions of conditions of data sharing with individuals outside of the research team, the data set and syntax of analysis was not uploaded. These conditions were set prior commencement of the study and were approved by the ethical bodies where recruitment took place (i.e., The Institute Review Board (IRB) at Columbia University Medical Center).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Metacognition; anosognosia; agency; cognition; Alzheimer's disease

1. INTRODUCTION

Individuals with Alzheimer's disease (AD) are often unaware of their deficits (see Agnew & Morris, 1998; Cosentino & Stern, 2005; Rosen, 2011). This disordered higher level self-awareness, or anosognosia, has been linked to a variety of negative personal and societal consequences, with "unaware" individuals engaging in and benefiting less from clinical management, demonstrating reduced capacity to make treatment decisions (Clare, Wilson, Carter, Roth, & Hodges, 2004; Cosentino, Metcalfe, Cary, De Leon, & Karlawish, 2011; Koltai, Welsh-Bohmer, & Schmechel, 2001), and evidencing more risky behaviors than those who are aware of their deficits (Cotrell & Wild, 1999; Wild & Cotrell, 2003). Moreover, those responsible for the care of unaware patients report higher degrees of stress and burden (Prigatano, 2005; Rymer et al., 2002; Seltzer, Vasterling, Yoder, & Thompson, 1997), even in the context of Mild Cognitive Impairment (MCI) (Kelleher, Tolea, & Galvin, 2016). As efforts to diagnose AD move toward a preclinical stage during which individuals have the capacity to be highly functional if aware of their cognitive deficits, understanding the specific metacognitive impairments leading to anosognosia in AD may be critical for enabling individuals to maintain their autonomy.

There is a growing yet incomplete understanding of the ways in which self-awareness breaks down in AD, as well as other conditions such as stroke. Existing models of anosognosia, or *global* awareness, have outlined the ways in which dysfunctional memory and executive systems can give rise to disordered awareness in AD (see Agnew & Morris, 1998; Ansell & Bucks, 2006; Morris & Mograbi, 2013). However, given that disruptions to memory (Derouesne et al., 1999; Reed, Jagust, & Coulter, 1993; Starkstein et al., 1995) and executive function (Lopez, Becker, Somsak, Dew, & DeKosky, 1994; Michon, Deweer, Pillon, Agid, & Dubois, 1994; Reed et al., 1993; Starkstein, Sabe, Chemerinski, Jason, & Leiguarda, 1996) do not fully explain anosognosia in AD, it is clear that other mechanisms are at play in the deterioration of higher levels of self-awareness. Currently, there is a drive in both the cognitive and motor literatures, towards a dynamic and multifaceted notion of self-awareness wherein factors specific to metacognition, not simply cognition, give rise to this fascinating disorder (Clare, Marcova, Roth, & Morris, 2011; Davies, Davies, & Coltheart, 2005; Fotopoulou, 2014; Levine, 1990; Rosen, 2011).

In this vein, research has begun to examine processes that may be uniquely self-evaluative, operating outside of primary cognitive abilities. One such process is self-monitoring or *local* awareness—the process by which one evaluates aspects of one's own individual thoughts, intentions and actions compared to those of others or those arising from the external world (Agnew & Morris, 1998; Fotopoulou, 2014; Jenkinson & Fotopoulou, 2010; McGlynn & Schacter, 1989; Rosen, 2011; Saj, Vocat, & Vuilleumier, 2014; Venneri & Shanks, 2004). Leading theorists in anosognosia for memory loss, have conceptualized awareness within a hierarchical structure in their models, with monitoring processes, or local awareness,

considered to be underlying anosognosia, or higher order global awareness (i.e., the Conscious Awareness Model (CAM) (Agnew & Morris, 1998; Mograbi & Morris, 2013) and the Biopsychosocial hierarchical model (Clare et al., 2011)). Previous work in AD has linked traditional clinical ratings of anosognosia, understood as a higher level of awareness (i.e., *global* awareness), with performance on memory monitoring or metamemory measures, understood as lower levels of awareness (i.e., *local* awareness) (Cosentino, Metcalfe, Butterfield & Stern, 2007; Clare et al., 2011). However, relatively little work has examined the extent to which anosognosia in AD, stroke, or other conditions is characterized by broad deficits in self-monitoring or domain specific deficits in self-monitoring.

In the current study, we examined the association between anosognosia for memory loss in AD, memory monitoring, and motor monitoring (i.e., agency judgments, or the extent to which individuals perceive themselves to be the agent of a determined outcome or action) (Gallagher, 2000). There is an inherent necessity of accessing self-specific information when making a judgment of agency related to an action or thought, and agency tasks have been used to understand unawareness of hemiplegia or other motor deficits following stroke (Fotopoulou et al., 2008), providing an ideal framework to examine self-referential monitoring in a non-memory domain. Indeed, much of the work dedicated to modeling anosognosia and examining the role of monitoring difficulties has occurred in the context of impaired motor functioning in individuals with stroke who are unaware of hemiplegia (Jenkinson, Edelstyn, Drakeford, & Ellis, 2009; Saj et al., 2014; Venneri & Shanks, 2004; Vocat, Saj, & Vuilleumier, 2013). Conceptually, it has been proposed that discrepancies in monitoring between one's intentions (i.e., motor plan) and one's actual motor performance may result in unawareness of hemiplegia (Berti, Spinazzola, Pia, & Rabuffetti, 2007; Cocchini, Beschin, Fotopoulou, & Della Sala, 2010; Fotopoulou et al., 2008; Moro, Pernigo, Zapparoli, Cordioli, & Aglioti, 2011). The *Comparator Model of motor control* posits that for each produced movement, an individual *implicitly* monitors their intentions and predicted outcome in relation to sensory and perceptual feedback about the actual outcome (Blakemore, Wolpert, & Frith, 2002). The comparison between these two processes allows the detection of a mismatch that would occur in the context of a movement error, and therefore allows correction of the error. The comparison also provides a neural basis for the perception of a distinction between internally driven movements (where the match between the two processes is high) and those movements caused by an external source (Feinberg, 1978; Frith, 2005).

Another explanatory model of judgments of agency or judgments of motor monitoring is the *Theory of mental causation* (Wegner, 2002; Wegner & Wheatley, 1999), by which individuals *consciously* assess the relationship between intentions and actions, and infer causal judgments of agency. This theory moves away from the underlying process of motor monitoring, arguing that such processes are unconscious. According to this theory, conscious processes such as the intention associated with the action and contextual cues of the outcome itself are utilized to derive an inferential judgment of agency or judgments of motor monitoring (Haggard & Tsakiris, 2009; Metcalfe, Eich, & Castel, 2010; Moore, 2016; Synofzik, Vosgerau, & Newen, 2008; Wegner, 2002; Wegner & Wheatley, 1999).

Several studies have supported the association of motor monitoring and anosognosia for hemiplegia (e.g., Jenkinson & Fotopoulou, 2010; Vocat et al., 2013). Interestingly, monitoring deficits in patients unaware of their motor deficits seem to relate to monitoring deficits in other cognitive domains (Feinberg, Roane, Kwan, Schindler, & Haber, 1994; Jenkinson et al., 2009; Venneri & Shanks, 2004). These cross-domain associations suggest that at least in the case of anosognosia for motor deficits, its underlying mechanisms may not be domain specific and that a combination of different processes may be key to the emergence of impaired awareness (e.g., deficient error prediction, encoding, monitoring and premorbid factors) (Cocchini, Beschin, & Sala, 2002; Davies et al., 2005; Fotopoulou, 2014; Levine, 1990; Marcel, Tegnér, & Nimmo-Smith, 2004; McGlynn & Schacter, 1989; Vuilleumier, 2004). The association of self-monitoring abilities across different task domains has also been demonstrated in non-demented cohorts in which the integrity of memory monitoring and motor monitoring (i.e., *agency*) judgments have been linked (Cosentino, Metcalfe, Holmes, Steffener, & Stern, 2011).

To our knowledge, there are no previous studies examining judgments of agency in AD. Given both the cross-domain monitoring deficits seen in individuals with anosognosia for hemiplegia, as well as the link between memory monitoring and agency monitoring in older adults, one might hypothesize that anosognosia in AD may be associated with compromised agency in AD. However, there is also reason to believe that these processes may be dissociated. While they are both self-referential, the substrates that contribute to each judgment are seemingly very different. For example, memory monitoring has been hypothesized to rely on memory abilities, executive functioning, and underlying implicit internal monitoring of mnemonic processes such as familiarity and partial access to information (Cosentino, Metcalfe, Holmes, et al., 2011; Koriat, 1993; Koriat & Levy-Sadot, 2001; Metcalfe, Schwartz, & Joaquim, 1993; Reder & Ritter, 1992; Schnyer et al., 2004; Schwartz & Metcalfe, 1992). In contrast, judgments of agency have been hypothesized to rely on the monitoring of sensory and perceptual stimuli of the action and the integration of different contextual cues such as, perceived success, temporal delay between intention and outcome and reward (Blakemore et al., 2002; Frith, Blakemore, & Wolpert, 2000; Kirkpatrick, Metcalfe, Greene, & Hart, 2008; Metcalfe, Van Snellenberg, DeRosse, Balsam, & Malhotra, 2014; Michotte, 1963; Moore, 2016; Schlottman & Shanks, 1992).

The purpose of this study is to clarify the association between different domains and levels of awareness in AD by examining the relationship between anosognosia for memory loss, memory monitoring and agency. For this purpose, we ran regression models examining the associations among these three self-evaluative measures including covariates such as memory, executive functions and mood (Ansell & Bucks, 2006; Bertrand et al., 2016; Cines et al., 2015; Clare et al., 2012; Conde-Sala et al., 2014; Cosentino, Metcalfe, Holmes, et al., 2011; Mograbi & Morris, 2013; Perrotin, Isingrini, Souchay, Clarys, & Tacconat, 2006; Reed et al., 1993). In doing so, this study will refine and build upon current models of self-awareness with the goal of improving their ultimate utility for guiding the management of anosognosia in AD.

2. METHODS

2.1. Participants

As part of a larger study, 51 participants with mild to moderate AD were recruited through the Department of Neurology at the Columbia University Medical Center. Participants had a diagnosis of Alzheimer's following the criteria of the Neurologic Disorders and Stroke - Alzheimer's disease and Related Disorders Association (NINDS-ADRDA). Participants were excluded from the study if there was evidence of moderate to severe psychiatric illness, history of acquired brain injury (traumatic and vascular), or any other neurological conditions that may have had an impact on cognition. Participants were also excluded if they scored under 20 in the Mini-Mental State Examination (Folstein, Folstein, & McHugh, 1975) to ensure comprehension of the tasks. Participants with atypical presentations of AD that were not characterized primarily by memory loss (i.e., language or frontal variant AD) were excluded. All participants provided informed consent and all procedures were approved by the Institutional Review Board at Columbia University Medical Center.

Of the original 51 participants, only 38 completed the agency task. Three participants were then excluded from final analysis due to missing or invalid data on the agency task resulting in a final sample size of 35 participants (69% females). The overall mean age was 77.72 ($SD = 9.40$; range = 57–99) and over 91% of the participants were white Caucasians; the remaining 9% were African American. All participants were assessed across three visits.

2.2. Measures

2.2.1. Anosognosia—Anosognosia was evaluated via a brief interview at the beginning of each of the three study visits, generating a Clinical Rating of Awareness (CRA) regarding memory functioning. We used a modified version of Reed et al.'s (1993) clinical awareness scoring categories. Participants were asked an open-ended question about their memory (e.g., "how is your memory?"). Based on participants' responses, the examiner rated their awareness in line with the following scoring system: 1.00 = Full Awareness (e.g., Patient spontaneously complains of significant memory loss and may discuss memory loss as consequential of the disease); 2.00 = "Moderate Awareness" (e.g., Patient spontaneously admits significant memory loss but attributes it to normal aging); 3.00 = "Shallow Awareness" (e.g., Patient is inconsistent or uncertain about memory loss); 4.00 = "No Awareness" (e.g., Patient denies memory loss). Repeated measures examined if there were significant differences of awareness across the three visits before averaging these into one score. For purposes of this study, the scoring ratings were then collapsed into two categories (1–2 = "Aware"; >2–4 = "Unaware") in line with previous publications (Cosentino et al., 2016).

2.2.2. Cognitive Measures—Participants underwent neuropsychological examination, which included measures of global cognition, memory, executive functions and attention. Memory measures consisted of the Philadelphia Verbal Learning Task (PVL - Price et al., 2009) for verbal memory and the Biber Learning Test (Glosser, Goodglass, & Biber, 1989) as a nonverbal memory measure. Executive function measures included a design fluency task (Glosser & Goodglass, 1990), a verbal fluency task (i.e., FAS) (Stuss & Benson, 1986)

and the Digit and Spatial backward spans from the Wechsler Memory Scale - Revised (WMS-R) (Wechsler, 1987). Attention was assessed with a visual scanning task. Cognitive index scores were obtained from these measures to represent three main cognitive domains: memory, executive functions and attention. A memory index score was obtained by averaging z scores of the total immediate recall and long delayed recall of both the PVLTL and Biber memory tests. An executive index score was derived from an average of the Digit and Spatial spans backward, FAS and Design fluency z scores. Finally, an attention score was derived from the z scores of the visual scanning task (Cosentino, Metcalfe, Holmes, et al., 2011).

2.2.3. Mood—Mood was assessed with the 30 item Geriatric Depression Scale (GDS). This measure includes a variety of items targeting symptoms of depression such as sadness, hopelessness, dissatisfaction with life and worthlessness. Higher scores represent higher endorsement of depressive items, with a cut off of 10 as indicative of at least mild depression. This measure has been shown to have high validity and reliability in measuring the construct of depression (Yesavage et al., 1982).

2.2.4. Self-Monitoring Measures

Memory Monitoring Task: A modified Feeling of knowing or FOK task was used in this study. As part of this task, participants underwent three different tasks conditions (standard, query, and feedback; described below) counterbalanced across three visits. Task condition and trivia set were compared and analyzed to determine that performance was not affected by condition before collapsing the scores. Each condition of the FOK task was comprised of four trials with five items per trial. Prior to commencing all trials, participants were instructed: (i.e., “During this task, I am going to tell you about five people. I will tell you their name and something about their background. Your task is to try to remember this information as best you can. Please listen carefully”). After hearing the information read aloud, participants were asked to give a global judgment of learning (JOL) (i.e., “Now I am going to test your memory for those names, giving you answer choices. Of the five names, how many do you think you will get right?”). Then, for each of the five items, participants were shown the individual question and asked to estimate the likelihood of knowing the right answer (FOK judgment; i.e., “There are eight possible answers on the next page. Will you know which one is right – Yes, Maybe, or No?”). After each FOK judgment, participants were shown eight answer choices which included the correct answer as well as seven distractors. These seven distractors included the other four names that had been presented in the learning trials, and three unrelated distractors. This procedure was the same for each condition (standard, query and feedback) except that for the query and feedback conditions, in which one more element was included. In the query condition, participants were also asked to make a judgment, after each item, regarding the accuracy of their answer. In the feedback condition, the examiner provided participants with verbal feedback on the accuracy of their response after each item. Each of the four global JOLs provided before each trial ranged from 0 to 5. Item level prediction judgments were given ordinal values of 0 = No; 0.5 = Maybe and 1 = Yes. Performance (i.e., memory) accuracy had values of 0 = incorrect and 1 = correct to enable the calculation of the measures below.

Resolution (Gamma): Resolution reflects the extent to which participants are able to adjust their predictions for performance on each item in line with actual memory performance on that item. Resolution was measured with the Goodman-Kruskal gamma statistic, a rank order correlation that is based on the total amount of *concordances* across the test (C ; predictions for performance on an item are heightened when performance on that item is high, and vice versa) and the total number of *discordances* (D ; predictions for performance on an item are lowered when performance on that item is high, and vice versa). Gamma is calculated as $(C-D)/(C+D)$. Following this formula, more concordances will result in a value of gamma closer to 1 (perfect resolution), whilst the opposite will result in a value of gamma closer to -1 . This calculation does not take in account the number of “ties” where predictions and accuracy are equal in two pairs. Therefore, if someone “ties” across all pairs, gamma cannot be calculated. To avoid losing data in these cases, a formula was developed so that a value of 0 was assigned to gamma, representing the randomness or no association between predictions and actual accuracy (see Cosentino et al., 2015).

Calibration: Calibration scores reflect the extent to which individuals are generally over or under confident in their predictions. For this study, two measures of calibration were obtained, global calibration judgments and item level calibration.

Global calibration judgments reflect the overall level of predictive confidence participants had in their upcoming performance for each 5-item learning trial. These scores were calculated for each of the four trials by subtracting predictions of accuracy (ranging from 0–5) from total accuracy (ranging from 0–5) and dividing by 5 (the total number of items in the trial). The Global calibration judgments represent the average score across all four trials. Values close to 0 represent accurate judgments. Positive values indicate overconfidence, and negative values indicate under confidence.

Item level calibration indicates the extent to which participants are under or over confident in their performance at the item by item level (i.e., “Will you know whether this item is right? Yes? Maybe? No?”). Predictions were given a score of 0 if the participant stated they would not recognize the correct choice, 0.5 if they were not sure and stated “maybe”, and a score of 1 if they were sure they would recognize the right answer. Memory recognition accuracy was scored 0 if they chose the wrong answer, and 1 if they chose the correct answer. Item level calibration was calculated by summing all predictions for performance within all trials, subtracting the sum of accuracy scores, and dividing by the total number of items (e.g., $(\sum \text{prediction} - \sum \text{accuracy}) / \text{total items}$). The resulting measure reflects the extent to which a patient is overconfident (positive values), or under confident (negative values) in their item-level predictions compared to their actual performance. Item level calibration was calculated across each of the four trials and averaged to create a single score. A final average score was computed across conditions (i.e., standard, query and feedback).

Agency Task: A computer task was used to measure patients’ ability to monitor when they were or were not in control of motor outcomes whilst playing a simple computerized game. A modified version of Metcalfe and Greene (2007) task was used (see Cosentino, Metcalfe, Holmes, et al., 2011). In this task, participants were required to move the cursor of a computer horizontally across the bottom of the screen to try to “catch” as many “X”s as

possible whilst avoiding the “O”s, both of which were falling vertically in the screen. In the modified version of Metcalfe and Greene's task, on some of the trials, participants were in complete control of the computer mouse, and so they should have said that they were 'in control.'; on other trials, the computer interfered with the position of the cursor, and so on these trials, to the extent that they correctly recognized their own lack of control over the cursor, they should have said that the computer was 'in control'. Participants were given 1 practice trial, 8 trials in which they were in complete control of the cursor, 8 trials in which the computer controlled the cursor, and 8 split trials in which they were in control half of the time and the computer took over the other half. In computer trials, the cursor on the screen moved directly towards the proximal target in a linear fashion without actively attempting to avoid O's. The person's own mouse movements had no effect on this trajectory. The trials were presented in random order and each had a duration 10 seconds.

To begin each trial, the participant had to move the cursor. If they failed to do so, a message would inform them that the game would not begin if they did not perform a movement. This avoided the strategy of waiting to see if the computer moved the cursor. At the end of each trial participants were required to make a judgment of agency (i.e., “who was in control”) between two dichotomous choices of themselves or the computer as being in control.

Agency judgments, or motor monitoring, was measured as the total accuracy of all judgments on self-and computer-based trials. A combined score of both trial types ranged from 0 to 16. Accuracy for each trial was also derived which ranged from 0 to 8 in each. Split trials were excluded from analysis.

2.2.5. Computer Experience Questionnaire—Three questions regarding computer experience were presented to participants about how often and how comfortable they felt using a mouse: (i) “How often did you use a mouse before the study?”, responses were recorded in a Likert scale from 0 = Never, 1 = A few times and 2 = Many times; (ii) “How comfortable are you using a mouse?”, responses were recorded in a Likert scale from 0 = Not comfortable, 1 = Somewhat comfortable, and 2 = Very comfortable; (iii) “How often did you use a mouse last year?”, responses were recorded in a Likert scale from 0 = Never; 1 = A few times; 2 = Several times a month; 3 = Several times a week; and 4 = Daily. A composite score, used as a measure of overall computer experience, was developed by averaging the results of the three questions.

2.3. Statistical analysis

GLM and non-parametric Friedman tests for repeated measures were used to explore differences between metacognitive and CRA scores administered across the three visits before averaging these into one score. Two-tailed independent t tests, and Mann-Whitney U tests were used to explore differences in cognitive and other self-evaluative measures (memory monitoring and judgments of agency) between participants aware and unaware of their memory deficits as defined by the CRA. Bivariate one and two-tailed Pearson's and Spearman correlations were then used to examine the relationship between the self and computer trials of the agency test, between agency and computer experience, and between

agency and memory monitoring measures. Finally, linear and logistic regression analyses were conducted to examine the correlates of CRA, memory monitoring and agency.

3. RESULTS

3.1. Descriptive Results

3.1.1. Anosognosia—Anosognosia was examined through CRA at each visit. A non-parametric Friedman test for repeated measures revealed no significant difference of awareness ratings across the three sessions $\chi^2(2) = .95, p = .62$. The scores of the three visits were averaged to provide a composite score, and the scores were then collapsed into two categories described in the methods (aware and unaware). 57% of our sample was classified as unaware (shallow or no awareness) and 43% as aware of their memory deficits (full or moderate awareness). The awareness groups did not differ significantly in demographic variables (see Table 1).

With regard to cognitive tasks, unaware participants appeared to perform somewhat worse on memory tasks, though this qualitative difference was not significant ($t(33) = -1.69, p = .10$). No differences were found in executive functions ($t(29) = .11, p = .90$), or attention ($t(32) = 1.61, p = .11$). Depressive symptoms were comparable across groups ($U = 94, z = -1.87, p = .06$).

3.1.2. Memory Monitoring Task—As noted in the methods section, the data presented in this paper are part of a larger study, and participants were exposed to three different FOK conditions (standard, query and feedback). GLM Repeated measures corrected for Green House Geisser showed no difference in memory monitoring as measured by gamma ($F(1.62, 50.24) = 1.72, p = .19$) across the conditions. Similarly, GLM repeated measures for prospective global calibration judgments revealed no differences across conditions for either the global or item level predictions ($F(2, 56) = .64, p = .53$; $F(2, 62) = 1.26, p = .28$). These metacognitive metrics were therefore averaged across visits to create composite scores for comparison with agency and anosognosia for memory loss. Within the memory monitoring scores, resolution (i.e., gamma) was not significantly correlated with item calibration ($r = .28, p = .11$) or global calibration judgments ($r = -.11, p = .55$).

3.1.3. Agency Task—Bivariate Pearson's correlation revealed no association between accuracy of agency judgments in self trials and computer trials ($r = -.10, p = .54$). Therefore, agency was broken down into two scores reflecting each trial type and examined separately in subsequent analyses. Overall, both unaware and aware participants performed significantly better on self trials ($M = 6.02, SD = 1.69$) as compared to computer trials ($M = 2.40, SD = 2.38$) ($t(34) = 7.02, p < .001$; $d = 1.75$).

3.1.4. Computer Experience Questionnaire—Computer mouse experience data were available for 25 participants. Out of these, 47% reported using a mouse before the study many times, whilst 26% had used it a few times, and 26% had never used one. More specifically, 73% of participants reported using the mouse at least once within the last year. Finally, participants were asked how comfortable they felt using a mouse, and 39% reported being very comfortable, 26% somewhat comfortable and 34% not comfortable. The

relationship between computer experience and agency was not significant for self ($r = 0.00$, $p = .99$) or computer trials ($r = .33$, $p = .11$).

3.2. Bivariate Relationships between Awareness Measures

Comparison of the three memory monitoring metrics (gamma, global, and item level calibrations) between unaware and aware participants showed a significant difference only for the gamma score ($t(33) = -3.02$, $p = .005$; $d = 1.06$; see Table 2) such that participants who were unaware of their deficits tended to have lower resolution scores—that is, they showed greater difficulties predicting their memory performance. This difference remained significant after Bonferroni correction for multiple comparisons. There were no differences between anosognosic and aware participants in the accuracy of their agency judgments for self trials ($t(33) = -.51$, $p = .61$) or computer trials ($t(33) = 0.00$, $p = 1.00$). Total judgments of agency showed a qualitative but not significant association with gamma ($r = .28$, $p = .0501$; $d = .57$). Although the correlation between the accuracy of the agency judgments on the computer trials and the resolution gamma correlations was not significant ($r = .11$, $p = .25$), an association was found between the accuracy of agency judgments for self trials and the resolution gamma correlations ($r = .30$, $p = .04$).

3.3. Regression analyses

Because our interest was in exploring the relation between the three measures of self-evaluation (i.e., memory monitoring as measured by gamma, CRA and agency), adjusting for potential covariates, these three variables were included in all models as dependent variables and/or predictors. Covariates were selected on theoretical bases as well based on previously shown associations. The first linear regression was conducted to examine the extent to which gamma could be predicted by scores on agency self trials, CRA, mood, memory and executive function indexes, entered in a single block. Results indicated that the overall model was significant and explained 50 % of the variance ($R^2 = .50$, $F(5, 28) = 5.77$, $p = .001$). It was found that higher memory ($B = .25$, $p = .002$), greater accuracy for agency self trials ($B = .08$, $p = .008$) and higher clinical rated awareness ($B = .23$, $p = .03$) significantly predicted higher gamma. When controlling for demographics, including age, sex and education, the model remained significant, as did the three predictors.

Two additional linear regressions were conducted to examine the predictors of accurate judgments of agency in the self trials and in the computer trials. Predictors included the executive function index, gamma, CRA and computer experience. The overall model, however, was not significant for either the self ($R^2 = .30$, $F(5, 19) = 1.68$, $p = .19$) or the computer trials ($R^2 = .21$, $F(4, 20) = 1.31$, $p = .30$).

Finally, a logistic regression was conducted to explore the extent to which CRA could be predicted by gamma, agency accuracy for self trials, mood, memory and executive function, entered in one block. Results indicated that the overall model was significant ($\chi^2(5) = 13.37$, $p = .02$) and explained 43.6% of the variance (Nagelkerke R^2) in clinically rated awareness. Increasing accuracy in gamma was associated with increased likelihood of being aware of their memory deficits ($B = 3.77$, Wald $\chi^2(1) = 4.42$, $p = .03$) as was endorsing more depression in the Geriatric Depression scale ($B = .21$, Wald $\chi^2(1) = 4.10$, $p = .04$). No

other predictors were significant. When controlling for demographics, only gamma remained a significant predictor of clinically rated awareness. All predictors, for each model, are summarized in Table 3.

4. DISCUSSION

This paper examined the extent to which anosognosia (i.e., a global marker of awareness) in AD is characterized by deficits in specific aspects of online self-monitoring (i.e. lower level of awareness) across domains. Moreover, we explored whether these specific forms of self-monitoring deteriorate in tandem or are dissociable processes. By exploring different measures of self-awareness, this study seeks to understand how different aspects of self-evaluation operate in the context of AD.

Traditionally, anosognosia in AD has been measured by acquiring a patient's global (or offline) self-evaluative judgments regarding his or her general or everyday level of memory functioning (Clare, Marcová, Verhey, & Kenny, 2005; Cosentino & Stern, 2005). These self evaluations can then be compared to an informant's judgment of the person's memory abilities (i.e., Subjective Rating Discrepancy – SRD) (Clare, Wilson, Carter, Roth, & Hodges, 2002), or can be evaluated by a clinician during an interview process (i.e. Clinically Rated Awareness – CRA) (Reed et al., 1993). In contrast, local (or online) measures of awareness include memory monitoring paradigms such as the FOK task (Hart, 1965). FOK is one of the most sensitive approaches to measuring online memory monitoring processes, and it reflects participants' ability to make predictions regarding the likelihood that they will recognize previously learned information. These paradigms have been used for decades to understand the mechanisms that contribute to self-evaluation in healthy individuals, and are increasingly being implemented to measure specific memory monitoring processes in clinical samples (Cosentino et al., 2007; Cosentino, Metcalfe, Cary, et al., 2011; Rosen, 2011; Rosen et al., 2014; Souchay, Isingrini, & Gil, 2002; Souchay, Isingrini, Pillon, & Gil, 2003). Measuring memory monitoring holds promise for developing a better understanding of the neurocognitive basis of anosognosia, and for further refining models of self-awareness in dementia. Indeed, as noted by Clare, Markova, Roth and Morris (2011), investigation of the links among various forms of self-awareness in AD has begun to increase our understanding of the metacognitive errors that give rise to the clinical symptom of anosognosia, and various efforts have been made toward this end (Cosentino et al., 2016).

From previous studies, we know that within a domain such as memory, anosognosia and monitoring deficits (considered as different levels of awareness) are associated with each other in individuals with AD (Cosentino et al., 2015; Cosentino et al., 2007; Cosentino, Metcalfe, Cary, et al., 2011). In anosognosia for motor deficits and in the healthy aging literatures, examination of online monitoring has spanned different cognitive domains (see Cosentino, Metcalfe, Holmes, et al., 2011; Jenkinson, Preston, & Ellis, 2011). However, to our knowledge, the study of cross-domain monitoring has not yet been undertaken in the context of anosognosia for memory impairment in individuals with AD. Following Cosentino, Metcalfe, Holmes, et al. (2011), we examined memory monitoring and motor monitoring (i.e., metacognitive judgments of agency) as they relate to each other, and to a global, offline measure of memory awareness (CRA) that is traditionally used as diagnostic

criteria for anosognosia or unawareness of memory deficits in AD. Further, commonly associated correlates of each measure were included in each model to assess the extent as to which they share common or distinct correlates.

As it was our main interest to explore different self-evaluation measures within the context of anosognosia, we first determined which participants were classified as aware and unaware of their memory deficits through our clinical measure. We found that over 50% of participants were classified as unaware of their deficits. Variable reports can be found throughout the literature regarding the incidence of unawareness for memory impairment in AD. This variability has been proposed to underlie sampling and methodological differences among studies (see Clare et al., 2005; Cosentino & Stern, 2005 for review).

The conceptualization of the association between local awareness (monitoring) and global awareness (anosognosia)—previously described as different levels of awareness—is represented in two of the most influential models of awareness, the Conscious Awareness Model (CAM) and Clare and colleagues' Hierarchical Model – (Agnew & Morris, 1998; Clare et al., 2011). In the CAM, domain specific monitoring processes are located at a lower level (i.e., *cognitive comparator mechanisms (CCMs)*) (see Figure 1). Supervising each of these domain-specific mechanisms is a central supervisory process, described to function under executive control. The CCMs are specified as those in charge of comparing recent errors in given domains with previous experiences, giving rise to global, higher order self-evaluation of one's abilities. Based on this formulation, dissociation between anosognosia (global awareness) across different domains would be due to a domain-specific comparator (Cm) impairment. On the other hand, a dysfunctional central supervisory system would lead to anosognosia across domains (i.e., *executive anosognosia*) (Agnew & Morris, 1998; McGlynn & Schacter, 1989; Morris & Hannesdottir, 2004; Morris & Mograbi, 2013). That said, this conceptualization has not been experimentally assessed.

In Clare et al.'s (2011) hierarchical model of awareness, ongoing monitoring processes during a task where current errors can be detected are defined as *performance monitoring*. Superior to this level lies the *evaluative judgment* and the *meta-representation* levels, where awareness can be reached through informant interview and in depth clinical interview with the patient. These superior levels of awareness can also be described as global levels of awareness, as they rely on lower levels to produce a stable representation of one self, one that provides the continuity of an individual through time.

The representation of a lower level or local level of awareness, as measured through memory monitoring judgments (i.e., gamma), and its association to a more global level of awareness, as measured through clinical interview (i.e., CRA) is supported in the current and previous studies (Cosentino et al., 2015; Cosentino et al., 2007; Cosentino, Metcalfe, Cary, et al., 2011; Morris et al., 2016). Similarly, there was a link between local levels of awareness across domains, as measured by gamma and the accuracy of the agency judgments for self trials. The main question that we attempted to answer in this paper was the extent to which individuals with anosognosia for memory loss in AD demonstrated deficits at the lower level of awareness (or metacognitive output) in self-monitoring mechanisms beyond memory. This was explored by assessing agency judgments in relation to anosognosia. If monitoring

deficits underlying anosognosia are not domain specific, agency should be distorted in anosognosic patients.

The lack of an observed association between anosognosia and judgments of agency in our study suggests that the mechanisms of awareness in AD are modular, at least to some extent, across the domains of memory and motor functioning. This conclusion, though, requires careful scrutiny. The pattern of performance on the agency task was very similar in both aware and unaware patients, with a clear trend for higher performance on self trials than computer trials. In the latter trials, both aware and unaware participants performed below chance. This pattern is broadly similar to that observed in non-demented older adults (Cioffi, Cocchini, Banissy, & Moore, 2017; Cosentino, Metcalfe, Holmes, et al., 2011) and might speak to a factor unaccounted for in our models that could obscure the association between metacognitive judgments of agency and anosognosia.

Taken together, the current results support the notion that within-domain awareness such as memory, may be associated across levels (i.e., CRA and gamma), but cross-domain monitoring (e.g., motor and memory monitoring) may be associated only within a given level of awareness (i.e., gamma and agency). Though Morris & Mograbi's (2013) CAM parallels Clare and colleagues' (2011) in terms of its hierarchical progress from 'unimodal to heteromodal processes', the CAM does not explicitly address the potential for different levels of metacognitive output (e.g., local contextual judgments of memory and motor performance versus global offline memory awareness). While the CAM does include conscious perception of error through the MAS, which also serves as an 'emergent' process that can represent metacognitive judgments in general, we suggest that it may be useful to conceptualize different levels of metacognitive output separately as the processes and factors associated with each level of output can differ (Clare et al., 2011; Perrotin, Belleville, & Isingrini, 2007). Figure 2 provides an attempt to incorporate our pattern of results into existing models of awareness. As shown in that figure, we propose a simplified model of the different levels of metacognitive output focused on the two domains explored in this paper.

Based on the CAM model and our findings, monitoring of performance depends on domain specific monitors (i.e., CCMs), identified as unconscious processes that can lead to a local metacognitive output of performance (e.g., context local judgment of motor or memory monitoring). At the same time, these monitors are part of the evaluative process by which an individual makes more global and stable judgments of their own abilities. Specific deficits to each CCM would contribute to a domain specific anosognosia. Following the CAM and the motor literature of anosognosia, some individuals may have a more generalized impairment in executive control leading to a generalized impairment of monitoring across domains. In our sample of individuals suffering from AD, we found support for a domain specific CCM deficit (i.e., Cm) contributing to a specific global awareness deficit. The relationship between Cm (memory) and Cn (motor), however, speaks to a shared variance at a local level of awareness.

Finally, our examination of the cognitive and mood correlates of each self-evaluation measure revealed different predictive factors associated with different levels of memory awareness. Specifically, within the cognitive factors, poorer memory performance was a

The association of both levels of awareness, memory and executive function, then, has not been consistent across studies (Correa, Graves, & Costa, 1996; Cosentino et al., 2007; Dalla Barba, Parlato, Iavarone, & Boller, 1995; DeBettignies, Mahurin, & Pirozzolo, 1990; Michon et al., 1994; Reed et al., 1993; Shaked et al., 2014; Starkstein et al., 1996). As mentioned earlier, these contradictory results can be partially explained through the differences between sampling methods and measures used (Clare, 2004; Cosentino & Stern, 2005). Another possibility is that impairment in memory or executive function alone is not sufficient to cause anosognosia or memory monitoring deficits, as other process likely contribute to self-reflection. It has also been suggested that cognitive and metacognitive processes may simply be concomitant deficits affected in the neurodegenerative process. For example, previous work by Shaked et al. (2014), showed that an index of non-verbal memory and non-verbal executive functions was more closely related to memory monitoring (i.e., gamma) than was a verbal index of these cognitive domains. These results were interpreted within a neuroanatomic framework as potentially pointing to differential disruption in right hemisphere networks critical for processing nonverbal information as well as for supporting self-reflective processes (Cosentino, 2014).

Lastly, in regards to mood, our results showed different relations between depression and our three measures of self-evaluation. Specifically, we found that endorsing higher levels of depression was associated with being globally more aware of one's memory deficits, consistent with a number of previous studies (Bertrand et al., 2016; Cines et al., 2015; Conde-Sala et al., 2014), but not with memory monitoring. These findings support a previously shown dissociation of the correlates found between these measures of self-evaluation. For example, Cosentino, Metcalfe, Cary, et al. (2011) found that decision making capacity pertaining to medication management was related to global memory awareness but not local (gamma). The authors suggested that global awareness likely reflects a general, context independent higher level of awareness. On the other hand, memory monitoring, as measured by gamma, is a specific and local, context dependent lower level of awareness. This argument might explain the observed dissociation in our study between mood and the two measures of self-awareness. As suggested by Clare and colleagues, and consistent with the Hierarchical model, global measures of awareness may in part reflect beliefs, premorbid factors, or psychological functioning as they are not constrained by the same type of specific contextual details that constrain self-evaluative judgments in the context of local awareness measures (Clare et al., 2012). As such, local metacognitive evaluations might reflect a more 'objective' or 'pure' measure of someone's self-evaluative ability.

To conclude, while global unawareness in AD (i.e., anosognosia) seems to break down in tandem with deficits in mnemonic self-monitoring (i.e., gamma), anosognosia for memory loss was not associated with local self-monitoring processes in the motor domain as measured by metacognition of agency in the data presented here. However, the local or online forms of self-evaluation that are at play during agency judgments, appear to relate to those used to make judgments about mnemonic self-monitoring. That is, it seems that within-domain awareness may be associated across levels, but cross-domain monitoring may be associated only within their level of awareness (i.e., local awareness). Future research should examine other types of monitoring processes as well as anosognosia for other deficits, to evaluate the extent and qualitative differences of self-monitoring at different

levels of awareness for memory and other deficits. Due to the important clinical implications of deficits in awareness among individuals suffering from AD, it is essential that we continue to refine our current theoretical models of self-awareness. Doing so will open the door to shaping specific rehabilitation programs in order to minimize the negative consequences of unawareness.

Acknowledgments

Statement of funding

This work was supported by a Paul B. Beeson Career Development in Aging Award (1 K23 AG032899) funded jointly by the National Institute on Aging and the American Federation of Aging Research.

References

- Agnew SK, Morris RG. The heterogeneity of anosognosia for memory impairment in Alzheimer's disease: A review of the literature and a proposed model. *Aging & Mental Health*. 1998; 2(1):7–19. DOI: 10.1080/13607869856876
- Ansell EL, Bucks RS. Mnemonic anosognosia in Alzheimer's disease: a test of Agnew and Morris (1998). *Neuropsychologia*. 2006; 44(7):1095–1102. DOI: 10.1016/j.neuropsychologia.2005.10.019 [PubMed: 16324727]
- Berti, A., Spinazzola, L., Pia, L., Rabuffetti, M. Motor awareness and motor intention in anosognosia for hemiplegia. In: Haggard, P., Rossetti, Y., Kawato, M., editors. *Sensorimotor foundations of higher cognition series: Attention and performance number XXII*. New York, NY: Oxford University Press; 2007.
- Bertrand E, Dourado MC, Laks J, Morris RG, Landeira-Fernandez J, Mograbi DC. Mood-congruent recollection and anosognosia in Alzheimer's disease. *Cortex*. 2016; 84:55–62. DOI: 10.1016/j.cortex.2016.09.001 [PubMed: 27697664]
- Blakemore SJ, Wolpert DM, Frith CD. Abnormalities in the awareness of action. *Trends in Cognitive Sciences*. 2002; 6(6):237–242. [PubMed: 12039604]
- Cines S, Farrell M, Steffener J, Sullo L, Huey E, Karlawish J, Cosentino S. Examining the Pathways Between Self-Awareness and Well-Being in Mild to Moderate Alzheimer's Disease. *American Journal of Geriatric Psychiatry*. 2015; 23(12):1297–1306. DOI: 10.1016/j.jagp.2015.05.005 [PubMed: 26560509]
- Cioffi MC, Cocchini G, Banissy MJ, Moore JW. Ageing and agency: age-related changes in susceptibility to illusory experiences of control. *Royal Society Open Science*. 2017; 4(5): 161065. doi: 10.1098/rsos.161065 [PubMed: 28572999]
- Clare L. Awareness in early-stage Alzheimer's disease: a review of methods and evidence. *British Journal of Clinical Psychology*. 2004; 43(Pt 2):177–196. DOI: 10.1348/014466504323088042 [PubMed: 15169617]
- Clare L, Marcová IS, Roth I, Morris RG. Awareness in Alzheimer's disease and associated dementias: Theoretical framework and clinical implications. *Aging & Mental Health*. 2011; 15(8):936–944. DOI: 10.1080/13607863.2011.583630 [PubMed: 21702711]
- Clare L, Marcová IS, Verhey F, Kenny G. Awareness in dementia: A review of assessment methods and measures. *Aging & Mental Health*. 2005; 9(5):394–413. [PubMed: 16024399]
- Clare L, Nelis SM, Martyr A, Roberts J, Whitaker CJ, Marcová IS, ... Morris RG. The influence of psychological, social and contextual factors on the expression and measurement of awareness in early-stage dementia: testing a biopsychosocial model. *International Journal of Geriatric Psychiatry*. 2012; 27(2):167–177. DOI: 10.1002/gps.2705 [PubMed: 21425345]
- Clare L, Wilson BA, Carter G, Roth I, Hodges JR. Assessing awareness in early-stage Alzheimer's disease: Development and piloting of the Memory Awareness Rating Scale. *Neuropsychological Rehabilitation*. 2002; 12(4):341–362.

- Clare L, Wilson BA, Carter G, Roth I, Hodges JR. Awareness in early stage Alzheimer's disease: Relationship to Outcome of Cognitive Rehabilitation. *Journal of Clinical and Experimental Neuropsychology*. 2004; 26(2):215–226. DOI: 10.1076/jcen.26.2.215.28088 [PubMed: 15202541]
- Cocchini G, Beschin N, Fotopoulou A, Della Sala S. Explicit and implicit anosognosia or upper limb motor impairment. *Neuropsychologia*. 2010; 48(5):1489–1494. DOI: 10.1016/j.neuropsychologia.2010.01.019 [PubMed: 20117119]
- Cocchini G, Beschin N, Sala SD. Chronic anosognosia: a case report and theoretical account. *Neuropsychologia*. 2002; 40(12):2030–2038. [PubMed: 12208000]
- Conde-Sala JL, Rene-Ramirez R, Turro-Garriga O, Gascon-Bayarri J, Campdelacreu-Fumado J, Juncadella-Puig M, ... Garre-Olmo J. Severity of dementia, anosognosia, and depression in relation to the quality of life of patients with Alzheimer's disease: discrepancies between patients and caregivers. *American Journal of Geriatric Psychiatry*. 2014; 22(2):138–147. DOI: 10.1016/j.jagp.2012.07.001 [PubMed: 23567444]
- Correa DD, Graves RE, Costa L. Awareness of memory deficit in Alzheimer's disease patients and memory-impaired older adults. *Aging, Neuropsychology, and Cognition*. 1996; 3(3):215–228. DOI: 10.1080/13825589608256625
- Cosentino S. Metacognition in Alzheimer's Disease. In: Fleming, SM., Frith, C., editors. *The cognitive neuroscience of metacognition*. New York, NY: Springer; 2014. p. 389–407.
- Cosentino S, Brickman AM, Griffith E, Habeck C, Cines S, Farrell M, ... Stern Y. The right insula contributes to memory awareness in cognitively diverse older adults. *Neuropsychologia*. 2015; 75:163–169. DOI: 10.1016/j.neuropsychologia.2015.05.032 [PubMed: 26049091]
- Cosentino S, Metcalfe J, Butterfield B, Stern Y. Objective metamemory testing captures awareness of deficit in Alzheimer's disease. *Cortex*. 2007; 43(7):1004–1019. [PubMed: 17941356]
- Cosentino S, Metcalfe J, Cary MS, De Leon J, Karlawish J. Memory Awareness Influences Everyday Decision Making Capacity about Medication Management in Alzheimer's Disease. *International Journal of Alzheimer's Disease*. 2011; 2011:9. doi: 10.4061/2011/483897
- Cosentino S, Metcalfe J, Holmes B, Steffener J, Stern Y. Finding the self in metacognitive evaluations: Metamemory and agency in nondemented elders. *Neuropsychology*. 2011; 25(5):602–612. DOI: 10.1037/a0023972 [PubMed: 21728426]
- Cosentino S, Stern Y. Metacognitive theory and assessment in dementia: do we recognize our areas of weakness? *Journal of the International Neuropsychological Society*. 2005; 11(7):910–919. [PubMed: 16519270]
- Cosentino S, Zhu C, Bertrand E, Metcalfe J, Janicki S, Cines S. Examination of the metacognitive errors that contribute to anosognosia in Alzheimer's disease. *Cortex*. 2016; 84:101–110. DOI: 10.1016/j.cortex.2016.08.003 [PubMed: 27750070]
- Cotrell V, Wild K. Longitudinal Study of Self-imposed Driving Restrictions and Deficit Awareness in Patients with Alzheimer's Disease. *Alzheimer's Disease and Associated Disorders*. 1999; 13(3): 151–156.
- Dalla Barba G, Parlato V, Iavarone A, Boller F. Anosognosia, intrusions and 'frontal' functions in Alzheimer's disease and depression. *Neuropsychologia*. 1995; 33(2):247–259. [PubMed: 7746367]
- Davies M, Davies AA, Coltheart M. Anosognosia and the Two-factor Theory of Delusions. *Mind & Language*. 2005; 20(2):209–236. DOI: 10.1111/j.0268-1064.2005.00283.x
- DeBettignies BH, Mahurin RK, Pirozzolo FJ. Insight for impairment in independent living skills in Alzheimer's disease and multi-infarct dementia. *Journal of Clinical and Experimental Neuropsychology*. 1990; 12(2):355–363. [PubMed: 2341561]
- Derouesne C, Thibault S, Lagha-Pierucci S, Baudouin-Madec V, Ancrì D, Lacomblez L. Decreased awareness of cognitive deficits in patients with mild dementia of the Alzheimer type. *International Journal of Geriatric Psychiatry*. 1999; 14(12):1019–1030. [PubMed: 10607969]
- Feinberg I. Efference copy and corollary discharge: implications for thinking and its disorders. *Schizophrenia Bulletin*. 1978; 4(4):636–640. [PubMed: 734369]
- Feinberg TE, Roane DM, Kwan PC, Schindler RJ, Haber LD. Anosognosia and visuo-verbal confabulation. *Archives of Neurology*. 1994; 51(5):468–473. [PubMed: 8179496]

- Fernandez-Duque D, Baird JA, Posner MI. Executive Attention and Metacognitive Regulation. *Consciousness and Cognition*. 2000; 9(2):288–307. DOI: 10.1006/ccog.2000.0447 [PubMed: 10924249]
- Folstein MF, Folstein SE, McHugh PR. “Mini-mental state”. A practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research*. 1975; 12(3):189–198. [PubMed: 1202204]
- Fotopoulou A. Time to get rid of the ‘Modular’ in neuropsychology: A unified theory of anosognosia as aberrant predictive coding. *Journal of Neuropsychology*. 2014; 8(1):1–19. DOI: 10.1111/jnp.12010 [PubMed: 23469983]
- Fotopoulou A, Tsakiris M, Haggard P, Vagopoulou A, Rudd A, Kopelman M. The role of motor intention in motor awareness: an experimental study on anosognosia for hemiplegia. *Brain*. 2008; 131(Pt 12):3432–3442. DOI: 10.1093/brain/awn225 [PubMed: 18812442]
- Frith CD. The self in action: Lessons from delusions of control. *Consciousness and Cognition*. 2005; 14(4):752–770. DOI: 10.1016/j.concog.2005.04.002 [PubMed: 16098765]
- Frith CD, Blakemore SJ, Wolpert DM. Explaining the symptoms of schizophrenia: Abnormalities in the awareness of action. *Brain Research Reviews*. 2000; 31(2–3):357–363. DOI: 10.1016/S0165-0173(99)00052-1 [PubMed: 10719163]
- Gallagher S. Philosophical conceptions of the self: implications for cognitive science. *Trends in Cognitive Sciences*. 2000; 4(1):14–21. DOI: 10.1016/S1364-6613(99)01417-5 [PubMed: 10637618]
- Glosser G, Goodglass H. Disorders in executive control functions among aphasic and other brain-damaged patients. *Journal of Clinical and Experimental Neuropsychology*. 1990; 12(4):485–501. DOI: 10.1080/01688639008400995 [PubMed: 1698809]
- Glosser G, Goodglass H, Biber C. Assessing visual memory disorders. *Psychological Assessment: A Journal of Consulting and Clinical Psychology*. 1989; 1(2):82–91. DOI: 10.1037/1040-3590.1.2.82
- Haggard P, Tsakiris M. The Experience of Agency. *Current Directions in Psychological Science*. 2009; 18(4):242–246. DOI: 10.1111/j.1467-8721.2009.01644.x
- Hertzog C, Dunlosky J, Sinclair SM. Episodic feeling-of-knowing resolution derives from the quality of original encoding. *Memory and Cognition*. 2010; 38(6):771–784. DOI: 10.3758/MC.38.6.771 [PubMed: 20852240]
- Jenkinson PM, Edelstyn NM, Drakeford JL, Ellis SJ. Reality monitoring in anosognosia for hemiplegia. *Consciousness and Cognition*. 2009; 18(2):458–470. DOI: 10.1016/j.concog.2008.12.005 [PubMed: 19195910]
- Jenkinson PM, Fotopoulou A. Motor awareness in anosognosia for hemiplegia: experiments at last! *Experimental Brain Research*. 2010; 204(3):295–304. DOI: 10.1007/s00221-009-1929-8 [PubMed: 19593552]
- Jenkinson PM, Preston C, Ellis SJ. Unawareness after stroke: A review and practical guide to understanding, assessing, and managing anosognosia for hemiplegia. *Journal of Clinical and Experimental Neuropsychology*. 2011; 33(10):1079–1093. DOI: 10.1080/13803395.2011.596822 [PubMed: 21936643]
- Kelleher M, Tolea MI, Galvin JE. Anosognosia increases caregiver burden in mild cognitive impairment. *International Journal of Geriatric Psychiatry*. 2016; 31(7):799–808. DOI: 10.1002/gps.4394 [PubMed: 26643996]
- Kirkpatrick MG, Metcalfe J, Greene MJ, Hart CL. Effects of intranasal methamphetamine on metacognition of agency. *Psychopharmacology*. 2008; 197:137–144. DOI: 10.1007/s00213-007-1018-2 [PubMed: 18043908]
- Koltai DC, Welsh-Bohmer KA, Schmechel DE. Influence of anosognosia on treatment outcome among dementia patients. *Neuropsychological Rehabilitation*. 2001; 11(3–4):455–475. DOI: 10.1080/09602010042000097
- Koriat A. How do we know that we know? The accessibility model of the feeling of knowing. *Psychological Review*. 1993; 100(4):609–639. [PubMed: 8255951]
- Koriat A. The feeling of knowing: Some metatheoretical implications for consciousness and control. *Consciousness and Cognition*. 2000; 9(2):149–171. DOI: 10.1006/ccog.2000.0433 [PubMed: 10924234]

- Koriat A, Levy-Sadot R. The combined contributions of the cue-familiarity and accessibility heuristics to feelings of knowing. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 2001; 27(1):34–53.
- Levine DN. Unawareness of visual and sensorimotor defects: a hypothesis. *Brain Cogn*. 1990; 13(2): 233–281. [PubMed: 2390236]
- Lopez OL, Becker JT, Somsak D, Dew MA, DeKosky ST. Awareness of cognitive deficits and anosognosia in probable Alzheimer's disease. *European Neurology*. 1994; 34(5):277–282. [PubMed: 7995303]
- Marcel AJ, Tegnér R, Nimmo-Smith I. Anosognosia for Plegia: Specificity, Extension, Partiality and Disunity of Bodily Unawareness. *Cortex*. 2004; 40(1):19–40. DOI: 10.1016/S0010-9452(08)70919-5 [PubMed: 15070001]
- McGlynn SM, Schacter DL. Unawareness of deficits in neuropsychological syndromes. *Journal of Clinical and Experimental Neuropsychology*. 1989; 11(2):143–205. [PubMed: 2647781]
- Metcalfe J. Feelings and judgments of knowing: Is there a special noetic state? *Consciousness and Cognition*. 2000; 9(2):178–186. DOI: 10.1006/ccog.2000.0451 [PubMed: 10924236]
- Metcalfe J, Eich TS, Castel AD. Metacognition of agency across the lifespan. *Cognition*. 2010; 116(2): 267–282. DOI: 10.1016/j.cognition.2010.05.009 [PubMed: 20570251]
- Metcalfe J, Greene MJ. Metacognition of agency. *Journal of Experimental Psychology: General*. 2007; 136(2):184–199. DOI: 10.1037/0096-3445.136.2.184 [PubMed: 17500645]
- Metcalfe J, Schwartz BL, Joaquim SG. The cue-familiarity heuristic in metacognition. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 1993; 19(4):851–861. DOI: 10.1037/0278-7393.19.4.851
- Metcalfe, J., Van Snellenberg, JX., DeRosse, P., Balsam, P., Malhotra, AK. Judgments of agency in schizophrenia: An impairment in auto-noetic metacognition. In: Fleming, SM., Frith, C., editors. *The cognitive neuroscience of metacognition*. New York, NY: Springer; 2014. p. 367-387.
- Michon A, Deweer B, Pillon B, Agid Y, Dubois B. Relation of anosognosia to frontal lobe dysfunction in Alzheimer's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*. 1994; 57(7):805–809.
- Michotte, A. *The perception of causality*. New York, NY: Basic books; 1963.
- Mograbi DC, Brown RG, Morris RG. Anosognosia in Alzheimer's disease – The petrified self. *Consciousness and Cognition*. 2009; 18(4):989–1003. DOI: 10.1016/j.concog.2009.07.005 [PubMed: 19683461]
- Mograbi DC, Morris RG. Implicit awareness in anosognosia: Clinical observations, experimental evidence, and theoretical implications. *Cognitive Neuroscience*. 2013; 4(3–4):181–197. DOI: 10.1080/17588928.2013.833899 [PubMed: 24251606]
- Moore JW. What Is the Sense of Agency and Why Does it Matter? *Frontiers in Psychology*. 2016; 7:1272.doi: 10.3389/fpsyg.2016.01272 [PubMed: 27621713]
- Moro V, Pernigo S, Zapparoli P, Cordioli Z, Aglioti SM. Phenomenology and neural correlates of implicit and emergent motor awareness in patients with anosognosia for hemiplegia. *Behavioural Brain Research*. 2011; 225(1):259–269. DOI: 10.1016/j.bbr.2011.07.010 [PubMed: 21777624]
- Morris, RG., Hannesdottir, K. Loss of 'awareness' in Alzheimer's disease. In: Morris, RG., Becker, JT., editors. *The Cognitive Neuropsychology of Alzheimer's Disease*. Oxford, England: Oxford University Press; 2004. p. 275-296.
- Morris RG, Mograbi DC. Anosognosia, autobiographical memory and self knowledge in Alzheimer's disease. *Cortex*. 2013; 49(6):1553–1565. DOI: 10.1016/j.cortex.2012.09.006 [PubMed: 23117055]
- Morris RG, Nelis SM, Martyr A, Marcová IS, Roth I, Woods RT, ... Clare L. Awareness of memory task impairment versus everyday memory difficulties in dementia. *Journal of Neuropsychology*. 2016; 10(1):130–142. DOI: 10.1111/jnp.12062 [PubMed: 25488044]
- Perrotin A, Belleville S, Isingrini M. Metamemory monitoring in mild cognitive impairment: Evidence of a less accurate episodic feeling-of-knowing. *Neuropsychologia*. 2007; 45(12):2811–2826. DOI: 10.1016/j.neuropsychologia.2007.05.003 [PubMed: 17597165]
- Perrotin A, Isingrini M, Souchay C, Clarys D, Taconnat L. Episodic feeling-of-knowing accuracy and cued recall in the elderly: evidence for double dissociation involving executive functioning and processing speed. *Acta Psychologica*. 2006; 122(1):58–73. DOI: 10.1016/j.actpsy.2005.10.003 [PubMed: 16309619]

- Price CC, Garrett KD, Jefferson AL, Cosentino S, Tanner JJ, Penney DL, ... Libon DJ. Leukoaraiosis severity and list-learning in dementia. *Clinical Neuropsychologist*. 2009; 23(6):944–961. DOI: 10.1080/13854040802681664 [PubMed: 19370451]
- Prigatano GP. Disturbances of self-awareness and rehabilitation of patients with traumatic brain injury: a 20-year perspective. *Journal of Head Trauma Rehabilitation*. 2005; 20(1):19–29. [PubMed: 15668568]
- Reder LM, Ritter FE. What determines initial feeling of knowing? Familiarity with question terms, not with the answer. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 1992; 18(3):435–451. DOI: 10.1037/0278-7393.18.3.435
- Reed BR, Jagust WJ, Coulter L. Anosognosia in Alzheimer's disease: Relationships to depression, cognitive function, and cerebral perfusion. *Journal of Clinical and Experimental Neuropsychology*. 1993; 15(2):231–244. DOI: 10.1080/01688639308402560 [PubMed: 8491848]
- Rosen HJ. Anosognosia in neurodegenerative disease. *Neurocase*. 2011; 17(3):231–241. DOI: 10.1080/13554794.2010.522588 [PubMed: 21667396]
- Rosen HJ, Alcantar O, Zakrzewski J, Shimamura AP, Neuhaus J, Miller BL. Metacognition in the behavioral variant of frontotemporal dementia and Alzheimer's disease. *Neuropsychology*. 2014; 28(3):436–447. DOI: 10.1037/neu0000012 [PubMed: 24548124]
- Rymer S, Salloway S, Norton L, Malloy P, Correia S, Monast D. Impaired awareness, behavior disturbance, and caregiver burden in Alzheimer's disease. *Alzheimer's Disease and Associated Disorders*. 2002; 16(4):248–253.
- Saj A, Vocat R, Vuilleumier P. Action-monitoring impairment in anosognosia for hemiplegia. *Cortex*. 2014; 61:93–106. DOI: 10.1016/j.cortex.2014.10.017 [PubMed: 25481468]
- Schacter DL. Toward a cognitive neuropsychology of awareness: implicit knowledge and anosognosia. *Journal of Clinical and Experimental Neuropsychology*. 1990; 12(1):155–178. DOI: 10.1080/01688639008400962 [PubMed: 2406281]
- Shlottman A, Shanks DR. Evidence of a distinction between judge and perceived causality. *Quarterly Journal of Experimental Psychology*. 1992; 44:321–342. DOI: 10.1080/02724989243000055 [PubMed: 1565803]
- Schnyer DM, Verfaellie M, Alexander MP, LaFleche G, Nicholls L, Kaszniak AW. A role for right medial prefrontal cortex in accurate feeling-of-knowing judgments: evidence from patients with lesions to frontal cortex. *Neuropsychologia*. 2004; 42(7):957–966. DOI: 10.1016/j.neuropsychologia.2003.11.020 [PubMed: 14998710]
- Schwartz BL, Metcalfe J. Cue familiarity but not target retrievability enhances feeling-of-knowing judgments. *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 1992; 18(5):1074–1083.
- Seltzer B, Vasterling JJ, Yoder JA, Thompson KA. Awareness of deficit in Alzheimer's disease: relation to caregiver burden. *Gerontologist*. 1997; 37(1):20–24. [PubMed: 9046701]
- Shaked D, Farrell M, Huey E, Metcalfe J, Cines S, Karlawish J, ... Cosentino S. Cognitive Correlates of Metamemory in Alzheimer's Disease. *Neuropsychology*. 2014; 28(5):695–705. DOI: 10.1037/neu0000078 [PubMed: 24819066]
- Shimamura AP. Memory and frontal lobe function. In: Gazzaniga, MS., editor. *The cognitive neurosciences*. Cambridge, MA: MIT Press; 1995.
- Shimamura AP. Toward a cognitive neuroscience of metacognition. *Consciousness and Cognition*. 2000; 9(2 Pt 1):313–323. discussion 324–316. DOI: 10.1006/ccog.2000.0450 [PubMed: 10924251]
- Souchay C, Isingrini M, Espagnet L. Aging, episodic memory feeling-of-knowing, and frontal functioning. *Neuropsychology*. 2000; 14(2):299–309. DOI: 10.1037/0894-4105.14.2.299 [PubMed: 10791869]
- Souchay C, Isingrini M, Gil R. Alzheimer's disease and feeling-of-knowing in episodic memory. *Neuropsychologia*. 2002; 40(13):2386–2396. DOI: 10.1016/S0028-3932(02)00075-1 [PubMed: 12417467]
- Souchay C, Isingrini M, Pillon B, Gil R. Metamemory accuracy in Alzheimer's disease and frontotemporal lobe dementia. *Neurocase*. 2003; 9(6):482–492. DOI: 10.1076/neur.9.6.482.29376 [PubMed: 16210230]

- Starkstein SE, Fedoroff JP, Price TR, Leiguarda R, Robinson RG. Anosognosia in patients with cerebrovascular lesions. A study of causative factors. *Stroke*. 1992; 23(10):1446–1453. [PubMed: 1412582]
- Starkstein SE, Sabe L, Chemerinski E, Jason L, Leiguarda R. Two domains of anosognosia in Alzheimer's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*. 1996; 61(5):485–490.
- Starkstein SE, Vazquez S, Migliorelli R, Teson A, Sabe L, Leiguarda R. A single-photon emission computed tomographic study of anosognosia in Alzheimer's disease. *Archives of Neurology*. 1995; 52(4):415–420. [PubMed: 7710378]
- Stuss, DT., Benson, DF. *The frontal lobes*. New York: Raven Press; 1986.
- Synofzik M, Vosgerau G, Newen A. Beyond the comparator model: A multifactorial two-step account of agency. *Consciousness and Cognition*. 2008; 17(1):219–239. DOI: 10.1016/j.concog.2007.03.010 [PubMed: 17482480]
- Venneri A, Shanks MF. Belief and awareness: reflections on a case of persistent anosognosia. *Neuropsychologia*. 2004; 42(2):230–238. DOI: 10.1016/S0028-3932(03)00171-4 [PubMed: 14644108]
- Vocat R, Saj A, Vuilleumier P. The riddle of anosognosia: does unawareness of hemiplegia involve a failure to update beliefs? *Cortex*. 2013; 49(7):1771–1781. DOI: 10.1016/j.cortex.2012.10.009 [PubMed: 23290635]
- Vuilleumier P. Anosognosia: the neurology of beliefs and uncertainties. *Cortex*. 2004; 40(1):9–17. [PubMed: 15070000]
- Wechsler, DA. *Wechsler Adult Intelligence Scale - Revised*. San Antonio, TX: The Psychological Corporation; 1987.
- Wegner, DM. *The Illusion of Conscious Will*. Cambridge, MA: MIT Press; 2002.
- Wegner DM, Wheatley T. Apparent mental causation. Sources of the experience of will. *American Psychologist*. 1999; 54(7):480–492. [PubMed: 10424155]
- Wild K, Cotrell V. Identifying Driving Impairment in Alzheimer's Disease: A Comparison of Self and Observer Reports Versus Driving Evaluation. *Alzheimer's Disease and Associated Disorders*. 2003; 17(1):27–34.
- Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, Leirer VO. Development and validation of a geriatric depression screening scale: a preliminary report. *Journal of Psychiatric Research*. 1982; 17(1):37–49. [PubMed: 7183759]

Highlights

- Hierarchical models of awareness for memory loss are assessed in Alzheimer's disease
- Self-monitoring in motor and memory domains are associated in Alzheimer' disease.
- Self-monitoring processes dissociate in anosognosia for memory loss.




Figure 1. Modified version of the revised CAM model from Morris and Mograbi (2013). Comparator mechanisms proposed to underlie monitoring of different cognitive domains highlighted.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2.

Metacognitive output of global and local levels of awareness in both memory and motor domains. Solid lines represent relationships found in this current study. Dotted arrows represent relationships previously shown in anosognosia for hemiplegia but not assessed in this study (see Jenkinson & Fotopoulou, 2010; Saj et al., 2014; Vocat et al., 2013). CRA – Clinically rated awareness; SRD – Subjective rating discrepancy; Cm – memory comparator; Cn – motor comparator.

Table 1

Mean and standard deviations of demographic and neuropsychological variables in participants unaware and aware of their memory difficulties.

<i>Demographics details, mood and neuropsychological performance</i>	Unaware (n=20)	Aware (n=15)	Sig. Two tailed	95 % Confidence intervals
Age	79.94 (8.02)	74.78 (10.54)	.10	- 1.21, 11.54
Education	16.00 (2.73)	16.73 (3.10)	.46	-1.55, 11.87
Gender (female/male)	14/6	10/5	.94	-
Race (Caucasian/African American)	19/1	13/2	.38	-
MMSE (0–30)	25.05 (1.93)	25.07 (2.18)	.98	-1.43, 1.40
Memory index (Z score)	-.20 (.58)	.19 (.79)	.10	-.86, .07
Executive Index (Z score)	.01 (.89)	-.01 (.80)	.69	-.28, .41
Attention Index (Z score)	.11 (1.03)	-.15 (1.02)	.48	-.49, 1.00
Mood (0–30) *	3.00 (6.75)	7.00 (9.00)	.06	-

Higher scores on MSMSE reflect better performance. Higher scores in the Mood variable reflect higher endorsement of depressive items.

* Non normal data is reported as median and interquartile ranges.

Table 2

Mean and standard deviations of metacognitive measures for memory and agency in participants unaware and aware of their memory deficits.

<i>Metacognitive measures of memory and motor domains (range)</i>	Unaware (n=20)	Aware (n=15)	Sig. Two tailed	95% Confidence intervals
Gamma (-1-1)	.18(.34)	.50(.26)	.005	-.53, -.10
Global calibration (-1-1)	.07(.18)	.09(.18)	.72	-.15, .10
Item level calibration (-1-1)	.01(.12)	.02(.08)	.78	-.08, .06
Agency total (0-16)	8.30(2.61)	8.60(3.04)	.76	-2.25, 1.65
Agency computer trials (0-8)	2.40(2.11)	2.40(2.77)	1.00	-1.67, 1.67
Agency self trials (0-8)	5.90(1.74)	6.20(1.65)	.61	-1.48, .88

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Regression models of self-awareness measures of memory monitoring (gamma), anosognosia (CRA), and the accuracy of agency judgments in self trials and in computer trials.

<i>Predictors of memory monitoring (gamma), CRA and agency</i>	Gamma (Std. error)	CRA (Std. error)	Agency self trials (Std. error)	Agency computer trials B (Std. error)
Gamma	-	3.77 (1.79)	1.47 (1.47)	2.10 (1.87)
CRA	.23 (.10)	-	.23 (.82)	-.81 (1.05)
Agency self	.08 (.03)	-.15 (.32)	-	-
Executive functions	-.02 (.09)	-.14 (.87)	-.01 (.77)	.69 (.97)
Memory	.25 (.07)	.12 (.88)	-	-
Mood	-.01 (.01)	.21 (.10)	-	-
Computer experience	-	-	-.03 (.13)	.24 (.16)

Unstandardized betas and standard errors of the individual predictors are included. Significant predictors are shown in bold ($p < .05$).